



Changing trends of ocular trauma in the time of COVID-19 pandemic

Marco Pellegrini¹ · Matilde Roda¹ · Natalie Di Geronimo¹ · Enrico Lupardi¹ · Giuseppe Giannaccare² · Costantino Schiavi¹

Received: 11 April 2020 / Revised: 23 April 2020 / Accepted: 24 April 2020 / Published online: 6 May 2020
© The Royal College of Ophthalmologists 2020

To the Editor:

To reduce the spread of the novel coronavirus (2019-nCoV), countries have promoted a range of unprecedented public health responses. These measures aim at reducing the final size of the epidemic as well as its peak in order to decrease the acute pressure on the health-care system [1]. In Italy, the government ordered people to stay home, restricting movements with the exception of work, urgent matters and health reasons. Furthermore, all commercial and productive activities, except those providing essential services, were obligated to remain closed [2].

Ocular trauma represents a serious public health problem and leading cause of visual impairment [3, 4]. The COVID-19 social distancing measures might have a significant impact on the risk of ocular trauma. In this study, we retrospectively reviewed the charts of all patients presenting to an Italian ophthalmological emergency department (the Ophthalmology Unit of the S.Orsola-Malpighi University Hospital in Bologna) to identify all eye injuries. Data were analysed from 10th March 2020 (i.e. the day in which the quarantine measures were applied in our city) to 10th April 2020, and confronted with those of the same period of the previous year (from 10th March 2019 to 10th April 2019).

In the 2019 study period, there were 354 eye injuries (15.6% of all patients presenting to the emergency department). In the 2020 study period, eye injuries decreased to 112 (19.9% of all patients). The characteristics of eye injuries in the two study periods are reported in Table 1. During quarantine, the proportion of children and

Table 1 Characteristics of eye injuries in the 2019 study period (from 10th March 2019 to 10th April, 2019) and 2020 study period (from 10th March 2020 to 10th April 2020).

Characteristic	2019 period	2020 period
Total number of eye injuries	354	112
Sex (m/f)	236/109	84/28
Mean age (\pm SD)	40.7 \pm 19.7	43.3 \pm 17.4
Mechanism of injury (% of the total)		
Sports	21 (5.9%)	3 (2.7%)
Manual works ^a	49 (13.8%)	18 (16.1%)
Animal care	8 (2.3%)	4 (3.6%)
Gardening/injuries with plants	30 (8.5%)	12 (10.7%)
Home activities	44 (12.4%)	19 (17.0%)
Falls	23 (6.5%)	1 (0.9%)
Burns/corrosive substances	26 (7.3%)	8 (7.1%)
Violence	15 (4.2%)	6 (5.4%)
Other/unknown	138 (39.0%)	41 (36.6%)
Diagnosis (% of the total)		
None	33 (9.3%)	1 (0.9%)
Foreign body on external eye	142 (40.1%)	56 (50.0%)
Superficial injury (cornea and conjunctiva)	97 (27.4%)	39 (34.8%)
Subconjunctival haemorrhage	23 (6.5%)	6 (5.4%)
Eyelid injury	33 (9.3%)	3 (2.7%)
Posterior vitreous detachment	2 (0.6%)	1 (0.9%)
Hyphema	4 (1.1%)	4 (3.6%)
Retinal oedema	15 (4.2%)	1 (0.9%)
Vitreous haemorrhage	1 (0.3%)	0 (0.0%)
Orbital fracture	3 (0.8%)	0 (0.0%)
Penetrating wound	1 (0.3%)	1 (0.9%)

^aGrinding, cutting, sanding, drilling, hammering, sawing.

✉ Marco Pellegrini
marco.pellegrini@hotmail.it

¹ Ophthalmology Unit, S.Orsola-Malpighi University Hospital, University of Bologna, Bologna, Italy

² Department of Ophthalmology, University of “Magna Graecia”, Catanzaro, Italy

adolescents with eye injuries decreased (from 14.7% to 8.0%, Fig. 1a), while the proportion of males increased (from 66.7% to 75.0%, Fig. 1b). Regarding the mechanisms of injury, the percentage of falls and sport injuries had the




Fig. 1 Number of eye injuries in the 2019 study period (from 10th March 2019 to 10th April, 2019) and 2020 study period (from 10th March 2020 to 10th April 2020). (a) Eye injuries categorized by age.

(b) Eye injuries categorized by sex. (c) Eye injuries categorized by mechanism of trauma. (d) Eye injuries categorized as minor or major injuries.

highest decrease (respectively, from 6.5% to 0.9% and from 5.9% to 2.7%), while injuries during home activities and injuries with plants had the highest increase (respectively, from 12.4% to 17.0% and from 8.5% to 10.7%, Fig. 1c). The percentage of minor injuries with low risk of vision loss increased (from 93.2% to 94.6%), while major injuries requiring monitoring decreased (from 6.8% to 5.4%, Fig. 1d).

There was a striking 68.4% decrease in the number of eye injuries seen in our Unit during the last month. Behavioural changes during the quarantine could be associated with lower risk of trauma. The decreases of sport injuries and of injuries in children during school closure seem to support this hypothesis. However, the drop of patients seeking emergency care affected all injuries, including serious ones potentially associated with vision loss. We believe that some patients may intentionally avoid urgent care rather than risking coronavirus exposure at hospitals. Anecdotal reports suggest that this is also happening for life-threatening medical emergencies such as myocardial infarction and stroke [5, 6]. Since ocular trauma is a major cause of vision loss, the importance of not delaying or

avoiding treatment should be stressed to all patients to prevent ocular morbidities.

Compliance with ethical standards

Conflict of interest The authors declare that they have no conflict of interest.

Publisher’s note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

References

1. Wilder-Smith A, Chiew CJ, Lee VJ. Can we contain the COVID-19 outbreak with the same measures as for SARS?. *Lancet Infect Dis.* 2020. [https://doi.org/10.1016/S1473-3099\(20\)30129-8](https://doi.org/10.1016/S1473-3099(20)30129-8).
2. Italian Government. decreto del presidente del Consiglio dei ministri (DPCM) March 11, 2020. <http://www.governo.it/it/articolo/coronavirus-conte-firma-il-dpcm-11-marzo-2020/14299>.
3. Matsa E, Shi J, Wheeler KK, McCarthy T, McGregor ML, Leonard JC. Trends in US emergency department visits for pediatric acute ocular injury. *JAMA Ophthalmol.* 2018;136:895–903.
4. Négrel AD, Thylefors B. The global impact of eye injuries. *Ophthalmic Epidemiol.* 1998;5:143–69.

5. Krumholz HM. Where have all the heart attacks gone? New York Times; 2020. <https://www.nytimes.com/2020/04/06/well/live/coronavirus-doctors-hospitals-emergency-care-heart-attack-stroke.html>.
6. American Heart Association. Health emergency? Don't hesitate to get help. 2020. <https://www.heart.org/en/news/2020/03/30/health-emergency-dont-hesitate-to-get-help>.