

# EndoCompass Project: Research Roadmap for Pituitary and Neuroendocrine Tumor Endocrinology

Justo P. Castano<sup>a, b, c</sup> Mehul Tulsidas Dattani<sup>d, e</sup>  
Simona Grozinsky-Glasberg<sup>f</sup> Niki Karavitaki<sup>g, h</sup> Marianne E. Pavel<sup>i</sup>  
Cynthia Andoniadou<sup>j, k</sup> Krystallenia Alexandraki<sup>l</sup> Cristina Capatina<sup>k, m</sup>  
Manuela Cerbone<sup>n, o</sup> Diego Ferone<sup>p</sup> Hoong-Wei Gan<sup>o, q</sup>  
Johannes Hofland<sup>r</sup> Leo J. Hofland<sup>r</sup> Alejandro Ibanez-Costa<sup>a, b, c</sup>  
Mirela-Diana Ilie<sup>s, t</sup> Andrea M. Isidori<sup>u, v</sup> Marta Korbonits<sup>w, x</sup>  
Beata Kos-Kudła<sup>y, z</sup> Mohamad Maghnie<sup>A, B</sup> Giovanna Mantovani<sup>C, D</sup>  
Monica Marazuela<sup>E</sup> Gerald Raverot<sup>F, G, H</sup> Aldo Scarpa<sup>I</sup>  
Katharina Schilbach<sup>J</sup> Marily Theodoropoulou<sup>K</sup> Hanneke M. van Santen<sup>L, M</sup>  
Maria Chiara Zatelli<sup>N</sup>

<sup>a</sup>Department of Cell Biology, Physiology, and Immunology, University of Córdoba, Córdoba, Spain; <sup>b</sup>Maimonides Biomedical Research Institute of Cordoba, Córdoba, Spain; <sup>c</sup>Reina Sofia University Hospital, Córdoba, Spain; <sup>d</sup>Genetics and Genomic Medicine Programme, UCL GOS Institute of Child Health, London, UK; <sup>e</sup>Endocrinology Department, Great Ormond Street Hospital for Children NHS Trust, London, UK; <sup>f</sup>Neuroendocrine Tumor Unit, ENETS Centre of Excellence, Division of Medicine, Hadassah University Hospital and Faculty of Medicine, The Hebrew University, Jerusalem, Israel; <sup>g</sup>Institute of Metabolism and Systems Research, University of Birmingham, and Centre for Endocrinology, Diabetes and Metabolism, Birmingham Health Partners, Birmingham, UK; <sup>h</sup>Department of Endocrinology, Queen Elizabeth Hospital, University Hospitals Birmingham NHS Foundation Trust, Birmingham, UK; <sup>i</sup>Medizinische Klinik 1, Schwerpunkt Endokrinologie und Diabetologie, Uniklinikum Erlangen, ENETS Center of Excellence Erlangen, CCC Erlangen-EMN, Friedrich-Alexander-Universität Erlangen-Nürnberg, Erlangen, Germany; <sup>j</sup>Faculty of Medicine Carl Gustav Carus, Department of Medicine III, Technische Universität Dresden, Dresden, Germany; <sup>k</sup>Department of Endocrinology, Carol Davila University of Medicine and Pharmacy, Bucharest, Romania; <sup>l</sup>2nd Department of Surgery, Aretaieio Hospital, National and Kapodistrian University of Athens, Athens, Greece; <sup>m</sup>Department of Pituitary and Neuroendocrine Pathology, C.I. Parhon National Institute of Endocrinology, Bucharest, Romania; <sup>n</sup>Genetics and Genomic Medicine Research and Teaching Department, Great Ormond Street Hospital for Children, London, UK; <sup>o</sup>Department of Endocrinology, Great Ormond Street Hospital for Children, London, UK; <sup>p</sup>Endocrinology, Department of Internal Medicine and Medical Specialties (DiMI), University of Genoa, Genoa, Italy; <sup>q</sup>Genetics and Genomic Medicine Research and Teaching Department, University College London Great Ormond Street Institute of Child Health, London, UK; <sup>r</sup>ENETS Center of Excellence, Section of Endocrinology, Department of Internal Medicine, Erasmus Medical Center Cancer Institute, Rotterdam, The Netherlands; <sup>s</sup>Inserm U1052, CNRS UMR5286, Claude Bernard Lyon 1 University, Cancer Research Center of Lyon, Lyon, France; <sup>t</sup>Endocrinology Department, “C.I. Parhon” National Institute of Endocrinology, Bucharest, Romania; <sup>u</sup>Section of Medical Pathophysiology and Endocrinology, Department of Experimental Medicine, Sapienza University of Rome, Rome, Italy; <sup>v</sup>Endocrine and Andrological

Regional Rare Disease Center (Endo-ERN accredited), Policlinico Umberto I, Rome, Italy; <sup>w</sup>Department of Endocrinology, Barts and The London School of Medicine, Queen Mary University of London, London, UK; <sup>x</sup>William Harvey Research Institute, London, UK; <sup>y</sup>Department of Endocrinology and Neuroendocrine Tumours, Medical University of Silesia, Katowice, Poland; <sup>z</sup>Department of Pathophysiology and Endocrinology, Medical University of Silesia, Katowice, Poland; <sup>A</sup>Pediatric Clinic and Endocrinology Unit, Istituto di Ricovero e Cura a Carattere Scientifico (IRCCS) Istituto Giannina Gaslini, Genoa, Italy; <sup>B</sup>DINOEMI (Department of Neuroscience, Rehabilitation, Ophthalmology, Genetics, Maternal and Child Health), University of Genoa, Genoa, Italy; <sup>C</sup>Department of Clinical Sciences and Community Health, University of Milan, Milan, Italy; <sup>D</sup>Endocrinology Unit, Fondazione IRCCS Ca' Granda Ospedale Maggiore Policlinico, Milan, Italy; <sup>E</sup>Universidad Autónoma de Madrid, Department of Endocrinology and Nutrition, Hospital Universitario de La Princesa, Instituto de Investigación Sanitaria de La Princesa, Madrid, Spain; <sup>F</sup>Endocrinology Department, Hospices Civils de Lyon, Reference Centre for Rare Pituitary Diseases HYPO, "Groupement Hospitalier Est," Bron, France; <sup>G</sup>Faculté de Médecine, Université Claude Bernard Lyon 1-Domaine de Rockefeller institution, Lyon, France; <sup>H</sup>Cancer Research Centre Lyon, Inserm U1052, CNRC UMR5286, Lyon, France; <sup>I</sup>Department of Diagnostics and Public Health, Section of Pathology, University of Verona, Verona, Italy; <sup>J</sup>Medizinische Klinik and Poliklinik IV, LMU Klinikum München, Munich, Germany; <sup>K</sup>Department of Medicine IV, LMU University Hospital, LMU Munich, Munich, Germany; <sup>L</sup>Division of Pediatric Endocrinology, Wilhelmina Children's Hospital, University Medical Centre, Utrecht, The Netherlands; <sup>M</sup>Division of Pediatric Oncology, Princess Máxima Center for Pediatric Oncology, Utrecht, The Netherlands; <sup>N</sup>Section of Endocrinology and Internal Medicine, Department of Medical Sciences, University of Ferrara, Ferrara, Italy

## Significance

The research roadmap for pituitary diseases and neuroendocrine tumor endocrinology project has been an exciting and fruitful collaborative work of many experts highlighting the multitude of challenges associated with the diagnosis and management of these conditions across the lifespan. Systematic, adequately supported by funders' and policymakers' work that focuses on the many identified research priorities is vital for enlightening the mechanisms of disease/tumor biology and ultimately improving patient outcomes. The application of technical and methodological advances, including precision medicine, preclinical models, machine learning and artificial intelligence, are promising, valuable tools for the progress of impactful translational research in modern neuroendocrinology.

## Keywords

EndoCompass · Thyroid · Funding · Roadmap

## Abstract

**Background:** Endocrine science remains underrepresented in European Union research programs despite the fundamental role of hormone health in human wellbeing. Analysis of the CORDIS database reveals a persistent gap between the societal impact of endocrine disorders and their research prioritization. At national funding level, endocrine societies report limited or little attention of national research funding toward endocrinology. The EndoCompass project – a joint initiative between the European Society of Endocrinology and the European Society of Paediatric Endocrinology, aimed to identify and promote stra-

tegic research priorities in endocrine science to address critical hormone-related health challenges. **Methods:** Research priorities were established through comprehensive analysis of the EU CORDIS database covering the Horizon 2020 framework period (2014–2020). Expert consultation in pituitary and neuroendocrine tumor endocrinology was conducted to identify key research priorities, followed by broader stakeholder engagement including society members and patient advocacy groups. **Results:** Research priorities encompass congenital and acquired disorders, tumor development and progression, and neuroendocrine neoplasms. Key areas include genetic/epigenetic factors, tumor microenvironment, personalized treatment approaches, developing preclinical models, and implementing artificial intelligence. Special emphasis is placed on aggressive tumors, hypothalamic syndromes, and establishing pan-European

biobanks and databases. **Conclusions:** This component of the EndoCompass project provides an evidence-based roadmap for strategic research investment. This framework identifies crucial investigation areas into pituitary-neuroendocrine disease pathophysiology, prevention, and treatment strategies, ultimately aimed at reducing the burden of these disorders on individuals and society. The findings support the broader EndoCompass objective of aligning research funding with areas of highest potential impact in endocrine health.

© 2025 The Author(s).  
Published by S. Karger AG, Basel

## Introduction

### *Pituitary Disease*

Pituitary disease can manifest across the whole lifespan, and includes a variety of pathologies, from congenital hypothalamo-pituitary disorders to pituitary tumors and other lesions in this area. The rarity of many of these conditions not only poses challenges for diagnosis and management, but also impacts research efforts to understand their pathogenesis and factors implicated in their prognosis. Comprehensive understanding of the mechanisms of tumor development, regrowth, and aggressive behavior remains deficient, making the discovery of more effective therapeutic agents more difficult.

Congenital hypothalamo-pituitary disorders, with their plethora of phenotypic manifestations, require further elucidation of their underlying pathophysiology, as does the often-daunting hypothalamic syndrome. Finally, the application of technical and methodological advances, including precision medicine, preclinical models, machine learning, and artificial intelligence (AI), constitutes a promising challenge. Their expected contribution to the progress of translational research is a major priority in modern neuroendocrinology.

### *Neuroendocrine Neoplasms*

Neuroendocrine neoplasms (NENs) are malignancies arising from the diffuse endocrine system, which comprises neuroendocrine cells dispersed throughout the bronchopulmonary and gastrointestinal tracts. Incidence and, particularly, prevalence of these neoplasms are steeply rising, making gastroenteropancreatic NENs the second most prevalent malignancy in the gut and the 10th most prevalent malignancy overall [1, 2].

Diagnosis and treatment of NENs are both challenging because of their heterogeneous biological behavior, with more than 60% being metastatic at diagnosis. One major unmet need arises from the paucity of curable/effective

therapeutic options. The only treatment that offers cure is surgery, possible in few patients with localized disease, while unresectable NENs are treated with a variety of noncurative options (somatostatin analogs, peptide receptor radioligand therapy, targeted agents, or chemotherapy), which eventually fail due to drug resistance, stressing the need for new therapeutic avenues [3]. Another unmet need is the lack of reliable biomarkers to guide management [4] as patients with the same tumor grade/stage behave heterogeneously in terms of time to tumor progression, tumor recurrence, or treatment response.

## Genetics and Epigenetics in Adult Pituitary and Hypothalamic Disease

There is a need to explore genetic and epigenetic factors, including the role of chromatin architecture and regulatory DNA sequences, in pituitary and hypothalamic disorders.

### *Epidemiology, Societal Impact, and Research State of the Art*

Understanding the interplay of genetic and epigenetic factors is essential to improve diagnosis and treatment in patients with pituitary adenomas and arginine vasopressin deficiency (AVP-D). The majority of pituitary adenomas are sporadic, while ~5% are familial or isolated, often presenting at a younger age and having a poorer prognosis [5, 6].

Recurrent somatic variants in *GNAS* and *USP8* exist in around 40% of somatotrophinomas and about 35% of corticotrophinomas, respectively [5]. Aggressive tumors show a relatively high prevalence of *TP53*, *ATRX*, and *SF3B1* variants [7, 8]. Epigenetic modifications in pituitary adenomas are cell lineage-dependent and less explored [9, 10]. Similarly, copy number variations and chromosomal gains and losses in pituitary adenomas are more likely to be histotype-specific [5]. AVP-D is characterized by genomic alterations on genes involved in the vasopressin/aquaporin axis [11, 12].

### *Future Research Priorities*

There is a need for refined diagnostic procedures for the early diagnosis of familial cases and timely prediction of aggressive tumor behavior and prognosis. Technical advances are expected to allow us to elucidate the mutational landscape of pituitary tumors beyond *GNAS* in somatotrophinomas and *USP8* in corticotrophinomas. In the long term, we need to intensify research into the epigenetic mechanisms that underpin the pathophysiology of pituitary adenomas, focusing on lineage-specific patterns of

gene demethylation and chromatin accessibility. We also need to decipher the genetic alterations contributing to hypothalamic disorders. Given the complexity and rarity of cases, interdisciplinary collaboration is essential to effectively address these challenges.

#### *Anticipated Impact of Future Research*

Advanced research in pituitary and hypothalamic disorders will have a major impact on understanding their pathophysiology and translation to effective clinical management. Early detection of familial cases of pituitary adenomas, hypothalamic disorders, and aggressive behavior of tumors is essential for timely intervention and improved patient outcome. Unraveling the complex genetic and epigenetic mechanisms underlying these disorders is expected to lead to earlier, more accurate diagnosis and personalized treatment strategies that are effective and have fewer side effects. These advancements will facilitate the precision management of patients with pituitary and hypothalamic disorders.

### **Tumor Microenvironment in Adult Pituitary Tumorigenesis**

An increased understanding is needed of the importance of the environment surrounding a tumor for pituitary tumor pathogenesis and clinical utility.

#### *Epidemiology, Societal Impact, and Research State of the Art*

Beyond genetics and epigenetics, the environment within which a tumor exists is defined as the tumor microenvironment. It includes immune and stromal non-tumor cells, vascular structures, and noncellular components (extracellular matrix) in a cross-talking continuum.

The tumor microenvironment opened a new area of research to shed light not only on the pathophysiology of neoplasms but also on the identification of those of its components that may provide drug targets [13]. Recently, the tumor microenvironment has been intensely investigated in pituitary tumors, particularly in the subset, which display an aggressive clinical course with rapid progression and resistance to conventional treatments. Some of the potential therapeutic targets among the cellular or non-cellular components of the tumor microenvironment have been used in solid tumors as well as in aggressive pituitary tumors. Hence, components of the tumor microenvironment may have a clinical utility, not only as targets for drugs but also as prognostic biomarkers for resistance to treatment, leading to more efficient management [14, 15].

#### *Future Research Priorities*

The future research priorities in studying the tumor microenvironment are based on investigation of the following:

1. New tumor microenvironment biomarkers that will identify the more aggressive pituitary tumors in the context of proliferation, and the pituitary tumors resistant to conventional treatment.
2. Genetic, histopathological, and molecular imprints, which enable selection of appropriate drugs for use in each case as a monotherapy or combination therapy, as well as the timing of early management before progression and the most effective sequence of each agent to be used.
3. AI to build an algorithm for individualized treatments, using a combination of omics, epigenetics and tumor microenvironment factors, including the immune component [16, 17].

The methodology to achieve the above goals includes establishment of a biobank (solid and liquid biopsies), as well as common European databases with basal information on patients and characteristics of pituitary tumors, including all the biomarkers studied. Thereafter, randomized, multicenter studies will need to establish the factors influencing aggressiveness and resistance to treatment [18].

#### *Anticipated Impact of Future Research*

Immunotherapy and antiangiogenic therapies that target components of the tumor microenvironment have been attempted in a few published cases of aggressive pituitary tumors, with positive outcomes. The rationale for using these therapies is based on the central premise that the tumor microenvironment impacts the pathophysiology of pituitary tumors.

On the other hand, the use of common centralized European pituitary tumor biobanks and databases will help facilitate and accelerate the identification of the tumor microenvironment components that can be used more efficiently as prognostic and predictive biomarkers or as drug targets for pituitary tumor management.

### **Treatment Options for Adult Pituitary Tumors**

(New) treatment options must be investigated for resistant prolactinomas, nonfunctioning pituitary adenomas and Cushing's disease, particularly in settings of recurrence.

### *Epidemiology, Societal Impact, and Research State of the Art*

#### Prolactinomas

Resistance to dopamine agonist (DA) treatment is uncommon in microprolactinomas and more frequent in macroprolactinomas and invasive tumors. Primary DA resistance occurs in around 20%–30% of the cases treated with bromocriptine and 10% of those treated with cabergoline [19]. Secondary DA resistance is rare and is usually associated with aggressive or malignant tumor behavior [19]. Management approaches in resistant prolactinomas include changing DA, surgery, radiotherapy, temozolomide (in aggressive tumors), or combinations of these. However, these approaches are not always successful [7, 19–21].

#### Nonfunctioning Pituitary Adenomas

Nonfunctioning pituitary adenomas have a prevalence of between 7/100,000 and 41.3/100,000 and are associated with various morbidities and increased mortality [22, 23]. Surgery combined, or not, with radiotherapy is the main means of treatment, aiming to safely remove the tumor and stop its regrowth. Despite these approaches, regrowth occurs in several patients (the risk of which is currently defined by clinical and pathological parameters). A subset of these tumors demonstrates aggressive behavior [24–26]. Following tumor regrowth, repeat surgery and/or radiotherapy are the first-line approaches, and DAs have been tried for prevention of growth of nonfunctioning pituitary adenomas [24, 27, 28].

#### Cushing's Disease

Cushing's disease is associated with increased morbidity and mortality [29]. After surgery, the recurrence rate is around 30%. For those cases, radiotherapy and combination drug therapy, targeted for biochemical control (steroidogenesis inhibitors and glucocorticoid-receptor blockers) and tumor control (pasireotide and DAs), are available. The latter have only limited effectiveness and, in some cases, severely impairing side effects.

In cases that are refractory to therapy, aggressively growing corticotrophinomas and adrenocorticotrophin-secreting carcinomas, there have, so far, only been off-label therapy trials with temozolomide, and attempts at individualized therapy with immune checkpoint inhibitors, tyrosine kinase inhibitors and angiogenesis inhibitors, all potentially associated with serious side effects [26].

Following the discovery of the activating mutation in the *USP8* gene and changes in regulators of the cell cycle (cyclin-dependent kinases) in Cushing's Disease, potential molecular targets are now available [30, 31].

### *Future Research Priorities*

#### Prolactinomas

- Understand the molecular mechanisms involved in resistance to DAs.
- Develop new alternative medical treatments and investigate the efficacy and safety of those proposed to date (e.g., estrogen receptor antagonists, aromatase inhibitors, pasireotide, lapatinib, everolimus, and immune checkpoint inhibitors).

#### Nonfunctioning Pituitary Adenomas

- Understand the mechanisms implicated in regrowth of nonfunctioning pituitary adenomas after primary treatment.
- Identify molecular prognostic factors associated with regrowth.
- Identify targets for pharmacotherapy that will prevent regrowth or be used as primary medical treatment, particularly for tumors not requiring prompt surgical intervention.

#### Cushing's Disease

- Identify further molecular mechanisms involved in corticotroph tumors.
- Further investigate existing drugs for molecular targets (i.e., seliciclib) and develop new, tumor-targeted, well-tolerated, and, ideally, oral therapy for recurrent Cushing's disease.

### *Anticipated Impact of Future Research*

In all of the above types of tumors, the outputs of the proposed research will improve management pathways by expanding treatment options and applying personalized medicine. In the long term, they will have a positive impact on the outcomes and prognosis of the patients.

### **Aggressive Behavior and Targeted Treatment in Adult Pituitary Adenomas**

There is a need to identify molecular factors associated with invasive behavior or in predicting aggressive behavior in pituitary adenomas, and to develop targeted therapies.

### *Epidemiology, Societal Impact, and Research State of the Art*

The prevalence of invasive and aggressive pituitary tumors varies widely depending on the definition or criteria used and on the type of patients included. Nevertheless, aggressive pituitary tumors are rare; a recent study concluded that around 0.5% of all clinically

detected pituitary tumors have an aggressive course [25]. On the other hand, invasive pituitary tumors are common (14–40% of surgically treated cases) [32, 33]. Both invasive and aggressive pituitary tumors incur increased morbidity and mortality.

Radiological evidence of tumor extension into the cavernous sinus is not synonymous with invasion per se as this extension can correspond solely to tumor expansion [34, 35]. Moreover, the modified Knosp classification, widely used to define invasiveness, was only intended to predict the probability of intraoperatively observed cavernous sinus invasion and the surgical outcome [34]. Thus, prior to identifying molecular factors associated with invasiveness, clear definitions are needed.

Predicting aggressive behavior remains challenging. A combined clinicopathological classification has proven its value in predicting progression or recurrence following surgery [36] but remains, nevertheless, limited in predicting future aggressive behavior. Recently, a Ki67 index  $\geq 10\%$  has been proposed to predict future malignancy [37], but its value has not yet been proved. Identification of targeted therapies (including combinations), along with establishment of predictors of response to treatment, are needed most, especially following temozolomide failure.

#### *Future Research Priorities*

1. It is important to homogenize how “invasiveness” and “expansion” are defined, for both clinical and research purposes.
2. Investigations should determine whether “invasiveness” and/or “expansion” is associated with worse prognosis, independent of the fact that invasiveness/expansion leads to incomplete surgical removal of the tumor.
3. From a biological point of view, investigations are needed to show whether true “invasiveness” has a different molecular and cellular basis compared with simple “expansion.” These mechanisms should be dissected to identify a marker that is useable in clinics, as well as new therapeutic targets. We strongly recommend the use of precision technological approaches (e.g., single-cell transcriptomics, as opposed to bulk transcriptomics), followed by validation at the protein level and, ideally, outcomes that are further investigated by mechanistic in vitro and/or in vivo studies. The application of other cutting-edge technologies (such as single-cell multiomics and spatial transcriptomics) is strongly encouraged.
4. Given the rarity of aggressive pituitary tumors and carcinomas, it is of utmost importance to create pan-European Union or, at least, national databases (with

clinical, pathology and molecular data), coupled with biobanks that are as comprehensive as possible (tumor tissue, blood). This will enable understanding of tumor biology, identification of therapeutic targets, and identification of predictive/prognostic markers. Including all surgical specimens available in the biobank (initial tumor, recurrences, and metastases) will be crucial in order:

- to investigate at what point the tumor phenotype has changed and, hence, test whether the identified markers of aggressiveness could serve as predictive makers, prior to clinical evidence of aggressive behavior
  - to study potential predictors of response to different treatments.
5. Similar to the above, a combination of multiomics studies and wet lab approaches should be applied, ideally within consortia with both strong bioinformatics and wet lab expertise.
  6. It is important to test the identified predictive markers of future aggressiveness among large cohorts of both aggressive and benign pituitary tumors with a long follow-up (at least 5 years), in order to assess the prevalence and the specificity of the markers. While waiting for additional markers, one marker that is already proposed in the literature and would be simple to test is an initial Ki67 index  $\geq 10\%$  (see above).
  7. Given the rarity of patients requiring therapeutic options beyond temozolomide, pan-European Union or, at least, national multicenter studies are needed to assess the efficacy and potential predictors of response to such treatments.

#### *Anticipated Impact of Future Research*

From a biological perspective, this research is expected to increase the understanding of the tumorigenesis-related processes linked to invasiveness and aggressiveness. From a clinical perspective, this research is expected to inform future guidelines and to directly benefit patients, leading to personalized approaches, decreased morbidity and mortality, and increased quality of life and survival.

### **Congenital Hypothalamo-Pituitary Disorders and New Treatment Options in Children**

#### *Epidemiology, Societal Impact, and Research State of the Art*

Congenital hypopituitarism is characterized by single/multiple pituitary hormone deficiencies, often associated with midline brain/facial/eye and/or forebrain abnormalities,

including septo-optic dysplasia (SOD) and holoprosencephaly. It affects 1/3,000 to 1/4,000 children, with an incidence of SOD in Europe of 1.9–2.5/100,000 births.

Patients with congenital hypopituitarism, particularly SOD, can manifest autism, complex behavioral disorders, sleep dysregulation, and hypothalamic dysfunction including inexorable obesity. The impact on their families and wider society can be considerable, with few patients able to take up full-time employment, and much probably requiring lifelong supervised care. There is an increased risk of mental health problems, including self-harm.

Research suggests that congenital hypopituitarism arises from abnormal hypothalamo-pituitary development. Variants in over 67 genes have been identified, but account for only 10% of patient phenotypes. These genes encode transcription factors, signaling molecules, eukaryotic translation initiation factors and components of splicing machinery. Inheritance can be X-linked, autosomal dominant or recessive with variable penetrance [38, 39].

While SOD represents a spectrum of malformations with unpredictable endocrine and neuroimaging phenotypes, congenital hypopituitarism associated with multiple pituitary hormone deficiencies is more homogeneous [40]. Stalk and posterior pituitary abnormalities increase the risk of early onset deficits.

#### *Future Research Priorities*

There needs to be a greater understanding of the phenotypic heterogeneity observed, including growth in the absence of growth hormone (GH), unusual patterns of GH secretion, evolution of hormonal deficiencies, and pubertal development and obesity in SOD. Clinical features, such as impaired olfaction, pain insensitivity, and autonomic dysfunction, need to be studied in more detail. Improved neuroimaging techniques with better visualization of the hypothalamus and pituitary stalk may help predict clinical/metabolic outcomes and aid targeted therapy. Long-acting GH and glucocorticoids may improve quality of life, alongside treatments for hypothalamic obesity.

Research should also focus on mental health, neurocognition, behavior, autism, attention deficit hyperactivity disorder, aggression with self-harm, and sleep disorders. Additionally, given that the majority of cases remain genetically unexplained, we need to improve our understanding of the etiology of congenital hypopituitarism using comprehensive, unbiased modern mul-

tiomic techniques, including genomic, epigenomic, transcriptomic, and proteomic approaches, with the inclusion of AI.

#### *Anticipated Impact of Future Research*

Better understanding of the underlying pathophysiology and underlying molecular basis will not just improve genetic counseling, but also lead to improved prediction of the clinical trajectory of endocrinopathies and associated hypothalamic dysfunction, with better clinical outcomes.

### **Acquired Hypothalamo-Pituitary Disorders and New Treatment Options in Children**

#### *Epidemiology, Societal Impact, and Research State of the Art*

Acquired hypothalamo-pituitary dysfunction can be caused by (supra)sellar tumors or their treatment, traumatic brain injury, infection, autoimmune processes, granulomatous disease, iron overload, or vascular insults. Endocrinopathies and hypothalamic dysfunction can evolve, causing significant morbidity and mortality. The close proximity to the optic chiasm may cause associated visual impairment.

The combination of hypothalamo-pituitary dysfunction, visual impairment, and neuro-behavioral disturbance can severely impact the quality of life of patients and their families, with significant socioeconomic cost, and increases the risk of mortality [41, 42]. Current research focuses heavily on the genetic etiology of these tumors to identify novel therapeutic targets.

#### *Future Research Priorities*

Targeted therapies may aid in limiting the role of surgery and radiotherapy (e.g., craniopharyngiomas [43], Cushing's disease) or in overcoming treatment resistance (e.g., prolactinomas and somatotrophinomas). Elucidation of tumorigenic pathways may help in the development of newer, less toxic treatments. Additionally, more intensive treatment may be offered for tumors deemed high risk at a molecular level.

Neuroimaging needs to improve to achieve more accurate diagnoses and to grade hypothalamic tumoral invasion and the degree of disruption of the surrounding hypothalamic neurocircuitry, in order to increase the precision of treatment strategies. New therapies also need to be developed for other, rarer causes of acquired hypothalamo-pituitary dysfunction, such as hypothalamitis and hypophysitis, and sarcoidosis or granulomatous disease.

### *Anticipated Impact of Future Research*

Improved neuroimaging and molecular techniques will improve diagnostic accuracy and risk stratification of acquired hypothalamo-pituitary lesions [44], as well as the development of new treatments for tumors in this region. It will lead to fewer off-target effects, reducing the injury to healthy brain tissue incurred by neurosurgery and radiotherapy. Inclusion of markers of hypothalamo-pituitary dysfunction in clinical trials for novel oncology treatments will clarify the effect of such interventions on long-term quality of life. Overall, this will improve progression-free and event-free survival, reduce toxicity and adverse effects on the hypothalamo-pituitary axis, optimize metabolic and reproductive outcomes and, ultimately, improve the quality of life of survivors and their families.

### **Hypothalamic Syndrome and New Treatment Options in Children**

#### *Epidemiology, Societal Impact, and Research State of the Art*

“Hypothalamic syndrome” encompasses a range of clinical features: dysregulated appetite and thirst, inexorable morbid obesity, multiple endocrinopathies (including deficiencies of adrenocorticotrophin and AVP), thermoregulation issues, fatigue, and behavioral, mental health, and sleep difficulties [45–48].

Due to our currently incomplete understanding of the complex neuroendocrine hypothalamic circuitries, many of these problems remain untreatable. This results in morbidity and disability, with a consequent burden on health systems as well as national economies.

Recently, newer molecular techniques have revealed the components of the hypothalamic circuitry, and the importance of hypothalamic hormones in appetite and weight regulation [45]. Correlation of function with neuroimaging markers has been difficult as conventional magnetic resonance imaging (MRI) cannot precisely define the hypothalamic nuclei.

#### *Future Research Priorities*

At present, there are no universal biochemical or neuroradiological diagnostic criteria for hypothalamic syndrome. More precise neuroendocrine, genomic and epigenomic markers to pinpoint pathogenic changes need to be identified, to better understand and treat the various domains of hypothalamic dysfunction. Similarly, there is a need to establish the etiology of other congenital hypothalamic syndromes, such as ROHHAD(-NET) (rapid-onset obesity with hypoventilation, hypo-

thalamic dysfunction, autonomic dysregulation, and neuroendocrine tumor [NET]).

Advanced neuroimaging is required to more precisely define hypothalamic damage and determine how this correlates with the phenotypes observed. Prospective intervention studies are needed for the treatment of hypothalamic obesity and fatigue. The high rate of neuropsychological disturbances and autistic spectrum and mental health disorders associated with hypothalamic syndromes also needs to be better understood to develop effective interventions.

### *Anticipated Impact of Future Research*

The development of successful therapies for hypothalamic obesity will reduce long-term complications, such as cardiovascular morbidity and type 2 diabetes. Identifying neuroradiological correlates of hypothalamic dysfunction will allow the development of medical treatments and reduce the risk of complications, by improving the precision of neurosurgery and radiotherapy. Understanding the nature of congenital hypothalamic syndromes such as ROHHAD(NET) will aid early diagnosis, genetic counseling, and development of targeted therapies for this life-limiting condition. Overall, achievement of these research priorities will improve the long-term quality of life and reduce long-term mortality for these patients, as well as decrease the significant socioeconomic burden on health- and social-care systems.

### **Developing Precision Medicine Approaches for Pituitary Tumors**

#### *Epidemiology, Societal Impact, and Research State of the Art*

Precision medicine for pituitary tumors involves tailoring diagnosis, treatment, and monitoring strategies to the individual characteristics of each patient and their tumor. This requires identification of biomarkers to guide patient treatment based on individual conditions, as is being pioneered in acromegaly [49].

Evolution of image guidance and visualization technologies in pituitary surgery, including intraoperative MRI suites and biological imaging strategies, will be critical in this regard [50, 51]. Indeed, the endoscopic transsphenoidal approach has improved earlier surgical outcomes [52]. However, the anatomical complexity and heterogeneous biology of pituitary tumors challenge their management, demanding multimodal integration of surgery, radiosurgery, radiation therapy, and medical therapy [53].

### *Future Research Priorities*

The development of precision pituitary medicine will require genomic sequencing of each tumor to identify specific mutations, gene expression patterns and molecular signatures, and to predict their impact on tumor behavior. This could facilitate the establishment of a true clinico-molecular classification and stratification of tumors and the identification of molecular targets in specific tumor subtypes.

Circulating biomarkers (e.g., hormones, cytokines, and metabolite levels) could leverage this approach, jointly providing information on tumor progression, invasion, and therapy response, improving patient monitoring and prediction of prognosis. Advancing this strategy requires the development of algorithms to integrate clinical, molecular, and imaging data, in order to stratify patients into risk groups based on tumor aggressiveness, recurrence potential and treatment responsiveness. In turn, this will allow surveillance protocols and treatment interventions to be tailored according to individual risk profiles.

Additionally, targeted therapies will have to be developed, such as small molecule inhibitors, monoclonal antibodies or peptide receptor ligands, selectively disrupting tumor growth and hormone secretion while minimizing off-target effects. This will only be possible through advanced integrative computational models that are able to predict treatment outcomes, recurrence risk, and long-term prognosis for individual patients, while enabling algorithm refinement by machine learning from patient outcomes and real-world data. Naturally, this necessitates interdisciplinary and international collaborations among researchers, clinicians, patient advocates, and industry partners.

### *Anticipated Impact of Future Research*

Integrating these strategies will accelerate discovery and translation and boost precision medicine approaches for pituitary tumors. This can revolutionize their management and offer personalized diagnostic, therapeutic, and prognostic solutions tailored to the unique characteristics of each patient. Implementation of precision pituitary tumor medicine will impact the future of research, healthcare systems and patient care by providing more accurate diagnosis and prognosis prediction tools, to help tailor treatment.

In addition, molecular profiling can lead to more effective and personalized therapies, improving patient outcomes and quality of life, and helping to overcome treatment resistance by applying molecularly targeted therapies to each tumor. Overall, the future impact of precision medicine on pituitary tumors promises to be transformative, offering patients more personalized and effective treatment options, while advancing our understanding of tumor biology and improving healthcare outcomes.

## **Developing Preclinical Models for Hypothalamic and Pituitary Pathophysiology**

### *Epidemiology, Societal Impact, and Research State of the Art*

Establishing *in vitro* or *in vivo* models to study hypothalamus/pituitary function is inherently challenging, due to the complexity of this neuroendocrine axis, which involves intricate interactions between diverse cell types and multiple hormones, regulated through dynamic feedback mechanisms. Accurately modeling these interactions in the laboratory require the precise control of differentiation and functional maintenance of specific cell types.

Pituitary tumor cell models first established in 1968 (rat GH3, mouse AtT-20 and, later, rat MMQ cells) are still some of the few cell lines that are widely used in research, despite the limitation of their rodent origin [54]. Numerous attempts to develop human cell lines have failed to faithfully replicate any specific type of pituitary tumor. To circumvent this, labs commonly employ primary cell cultures derived from human pituitary tumors, a more pathophysiologically relevant model compared with immortalized cell lines, although with a limited lifespan and high variability [55, 56].

Fortunately, recent advances in cell culture techniques and stem cell and organoid research have facilitated the development of human and murine pituitary organoids derived from embryonic stem cells, induced pluripotent stem cells, and fetal or adult stem cells [57]. Indeed, various models are currently available for the normal and affected pituitary, including hypopituitarism, craniopharyngioma and anterior pituitary tumors, especially corticotrophinomas and prolactinomas. These may serve to evaluate drugs, assess tumor growth and restore pituitary function [58]. Similar approaches have been employed for hypothalamic studies [59] and, although no perfect models can recapitulate pituitary/hypothalamus dysfunction, several recent models can be used, depending on the disease [60].

*In vitro* systems lack the capacity to mimic the complex feedback loops and environmental influences (stress, nutrition, and circadian rhythms), further limiting hypothalamo-pituitary studies. Nevertheless, 3D chips are being developed to emulate hypothalamus-pituitary crosstalk [61].

### *Future Research Priorities*

Future research should prioritize the development of pituitary and hypothalamic organoid models to better replicate the complex cellular interactions and dynamic

feedback mechanisms inherent to these systems. Advances in stem cell technology and bioengineering must be leveraged to create more accurate and functional in vitro models. Additionally, developing 3D culture systems and organ-on-a-chip technologies will be crucial to simulate physiological conditions such as stress, nutrition and circadian rhythms, as well as enabling detailed investigation of disease mechanisms, leading to the discovery of new therapeutic targets and more effective treatments. Efforts should be focused on creating human-derived cell lines to improve translational relevance and facilitate personalized medicine applications for neuroendocrine disorders.

#### *Anticipated Impact of Future Research*

Development of more accurate and functional models should revolutionize the understanding of pituitary and hypothalamic pathophysiology. These advancements will improve drug screening processes, reduce the dependence on animal models, and facilitate personalized treatment strategies. Ultimately, this research holds the potential to restore pituitary function in patients with related disorders, significantly improving their quality of life and clinical outcomes.

### **Machine Learning and AI in Research and Management of Pituitary Diseases**

#### *Epidemiology, Societal Impact, and Research State of the Art*

Artificial intelligence and machine learning have gained momentum in recent years as integrative techniques for diagnosis and intervention in pituitary diseases. They enable more holistic and multidimensional approaches and facilitate the management of the increasingly large amount of data associated with clinical and basic research [62].

For clinical research, the main areas for application of machine learning are predictive algorithms for tumor aggressiveness and clinical outcomes of surgery and medical treatment [51, 63]. Algorithms may become helpful tools in complex differential diagnoses, but differences in training sets could affect the replicability of results. In basic research, machine learning is usually applied to pathomic and transcriptomic studies.

#### *Future Research Priorities*

Two critical issues in studies involving machine learning are appropriate sample size and algorithm/model homogeneity. Many different algorithms have been used in pituitary research to train datasets, including a broad range of subjects. Moreover, predictive algorithms are often designed

to replicate clinician-made evaluations (texture, development of pituitary deficiencies), which limits their impact on clinical decisions. Surgical outcomes are usually center-specific. With larger, collaborative datasets, radiomic studies could help unveil other disease-specific characteristics of pituitary neoplasms. Moreover, longitudinal studies with standardized follow-up intervals might help identify patients who require more proactive interventions than those in which “watchful waiting” could be a reasonable approach.

For basic research, pathomic studies could provide insightful results on pituitary neoplasm recurrence and aggressiveness, and patient response rate to standard medical regimens. Algorithm analysis of pathological images, ideally integrated with clinical and transcriptomic data, could help identify reliable predictors guiding treatment and follow-up regimens. Furthermore, identifying neoplasm-specific biomarkers in peripheral blood could help develop reliable “liquid biopsies,” allowing less invasive diagnosis and surveillance.

Complex machine learning techniques, such as neural networks, are often burdened by a lack of explainability, which is a mainstay of transparency in the use of AI algorithms in healthcare. However, machine learning-assisted clinical decisions will become increasingly prevalent in the coming decades. Therefore, the aim should be to combine data from different countries and settings, minimizing training-related biases, such as the underrepresentation of minorities.

#### *Anticipated Impact of Future Research*

Patient-centered precision medicine is the future of healthcare, where each person will be characterized by an increasing amount of data. Patients with pituitary disorders face numerous challenges during their lifespan, often with little knowledge at diagnosis of what their path will be. Predictive algorithms could help healthcare professionals identify “high-risk” and “low-risk” patients, probability of remission, and development of complications, so empowering people with pituitary disorders to be part of the decision-making process. Moreover, machine learning techniques could help bridge knowledge gaps in rare pituitary diseases by unveiling new patterns and developing novel therapeutic strategies.

### **Understanding NENs Biology**

#### *Epidemiology, Societal Impact, and Research State of the Art*

Neuroendocrine neoplasms are divided into well-differentiated NETs and poorly differentiated neuroendocrine carcinomas (NECs) [64]. Ki67 is the most

important prognostic biomarker of NENs, marking cycling cells, a crucial tool for grading [65]. For diagnostic purposes, alterations in TP53 and RB1 are considered biomarkers of NECs [66]. In contrast, MEN1, DAXX, and/or ATRX mutations are typically associated with NETs in some districts [67]. DNA methylation profiling has recently been proposed as a powerful source of diagnostic and prognostic biomarkers [68]. With the introduction of new therapeutic strategies into the clinic, new biomarkers with predictive value are urgently needed.

#### *Future Research Priorities*

Beyond the distinction between NETs and NECs, the entire NENs category should be considered a continuous entity, without the well-demarcated boundaries imposed by classification needs. Thus, one of the most crucial research priorities is to address the large gray areas within each neoplasm subgroup. One example is the G2 category, which includes tumors with totally different proliferation and clinical behavior. The most advanced next-generation techniques should be implemented in this category, including single-cell omics, with epigenomics and transcriptomics [69]. In addition, the use of spatial transcriptomics to investigate the tumor microenvironment is a promising methodology to further explore this heterogeneous tumor context [70].

Historically, improvements in establishing experimental models of NETs have been modest. Cell lines derived from different NETs have been generated, but most cell lines are only partially representative for their NETs of origin. Similar issues relate to the use of NETs animal models. NETs xenograft models are difficult to establish and generally slow-growing [71]. In recent years, promising 3D spheroid and organoid models of mainly high-grade NETs and NECs have been established [72, 73], but there remains an unmet need for preclinical models of low-grade NETs. The majority of NET models does not address tumor heterogeneity or incorporate the complex tumor microenvironment. Future research should address these issues, by 3D techniques and novel technologies such as tumor-on-a-chip [74].

#### *Anticipated Impact of Future Research*

Developments in molecular and preclinical research in the field of NENs are expected to foster progress in understanding the mechanisms underlying neoplasm formation, and, ultimately, to fundamentally improve their diagnostics, therapeutic methods and patient outcomes. Thus, identification of new biomarkers, ac-

companied by designing reliable molecular tests, and advances in anatomical and functional imaging techniques, will improve the current methods to diagnose and monitor NENs and to predict their progression.

Likewise, advancements in treatment strategies, including new therapeutic targets in the field of biotherapy, molecular targeted therapies, radioisotope therapies, chemotherapy, and immunotherapy (as well as combinations of these therapies) should provide more effective treatment options, especially for advanced NENs. Additionally, innovative surgical techniques, including minimally invasive methods, will foster improved outcomes.

However, as we embrace the era of precision medicine, personalized, individualized treatment approaches based on genetic tests will become available to optimize therapy selection and sequencing and, thereby, minimize side effects. Most importantly, research and technological advances will focus on the patient, leading not only to better treatment outcomes and enhanced survival rates, but also to an improvement in the quality of life of patients. In summary, research advances should facilitate a better understanding of NEN pathobiology and leverage the development of improved and early diagnostic tools and more effective treatments, ultimately benefiting patients with NENs and healthcare systems alike.

### **How to Improve Therapy in NENs**

#### *Epidemiology, Societal Impact, and Research State of the Art*

The existing treatment options in advanced NENs are limited, highlighting the unmet need for novel therapeutic approaches. One of the new options includes immune-based treatments, which have been at the epicenter of cancer therapy in recent years. Immunotherapy has been evaluated in NENs of different origin and differentiation, but with limited efficacy shown only in some high-grade NENs, and particularly as combination therapy [75]. Another option includes cancer vaccines, based either on tumor-associated or on tumor-specific antigens. Carcinoembryonic antigens and chimeric antigen receptors are under development [76]. The molecular profiles of patients have also been exploited with limited success in rare individual cases where mutations have been identified that could provide drug targets [77]. Additionally, identification of the preferred sequence in existing therapies is another major unmet need. Despite initial insights from recent studies, a general approach for application in clinical practice is missing [78].

### *Future Research Priorities*

Future research priorities in NENs treatment are based on personalized medicine, specifically focusing on the investigation of: (1) new biomarkers to identify the disease and appropriate timing of early management before progression, (2) patient-derived organoids for closer to real-life screening of drug efficacy, and (3) pathological and/or molecular imprints to individualize the sequence and combinations of treatments (omics, immunological, epigenetic, and microenvironmental factors).

The methodology should include establishing biobanks (with solid and liquid biopsies) inside a common European network of expert centers, to integrate information on patients and their respective NENs, including disease features and treatment sequence/outcomes. Moreover, this approach will allow randomized, multicenter studies to be undertaken, focusing on molecular profiles pre-/post-surgery and pre-/post-therapies, through the investigation of omics, metabolic alterations, cancer immunology and microenvironment, and mapping the different neoplasms. Integrating the use of novel models such as organoids, 3D tumor cultures and xenografts into this research network will enable testing of novel drugs and potential cancer vaccines. Finally, AI can help pinpoint the right algorithms for combination and sequential therapies for each patient group.

### *Anticipated Impact of Future Research*

Establishing a common European biobank and registry network will help identify novel prognostic and predictive biomarkers, the right timing and sequencing of existing drugs, and novel, efficient, individualized approaches, ultimately improving the prognosis and quality of life of patients with NENs.

## **Fibrosis in NETs**

### *Epidemiology, Societal Impact, and Research State of the Art*

Neuroendocrine tumors retain the ability to secrete hormonally active substances and consequently to elicit hormonal symptoms in patients. The most common hormonal syndrome, encountered in ~20% of patients with NETs, is carcinoid syndrome, which is characterized by secretory diarrhea, flushing and fibrotic complications [79], and reduced quality of life compared with patients who do not have the syndrome [80, 81]. Carcinoid syndrome is predominantly encountered in patients with a metastatic NET originating from the small intestine, followed by those of lung, pancreatic,

ovarian, or thymic origin. Although serotonin is considered a critical mediator of carcinoid syndrome, the condition's precise pathophysiology and determinants are incompletely understood [82].

Importantly, fibrotic complications may be observed in patients with NETs, usually, but not exclusively, within the context of carcinoid syndrome. These include mesenteric fibrosis and carcinoid heart disease.

Mesenteric fibrosis is a pathognomonic feature of mesenteric metastases of small intestinal NETs, characterized by radiating fibrotic strands from a central mesenteric mass. Local fibrosis can induce venous ischemia, ileus, obstruction, and perforation of the bowel, and is associated with an impaired overall survival.

Carcinoid heart disease is encountered in 20%–30% of patients with carcinoid syndrome. It is characterized by right-sided heart valve and endocardial fibrosis, potentially leading to tricuspid valve and pulmonary valve regurgitation and stenosis and, ultimately, right-sided heart failure with increased mortality and morbidity.

In both mesenteric fibrosis and carcinoid heart disease, there appears to be a crucial role for serotonin derived from local and/or systemic NETs cells, and activation of its 5-HT<sub>2b</sub> receptor [83]. Besides serotonin, several other profibrotic factors have been postulated to contribute to the development of NET-associated fibrosis, including prostaglandins, histamine, bradykinin, tachykinins (substance P, neurokinin A, and neuropeptide K), and growth factors (transforming growth factor- $\beta$ , platelet-derived growth factor, and connective tissue growth factor). Moreover, mitochondrial dysfunction also reportedly contributes to the fibrotic process, but the specific mechanisms of mitochondrial involvement in this setting are unclear.

The use of 5-hydroxyindoleacetic acid, a serotonin metabolite, for patients at risk of diagnosis with carcinoid heart disease, is possible only in individuals with serotonin hypersecretion, within specific diet and laboratory constraints. To this end, there are no specific/sensitive markers to assess the presence, severity, and risk of fibrosis in clinical practice to date.

### *Future Research Priorities*

A better understanding of the pathophysiological mechanisms within the tumor cell and its microenvironment will help delineate the therapeutic targets for NET-associated fibrosis.

Specifically, for carcinoid heart disease, we propose that the priorities should be:

- Establishing a multinational, multidisciplinary working group to study basic, translational, and clinical aspects of carcinoid heart disease by

combining patient databases, imaging records, and biobank samples of blood and tissue.

- Identifying NET-secreted endocrine factors and local cellular and paracrine mediators of carcinoid heart disease in ex vivo and in vitro experimental models.
- Improving clinical risk stratification of carcinoid heart disease by combining patient and tumor characteristics, circulating biomarkers, and imaging features.
- Developing therapeutic strategies for effective identification and prevention of carcinoid heart disease (progression) in high-risk patients with carcinoid syndrome through randomized controlled trials.
- Determining the optimal timing of surgical valve replacement for carcinoid heart disease.

For mesenteric fibrosis, the priorities should be:

- Developing an experimental model that successfully recapitulates the tumor microenvironment.
- Identifying tumor-/patient-specific factors within the tumor microenvironment that determine the development of mesenteric fibrosis.
- Developing a risk assessment tool to identify, which patients with mesenteric fibrosis require preventive surgical resection.
- Discovering novel medical targets to prevent progression of mesenteric fibrosis and testing these in multicenter randomized controlled trials.

#### *Anticipated Impact of Future Research*

Given the paucity of current knowledge on the pathophysiology of NET-associated fibrosis, these future research endeavors will increase our understanding of the most relevant molecular and cellular contributors. Together with superior risk stratification for the development or progression of fibrosis, they will lead to the first interventional clinical trials aimed specifically at treating/preventing NET-associated fibrotic complications. This will ensure high-grade evidence needed to change clinical practice and improve patient outcomes.

#### **Conclusions**

There is no doubt that hypothalamo-pituitary disease remains enigmatic in many of its aspects, ultimately affecting patients' prognoses and outcomes. The many unmet gaps in research that are described above require attention and action from funders, policymakers, healthcare systems, clinicians, and academics, for the benefit of those living with these conditions in the years to come.

Likewise, despite some recent advances in the field of NENs diagnosis and treatment, there is still limited improvement in the real-life approach to these diseases and the related patient outcomes. International funding should focus on promoting further research to fill the gap that exists today between our knowledge and the heterogeneous and still unsolved biological behavior of these rare cancers.

#### **Conflict of Interest Statement**

The authors have no conflicts of interest to declare.

#### **Funding Sources**

No funding was received.

#### **Author Contributions**

Justo Castano (investigation [equal], methodology [equal], project administration [equal], writing – original draft [equal], writing – review and editing [equal]); Mehul Tulsidas Dattani (investigation [equal], methodology [equal], project administration [equal], writing – original draft [equal], writing – review and editing [equal]); Simona Grozinsky-Glasberg (investigation [equal], methodology [equal], project administration [equal], writing – original draft [equal], writing – review and editing [equal]); Niki Karavitaki (investigation [equal], methodology [equal], project administration [equal], writing – original draft [equal], writing – review and editing [equal]); Marianne E. Pavel (investigation [equal], methodology [equal], project administration [equal], writing – original draft [equal], writing – review and editing [equal]); Cynthia Andoniadou (writing – review and editing [equal]). Krystallenia Alexandraki (writing – review and editing [equal]); Cristina Capatina (investigation [equal], writing – original draft [equal]); Manuela Cerbone (investigation [equal], writing – original draft [equal]); Diego Ferone (investigation [equal], writing – original draft [equal]); Hoong Wei Gan (investigation [equal], writing – original draft [equal]); Johannes Hofland (investigation [equal], writing – original draft [equal]); Leo J. Hofland (investigation [equal], writing – original draft [equal]); Alejandro Ibanez-Costa (investigation [equal], writing – original draft [equal], writing – review and editing [equal]); Mirela Diana Ilie (writing – review and editing [equal]); Andrea M. Isidori (investigation [equal], writing – original draft [equal]); Marta Korbonits (investigation [equal], writing – original draft [equal]); Beata Kos-Kudla (investigation [equal], writing – original draft [equal]); Mohamad Maghnie (investigation [equal], writing – original draft [equal]); Giovanna Mantovani (investigation [equal], writing – original draft [equal]); Monica Marazuela (investigation [equal], writing – original draft [equal]); Gerald Raverot (investigation [equal], writing – original draft [equal]); Aldo Scarpa (investigation [equal], writing – original draft [equal]); Katharina Schilbach (investigation [equal], writing – original draft [equal]); Marily Theodoropoulou (investigation [equal], writing – original draft [equal]); Hanneke van Santen (investigation [equal], writing – original draft [equal]); and Maria Chiara Zatelli (Investigation [equal], Writing – original draft [equal]).

## References

- White BE, Rous B, Chandrakumar K, Wong K, Bouvier C, Van Hemelrijck M, et al. Incidence and survival of neuroendocrine neoplasia in England 1995–2018: a retrospective, population-based study. *Lancet Reg Health Eur.* 2022;23:100510. <https://doi.org/10.1016/j.lanepe.2022.100510>
- Chauhan A, Kohn E, Del Rivero J. Neuroendocrine tumors-less well known, often misunderstood, and rapidly growing in incidence. *JAMA Oncol.* 2020;6(1):21–2. <https://doi.org/10.1001/jamaoncol.2019.4568>
- Grozinsky-Glasberg S, Grossman AB, Korbonits M. The role of somatostatin analogues in the treatment of neuroendocrine tumours. *Mol Cell Endocrinol.* 2008;286(1–2):238–50. <https://doi.org/10.1016/j.mce.2007.10.006>
- Oberg K, Modlin IM, De Herder W, Pavel M, Klimstra D, Frilling A, et al. Consensus on biomarkers for neuroendocrine tumour disease. *Lancet Oncol.* 2015;16(9):e435–46. [https://doi.org/10.1016/S1470-2045\(15\)00186-2](https://doi.org/10.1016/S1470-2045(15)00186-2)
- Ho KKY, Fleseriu M, Wass J, Katznelson L, Raverot G, Little AS, et al. A proposed clinical classification for pituitary neoplasms to guide therapy and prognosis. *Lancet Diabetes Endocrinol.* 2024;12(3):209–14. [https://doi.org/10.1016/S2213-8587\(23\)00382-0](https://doi.org/10.1016/S2213-8587(23)00382-0)
- Korbonits M, Blair JC, Boguslawska A, Ayuk J, Davies JH, Druce MR, et al. Consensus guideline for the diagnosis and management of pituitary adenomas in childhood and adolescence: part 1, general recommendations. *Nat Rev Endocrinol.* 2024;20(5):278–89. <https://doi.org/10.1038/s41574-023-00948-8>
- Burman P, Casar-Borota O, Perez-Rivas LG, Dekkers OM. Aggressive pituitary tumors and pituitary carcinomas: from pathology to treatment. *J Clin Endocrinol Metab.* 2023;108(7):1585–601. <https://doi.org/10.1210/clinem/dgad098>
- Theodoropoulou M, Petersenn S, Chanson P, Raverot G. Evidence for somatic mutation screening on aggressive prolactinomas. *Nat Rev Endocrinol.* 2024;20(10):565–6. <https://doi.org/10.1038/s41574-024-01010-x>
- Neou M, Villa C, Armignacco R, Jouinot A, Raffin-Sanson ML, Septier A, et al. Pan-genomic classification of pituitary neuroendocrine tumors. *Cancer Cell.* 2020;37(1):123–34.e5. <https://doi.org/10.1016/j.ccell.2019.11.002>
- Marrero-Rodríguez D, Vela-Patiño S, Martínez-Mendoza F, Valenzuela-Perez A, Peña-Martínez E, Cano-Zaragoza A, et al. Genomics, transcriptomics, and epigenetics of sporadic pituitary tumors. *Arch Med Res.* 2023;54(8):102915. <https://doi.org/10.1016/j.arcmed.2023.102915>
- Angelousi A, Alexandraki KI, Mytareli C, Grossman AB, Kaltsas G. New developments and concepts in the diagnosis and management of diabetes insipidus (AVP-deficiency and resistance). *J Neuroendocrinol.* 2023;35(1):e13233. <https://doi.org/10.1111/jne.13233>
- Spiess M, Beuret N, Rutishauser J. Genetic forms of neurohypophysal diabetes insipidus. *Best Pract Res Clin Endocrinol Metab.* 2020;34(5):101432. <https://doi.org/10.1016/j.beem.2020.101432>
- Ilie MD, Vasiljevic A, Bertolino P, Raverot G. Biological and therapeutic implications of the tumor microenvironment in pituitary adenomas. *Endocr Rev.* 2023;44(2):297–311. <https://doi.org/10.1210/endo/bnac024>
- Ilie MD, Vasiljevic A, Raverot G, Bertolino P. The microenvironment of pituitary tumors-biological and therapeutic implications. *Cancers.* 2019;11(10):1605. <https://doi.org/10.3390/cancers11101605>
- Marques P, Barry S, Carlsen E, Collier D, Ronaldson A, Grieve J, et al. The expression of neural cell adhesion molecule and the microenvironment of pituitary neuroendocrine tumours. *J Neuroendocrinol.* 2021;33(12):e13052. <https://doi.org/10.1111/jne.13052>
- Marques P, Grossman AB, Korbonits M. The tumour microenvironment of pituitary neuroendocrine tumours. *Front Neuroendocrinol.* 2020;58:100852. <https://doi.org/10.1016/j.yfrne.2020.100852>
- Marques P, Silva AL, López-Presa D, Faria C, Bugalho MJ. The microenvironment of pituitary adenomas: biological, clinical and therapeutic implications. *Pituitary.* 2022;25(3):363–82. <https://doi.org/10.1007/s11102-022-01211-5>
- Papadimitriou E, Chatzellis E, Dimitriadi A, Kaltsas GA, Theocharis S, Alexandraki KI. Prognostic biomarkers in pituitary tumours: a systematic review. *Touchrev Endocrinol.* 2023;19(2):42–53. <https://doi.org/10.17925/EE.2023.19.2.12>
- Souteiro P, Karavitaki N. Dopamine agonist resistant prolactinomas: any alternative medical treatment? *Pituitary.* 2020;23(1):27–37. <https://doi.org/10.1007/s11102-019-00987-3>
- Petersenn S, Fleseriu M, Casanueva FF, Giustina A, Biermasz N, Biller BMK, et al. Diagnosis and management of prolactin-secreting pituitary adenomas: a pituitary Society International consensus statement. *Nat Rev Endocrinol.* 2023;19(12):722–40. <https://doi.org/10.1038/s41574-023-00886-5>
- Urwiler SA, Karavitaki N. Refractory lactotroph adenomas. *Pituitary.* 2023;26(3):273–7. <https://doi.org/10.1007/s11102-023-01305-8>
- Ntali G, Wass JA. Epidemiology, clinical presentation and diagnosis of non-functioning pituitary adenomas. *Pituitary.* 2018;21(2):111–8. <https://doi.org/10.1007/s11102-018-0869-3>
- Tampourlou M, Fountas A, Ntali G, Karavitaki N. Mortality in patients with non-functioning pituitary adenoma. *Pituitary.* 2018;21(2):203–7. <https://doi.org/10.1007/s11102-018-0863-9>
- Tampourlou M, Ntali G, Ahmed S, Arlt W, Ayuk J, Byrne JV, et al. Outcome of non-functioning pituitary adenomas that regrow after primary treatment: a study from two large UK centers. *J Clin Endocrinol Metab.* 2017;102(6):1889–97. <https://doi.org/10.1210/jc.2016-4061>
- Dekkers OM, Karavitaki N, Pereira AM. The epidemiology of aggressive pituitary tumors (and its challenges). *Rev Endocr Metab Disord.* 2020;21(2):209–12. <https://doi.org/10.1007/s11154-020-09556-7>
- Raverot G, Ilie MD, Lasolle H, Amodru V, Trouillas J, Castinetti F, et al. Aggressive pituitary tumours and pituitary carcinomas. *Nat Rev Endocrinol.* 2021;17(11):671–84. <https://doi.org/10.1038/s41574-021-00550-w>
- Ilie MD, Raverot G. Treatment options for gonadotroph tumors: current state and perspectives. *J Clin Endocrinol Metab.* 2020;105(10):dgaa497. <https://doi.org/10.1210/clinem/dgaa497>
- Greenman Y, Cooper O, Yaish I, Robenshtok E, Sagiv N, Jonas-Kimchi T, et al. Treatment of clinically nonfunctioning pituitary adenomas with dopamine agonists. *Eur J Endocrinol.* 2016;175(1):63–72. <https://doi.org/10.1530/EJE-16-0206>
- Fleseriu M, Auchus R, Bancos I, Ben-Shlomo A, Bertherat J, Biermasz NR, et al. Consensus on diagnosis and management of Cushing's disease: a guideline update. *Lancet Diabetes Endocrinol.* 2021;9(12):847–75. [https://doi.org/10.1016/S2213-8587\(21\)00235-7](https://doi.org/10.1016/S2213-8587(21)00235-7)
- Simon J, Theodoropoulou M. Genetics of Cushing's disease. *J Neuroendocrinol.* 2022;34(8):13148. <https://doi.org/10.1111/jne.13148>
- Theodoropoulou M, Reincke M. Tumor-directed therapeutic targets in cushing disease. *J Clin Endocrinol Metab.* 2019;104(3):925–33. <https://doi.org/10.1210/jc.2018-02080>
- Melmed S. Pituitary-tumor endocrinopathies. *N Engl J Med.* 2020;382(10):937–50. <https://doi.org/10.1056/NEJMra1810772>
- Melmed S, Kaiser UB, Lopes MB, Bertherat J, Syro LV, Raverot G, et al. Clinical biology of the pituitary adenoma. *Endocr Rev.* 2022;43(6):1003–37. <https://doi.org/10.1210/endo/bnac010>
- Micko AS, Wöhrer A, Wolfsberger S, Knosp E. Invasion of the cavernous sinus space in pituitary adenomas: endoscopic verification and its correlation with an MRI-based classification. *J Neurosurg.* 2015;122(4):803–11. <https://doi.org/10.3171/2014.12.JNS141083>

- 35 Seriola S, Doglietto F, Fiorindi A, Biroli A, Mattavelli D, Buffoli B, et al. Pituitary adenomas and invasiveness from anatomo-surgical, radiological, and histological perspectives: a systematic literature review. *Cancers*. 2019;11(12):1936. <https://doi.org/10.3390/cancers11121936>
- 36 Trouillas J, Roy P, Sturm N, Dantony E, Cortet-Rudelli C, Viennet G, et al. A new prognostic clinicopathological classification of pituitary adenomas: a multicentric case-control study of 410 patients with 8 years post-operative follow-up. *Acta Neuropathol*. 2013;126(1):123–35. <https://doi.org/10.1007/s00401-013-1084-y>
- 37 Raymond P, Raverot G, Ilie MD. Outcome and prognostic factors for pituitary carcinomas: lessons from a systematic review. *Endocr Relat Cancer*. 2023;30(5):220338. <https://doi.org/10.1530/ERC-22-0338>
- 38 Gregory LC, Dattani MT. The molecular basis of congenital hypopituitarism and related disorders. *J Clin Endocrinol Metab*. 2020;105(6):dgaz184. <https://doi.org/10.1210/clinem/dgz184>
- 39 Gregory LC, Cionna C, Cerbone M, Dattani MT. Identification of genetic variants and phenotypic characterization of a large cohort of patients with congenital hypopituitarism and related disorders. *Genet Med*. 2023;25(9):100881. <https://doi.org/10.1016/j.gim.2023.100881>
- 40 Cerbone M, Güemes M, Wade A, Improda N, Dattani M. Endocrine morbidity in midline brain defects: differences between septo-optic dysplasia and related disorders. *eClinicalMedicine*. 2020;19:100224. <https://doi.org/10.1016/j.eclinm.2019.11.017>
- 41 Gan HW, Phipps K, Aquilina K, Gaze MN, Hayward R, Spoudeas HA. Neuroendocrine morbidity after pediatric optic gliomas: a longitudinal analysis of 166 children over 30 years. *J Clin Endocrinol Metab*. 2015;100(10):3787–99. <https://doi.org/10.1210/jc.2015-2028>
- 42 Beckhaus J, Friedrich C, Boekhoff S, Calaminus G, Bison B, Eveslage M, et al. Outcome after pediatric craniopharyngioma: the role of age at diagnosis and hypothalamic damage. *Eur J Endocrinol*. 2023;188(3):lvad027. <https://doi.org/10.1093/ejendo/lvad027>
- 43 Gan HW, Morillon P, Albanese A, Aquilina K, Chandler C, Chang YC, et al. National UK guidelines for the management of paediatric craniopharyngioma. *Lancet Diabetes Endocrinol*. 2023;11(9):694–706. [https://doi.org/10.1016/S2213-8587\(23\)00162-6](https://doi.org/10.1016/S2213-8587(23)00162-6)
- 44 Cerbone M, Visser J, Bulwer C, Ederies A, Vallabhaneni K, Ball S, et al. Management of children and young people with idiopathic pituitary stalk thickening, central diabetes insipidus, or both: a National clinical practice consensus guideline. *Lancet Child Adolesc Health*. 2021;5(9):662–76. [https://doi.org/10.1016/S2352-4642\(21\)00088-2](https://doi.org/10.1016/S2352-4642(21)00088-2)
- 45 van Santen HM, van Schaik J, van Roessel IMAA, Beckhaus J, Boekhoff S, Müller HL. Diagnostic criteria for the hypothalamic syndrome in childhood. *Eur J Endocrinol*. 2023;188(2):lvad009. <https://doi.org/10.1093/ejendo/lvad009>
- 46 Gan HW, Cerbone M, Dattani MT. Appetite- and weight-regulating neuroendocrine circuitry in hypothalamic obesity. *Endocr Rev*. 2024;45(3):309–42. <https://doi.org/10.1210/andrev/bnad033>
- 47 van Iersel L, Brokke KE, Adan RAH, Bulhuis LCM, van den Akker ELT, van Santen HM. Pathophysiology and individualized treatment of hypothalamic obesity following craniopharyngioma and other suprasellar tumors: a systematic review. *Endocr Rev*. 2019;40(1):193–235. <https://doi.org/10.1210/er.2018-00017>
- 48 Müller HL, Tauber M, Lawson EA, Özyurt J, Bison B, Martínez-Barbera JP, et al. Hypothalamic syndrome. *Nat Rev Dis Primers*. 2022;8(1):24. <https://doi.org/10.1038/s41572-022-00351-z>
- 49 Kasuki L, Wildemberg LE, Gadelha MR. Management of endocrine disease: personalized medicine in the treatment of acromegaly. *Eur J Endocrinol*. 2018;178(3):R89–100. <https://doi.org/10.1530/EJE-17-1006>
- 50 Bi WL, Dunn IF, Laws ER. Image guidance and visualization in pituitary surgery. In: Golby AJ, editor. *Image-guided neurosurgery*. Boston, MA, USA: Academic Press; 2015. p. 245–61. [Chapter 10].
- 51 Gil J, Marques-Pamies M, Sampedro M, Webb SM, Serra G, Salinas I, et al. Data mining analyses for precision medicine in acromegaly: a proof of concept. *Sci Rep*. 2022;12(1):8979. <https://doi.org/10.1038/s41598-022-12955-2>
- 52 Lucas JW, Zada G. Endoscopic surgery for pituitary tumors. *Neurosurg Clin N Am*. 2012;23(4):555–69. <https://doi.org/10.1016/j.nec.2012.06.008>
- 53 Mortini P, Albano L, Barzaghi LR, Losa M. Pituitary surgery. *Presse Med*. 2021;50(4):104079. <https://doi.org/10.1016/j.lpm.2021.104079>
- 54 Zhu Z, Cui W, Zhu D, Gao N, Zhu Y. Common tools for pituitary adenomas research: cell lines and primary cells. *Pituitary*. 2020;23(2):182–8. <https://doi.org/10.1007/s11102-019-01003-4>
- 55 Albani A, Perez-Rivas LG, Tang S, Simon J, Lucia KE, Colón-Bolea P, et al. Improved pasireotide response in USP8 mutant corticotroph tumours in vitro. *Endocr Relat Cancer*. 2022;29(8):503–11. <https://doi.org/10.1530/ERC-22-0088>
- 56 Vázquez-Borrego MC, Gupta V, Ibáñez-Costa A, Gahete MD, Venegas-Moreno E, Toledano-Delgado Á, et al. A somatostatin receptor subtype-3 (SST3) peptide agonist shows antitumor effects in experimental models of nonfunctioning pituitary tumors. *Clin Cancer Res*. 2020;26(4):957–69. <https://doi.org/10.1158/1078-0432.CCR-19-2154>
- 57 Laporte E, Vankelecom H. Organoid models of the pituitary gland in health and disease. *Front Endocrinol*. 2023;14:1233714. <https://doi.org/10.3389/fendo.2023.1233714>
- 58 Cui R, Duan H, Hu W, Li C, Zhong S, Liang L, et al. Establishment of human pituitary neuroendocrine tumor derived organoid and its pilot application for drug screening. *J Clin Endocrinol Metab*. 2025;110(3):e827–40. <https://doi.org/10.1210/clinem/dgae228>
- 59 Miwata T, Suga H, Kawaguchi Y, Sakakibara M, Kano M, Taga S, et al. Generation of hypothalamic neural stem cell-like cells in vitro from human pluripotent stem cells. *Stem Cell Rep*. 2023;18(4):869–83. <https://doi.org/10.1016/j.stemcr.2023.02.006>
- 60 Gahete MD, Jiménez-Vacas JM, Alors-Pérez E, Herrero-Aguayo V, Fuentes-Fayos AC, Pedraza-Arévalo S, et al. Mouse models of endocrine tumors. *J Endocrinol*. 2019;240(3):R73–96. <https://doi.org/10.1530/JOE-18-0571>
- 61 Park SR, Kook MG, Kim SR, Lee JW, Yu YS, Park CH, et al. A microscale 3D organ on a chip for recapitulating reciprocal neuroendocrine crosstalk between the hypothalamus and the pituitary gland. *Biofabrication*. 2024;16(2):025011. <https://doi.org/10.1088/1758-5090/ad22f1>
- 62 Maroufi SF, Doğruel Y, Pour-Rashidi A, Kohli GS, Parker CT, Uchida T, et al. Current status of artificial intelligence technologies in pituitary adenoma surgery: a scoping review. *Pituitary*. 2024;27(2):91–128. <https://doi.org/10.1007/s11102-023-01369-6>
- 63 Wildemberg LE, da Silva Camacho AH, Miranda RL, Elias PCL, de Castro Musolino NR, Nazato D, et al. Machine learning-based prediction model for treatment of acromegaly with first-generation somatostatin receptor ligands. *J Clin Endocrinol Metab*. 2021;106(7):2047–56. <https://doi.org/10.1210/clinem/dgab125>
- 64 Klöppel G. Neuroendocrine neoplasms: dichotomy, origin and classifications. *Visc Med*. 2017;33(5):324–30. <https://doi.org/10.1159/000481390>
- 65 Rindi G, Mete O, Uccella S, Basturk O, La Rosa S, Broens LAA, et al. Overview of the 2022 WHO classification of neuroendocrine neoplasms. *Endocr Pathol*. 2022;33(1):115–54. <https://doi.org/10.1007/s12022-022-09708-2>
- 66 Prisciandaro M, Antista M, Raimondi A, Corti F, Morano F, Centonze G, et al. Biomarker landscape in neuroendocrine tumors with high-grade features: current knowledge and future perspective. *Front Oncol*. 2022;12:780716. <https://doi.org/10.3389/fonc.2022.780716>
- 67 Mafficini A, Scarpa A. Genetics and epigenetics of gastroenteropancreatic neuroendocrine neoplasms. *Endocr Rev*. 2019;40(2):506–36. <https://doi.org/10.1210/er.2018-00160>

- 68 English KA, Thakker RV, Lines KE. The role of DNA methylation in human pancreatic neuroendocrine tumours. *Endocr Oncol.* 2023;3(1):e230003. <https://doi.org/10.1530/EO-23-0003>
- 69 Webster AP, Thirlwell C. The molecular biology of midgut neuroendocrine neoplasms. *Endocr Rev.* 2024;45(3):343–50. <https://doi.org/10.1210/endrev/bnad034>
- 70 Matsumoto R, Yamamoto T. Single-cell and spatial transcriptomics in endocrine research. *Endocr J.* 2024;71(2):101–18. <https://doi.org/10.1507/endocrj.EJ23-0457>
- 71 Sedlack AJH, Saleh-Anaraki K, Kumar S, Ear PH, Lines KE, Roper N, et al. Preclinical models of neuroendocrine neoplasia. *Cancers.* 2022;14(22):5646. <https://doi.org/10.3390/cancers14225646>
- 72 Dayton TL, Alcalá N, Moonen L, den Hartigh L, Geurts V, Mangiante L, et al. Druggable growth dependencies and tumor evolution analysis in patient-derived organoids of neuroendocrine neoplasms from multiple body sites. *Cancer Cell.* 2023;41(12):2083–99.e9. <https://doi.org/10.1016/j.ccell.2023.11.007>
- 73 Dijkstra KK, van den Berg JG, Weeber F, van de Haar J, Velds A, Kaing S, et al. Patient-derived organoid models of human neuroendocrine carcinoma. *Front Endocrinol.* 2021;12:627819. <https://doi.org/10.3389/fendo.2021.627819>
- 74 Hwangbo H, Chae S, Kim W, Jo S, Kim GH. Tumor-on-a-chip models combined with mini-tissues or organoids for engineering tumor tissues. *Theranostics.* 2024;14(1):33–55. <https://doi.org/10.7150/thno.90093>
- 75 Mosalem O, Sonbol MB, Halfdanarson TR, Starr JS. Tyrosine kinase inhibitors and immunotherapy updates in neuroendocrine neoplasms. *Best Pract Res Clin Endocrinol Metab.* 2023;37(5):101796. <https://doi.org/10.1016/j.beem.2023.101796>
- 76 Kaur J, Vijayvergia N. Narrative review of immunotherapy in gastroentero-pancreatic neuroendocrine neoplasms. *Curr Oncol.* 2023;30(9):8653–64. <https://doi.org/10.3390/curroncol30090627>
- 77 Boilève A, Faron M, Fodil-Cherif S, Bayle A, Lamartina L, Planchard D, et al. Molecular profiling and target actionability for precision medicine in neuroendocrine neoplasms: real-world data. *Eur J Cancer.* 2023;186:122–32. <https://doi.org/10.1016/j.ejca.2023.03.024>
- 78 Tögel L, Schubart C, Lettmaier S, Neufert C, Hoyer J, Wolff K, et al. Determinants affecting the clinical implementation of a molecularly informed molecular tumor board recommendation: experience from a tertiary cancer center. *Cancers.* 2023;15(24):5892. <https://doi.org/10.3390/cancers15245892>
- 79 Grozinsky-Glasberg S, Davar J, Hofland J, Dobson R, Prasad V, Pascher A, et al. European Neuroendocrine Tumor Society (ENETS) 2022 guidance paper for carcinoid syndrome and carcinoid heart disease. *J Neuroendocrinol.* 2022;34(7):e13146. <https://doi.org/10.1111/jne.13146>
- 80 Beaumont JL, Cella D, Phan AT, Choi S, Liu Z, Yao JC. Comparison of health-related quality of life in patients with neuroendocrine tumors with quality of life in the general US population. *Pancreas.* 2012;41(3):461–6. <https://doi.org/10.1097/MPA.0b013e3182328045>
- 81 Halperin DM, Shen C, Dasari A, Xu Y, Chu Y, Zhou S, et al. Frequency of carcinoid syndrome at neuroendocrine tumour diagnosis: a population-based study. *Lancet Oncol.* 2017;18(4):525–34. [https://doi.org/10.1016/S1470-2045\(17\)30110-9](https://doi.org/10.1016/S1470-2045(17)30110-9)
- 82 Mulders MCF, de Herder WW, Hofland J. What is carcinoid syndrome? A critical appraisal of its proposed mediators. *Endocr Rev.* 2024;45(3):351–60. <https://doi.org/10.1210/endrev/bnad035>
- 83 Grozinsky-Glasberg S, Grossman AB, Gross DJ. Carcinoid heart disease: from pathophysiology to treatment—‘Something in the way it moves. *Neuroendocrinology.* 2015;101(4):263–73. <https://doi.org/10.1159/000381930>