



Human herpes 6 encephalitis in co-infection with Covid-19

A. Di Nora¹ · F. Pizzo¹ · G. Costanza¹ · M. Ruggieri² · R. Falsaperla³

Received: 23 January 2022 / Accepted: 20 April 2022 / Published online: 2 May 2022
© The Author(s) under exclusive licence to Belgian Neurological Society 2022

Keywords COVID-19 · Encephalitis · Herpes virus 6

A 2-year-old-boy presented with ensuing convulsions lasting 40 min (neurological disorders never referred in anamnesis), 3-day history of low-grade fever, erythematous rash on the trunk. Treatment with Midazolam in continuous infusion was started (0.01 mg/kg/h). For the strong suspect of meningoencephalitis, a magnetic resonance imaging (MRI) in emergency was performed, showing an altered signal in temporal lobe (imagine 1), and a lumbar puncture which revealed clear, non-hemorrhagic cerebrospinal fluid (CSF) with 278 cells/ μ L (70% lymphocytes). The real-time polymerase chain reaction (qPCR) assay from a nasopharyngeal swab revealed the presence of the Sars-Cov2 in the child and the mother. Treatment with intravenous acyclovir (10 mg/kg three times a day) and ceftriaxone (70 mg/kg/day) was started. The analysis of the CSF revealed the presence of HSV-6 DNA by PCR. The antibody titer in the blood showed IgM for HSV-6. The immune system was studied revealing that he was an immunocompetent subject. After 2 weeks of treatment with acyclovir, child was discharged in good conditions. After treatment, a control MRI showed a partial disappearance of the strong signal in the temporal lobe precedent described.

In children, primary infection with HSV-6 consists in roseola, a self-limited disease characterized by fever, rash, and pharyngitis. Rare cases of encephalitis to HSV-6 have

been reported in immunocompetent individuals, all infants younger than 3 years [1]. The damage seems related to the indirect effects of cytokine release rather than from direct viral CNS infection [2]. Thus, a co-infection viral may worsen cytokine release and contribute to clinical manifestations.

Many neurological manifestations are described in children affected by Covid-19 and some cases reported encephalitis. Although, most of these reports have had normal neuroimaging. Interestingly, a single study reported a neonatal case affected by seizures and MRI suggestive of viral encephalitis [3].

Diverse and different pathophysiological mechanisms behind neurologic manifestations associated with SARS-CoV-2 infection have been supposed. Some authors proposed an immune-mediated mechanism, reporting Guillain-Barré syndrome or acute disseminated encephalomyelitis [4]. Others supposed the possibility of invasion of Sars-Cov19 because the virus binds the surface spike protein to the human angiotensin-converting enzyme 2 receptor (ACE-2) and ACE-2 is present in the brain vascular endothelium [5]. Differently, others correlate the neurologic features with a severe state of inflammation. Our experience contributes to giving value to this last hypothesis.

Finally, it is interesting observing that multiple studies have been published regarding coinfections with other respiratory pathogens or bacterial infections among patients with COVID-19 [6]. To date, it is the first case of HSV-6 encephalitis in the context of co-infection with Sars-Cov19. In addition, it is a rare study describing MRI abnormality in a child with COVID-19-associated encephalitis (Fig. 1).


✉ A. Di Nora
alessandradinora@gmail.com

¹ Department of Clinical and Experimental Medicine, University of Catania, Via S. Sofia 78, 95123 Catania, Italy

² Unit of Rare Diseases of the Nervous System in Childhood, Department of Clinical and Experimental Medicine, Section of Pediatrics and Child Neuropsychiatry, AOU “Policlinico”, PO “G. Rodolico”, University of Catania, Via S. Sofia, 87, 95128 Catania, Italy

³ Unit of Pediatrics, Neonatology and Neonatal Intensive Care, and Pediatric Emergency, AOU “Policlinico”, PO “San Marco”, University of Catania, Catania, Italy

Fig. 1 MRI of the patient. **a** Axial T2-weighted MR image (left), **b** Coronal T2-weighted MR image (right); images show a hyperintense lesion on the insular cortex with swollen appearance (white arrows)



Acknowledgements We thank the family for taking part in the investigation as well as giving consent for publication of the data and the images.

Author contributions Each author has done a substantial contribution to the conception or design of the work and to revise it critically for important intellectual content. In addition, each author has approved the final of the version to be published. Conversely, each author is agree to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

Funding No funding sources.

Declarations

Conflict of interest The authors have no ethical conflicts to disclose.

References

- Hall CB, Long CE, Schnabel KC et al (1994) Human herpesvirus-6 infection in children. A prospective study of complications and reactivation. *N Engl J Med* 331(4):32–438
- Kawamura Y, Yamazaki Y, Ohashi M, Ihira M, Yoshikawa T (2014) Cytokine and chemokine responses in the blood and cerebrospinal fluid of patients with human herpesvirus 6B-associated acute encephalopathy with biphasic seizures and late reduced diffusion. *J Med Virol* 86:512–518
- Martin PJ, Felker M, Radhakrishnan R (2021) MR imaging findings in a neonate with COVID-19-associated encephalitis. *Pediatr Neurol* 119:48–49
- Frank CHM, Almeida TVR, Marques EA et al (2021) Guillain-Barré syndrome associated with SARS-CoV-2 infection in a pediatric patient. *J Trop Pediatr*. <https://doi.org/10.1093/tropej/fmaa044> (Published online April 9, 2020)
- Zhang H, Penninger JM, Li Y, Zhong N, Slutsky AS (2020) Angiotensin-converting enzyme 2 (ACE2) as a SARS-CoV-2 receptor: molecular mechanisms and potential therapeutic target. *Intensive Care Med* 46(4):586–590
- Cimolai N (2021) The complexity of co-infections in the era of COVID-19. *SN Compr Clin Med* 23:1–13

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.