



## VARIATIONS ON THE CAPACITARY INRADIUS

FRANCESCO BOZZOLA<sup>✉1</sup> AND LORENZO BRASCO<sup>✉\*2</sup>

<sup>1</sup>Dipartimento di Ingegneria Meccanica, Energetica, Gestionale e dei Trasporti,  
Università degli Studi di Genova,  
via all'Opera Pia 15, 16145 Genova, Italy

<sup>2</sup>Dipartimento di Matematica e Informatica,  
Università degli Studi di Ferrara,  
via Machiavelli 30, 44121 Ferrara, Italy

*Remembering Roberto Franceschi, gone too soon*

**ABSTRACT.** We discuss some properties of the *capacitary inradius* for an open set. This is an extension of the classical concept of *inradius* (i.e. the radius of a largest inscribed ball), which takes into account capacitary effects. Its introduction dates back to the pioneering works of Vladimir Maz'ya. We present some variants of this object and their mutual relations, as well as their connections with Poincaré inequalities. We also show that, under a mild regularity assumption on the boundary of the sets, the capacitary inradius is equivalent to the classical inradius. This comes with an explicit estimate and it permits to get a Buser-type inequality for a large class of open sets, whose boundaries may have power-like cusps of arbitrary order. Finally, we present a couple of open problems.

### 1. Introduction.

**1.1. Motivation.** In this note, we pursue our study on the *capacitary inradius* of open subsets of  $\mathbb{R}^N$  that we started in our recent paper [3]. In order to gently introduce the reader to the subject, it is certainly useful to start by recalling the definition of *inradius* of an open set  $\Omega \subseteq \mathbb{R}^N$ : This is the geometric quantity given by

$$r_\Omega = \sup \left\{ r > 0 : \text{there exists a ball } B_r(x_0) \subseteq \Omega \right\}.$$

It can be seen as a simple (and quite rough) measure of “fatness” of an open set. In connection with functional inequalities, its importance is due to the following fact: The condition  $r_\Omega < +\infty$  is *necessary* in order to infer the existence of a constant  $c_\Omega > 0$  such that

$$c_\Omega \int_\Omega |\varphi|^p dx \leq \int_\Omega |\nabla \varphi|^p dx, \quad \text{for every } \varphi \in C_0^\infty(\Omega),$$

where  $1 \leq p < \infty$  (see for example [21, Proposition 2.1]). Thus, the finiteness of the inradius is a necessary condition for the validity of the Poincaré inequality. If

---

2020 *Mathematics Subject Classification.* Primary: 46E35, 35P30, 31C45.

*Key words and phrases.* Poincaré-Sobolev inequality, inradius, capacity, Buser's inequality.

\*Corresponding author: Lorenzo Brasco.

we define the sharp constant in the previous inequality, i.e.,

$$\lambda_p(\Omega) := \inf_{\varphi \in C_0^\infty(\Omega)} \left\{ \int_\Omega |\nabla \varphi|^p dx : \|\varphi\|_{L^p(\Omega)} = 1 \right\},$$

the previous condition can be expressed in a quantitative way through the following well-known sharp estimate:

$$\lambda_p(\Omega) \leq \frac{\lambda_p(B_1)}{r_\Omega^p}.$$

We briefly recall that in general it is not always possible to reverse the previous inequality, unless some further restrictions on the open sets are imposed. We refer to the introductions of our previous papers [2, 3] for a list of results in this direction (see also [1] and the references therein). The crucial obstruction to the validity of a lower bound of the type

$$\frac{C_{N,p}}{r_\Omega^p} \leq \lambda_p(\Omega), \tag{1}$$

is a “removability issue”: Roughly speaking, while  $r_\Omega$  can be altered by the removal of a single point, the Poincaré constant  $\lambda_p(\Omega)$  is only affected by the removal of “sufficiently large” sets. That said, it is quite easy to produce counterexamples to the validity of (1). The previous largeness condition depends on the exponent  $p$  and is expressed in terms of  $p$ -capacity. In this paper, we will mainly work with the so-called *relative  $p$ -capacity*, defined by

$$\text{cap}_p(\Sigma; E) = \inf_{\varphi \in C_0^\infty(E)} \left\{ \int_E |\nabla \varphi|^p dx : \varphi \geq 1 \text{ on } \Sigma \right\},$$

for every open bounded set  $E \subseteq \mathbb{R}^N$  (typically, a ball) and every compact set  $\Sigma \Subset E$ .

Under this premise, it is natural to introduce a variant of  $r_\Omega$  which disregards sets having zero  $p$ -capacity, i.e. we wish to consider a relaxed notion of inradius which shares with  $\lambda_p(\Omega)$  the same removable sets. This is the idea for introducing the *capacitary inradius* of an open set  $\Omega$ , whose precise definition is as follows: For every  $0 \leq \gamma < 1$  and for every  $1 \leq p < \infty$ , this is defined by

$$R_{p,\gamma}(\Omega) := \sup \left\{ r > 0 : \exists x_0 \in \mathbb{R}^N \text{ such that } \overline{B_r(x_0)} \setminus \Omega \text{ is } (p, \gamma)\text{-negligible} \right\}. \tag{2}$$

Here  $(p, \gamma)$ -negligible (in the sense of Molchanov) means that  $\overline{B_r(x_0)} \setminus \Omega$  occupies a portion of a reference concentric ball, say  $B_{2r}(x_0)$ , which is at most  $\gamma$  in the sense of  $p$ -capacity. More precisely, we have

$$\text{cap}_p \left( \overline{B_r(x_0)} \setminus \Omega; B_{2r}(x_0) \right) \leq \gamma \text{cap}_p \left( \overline{B_r(x_0)}; B_{2r}(x_0) \right).$$

The quantity (2) was explicitly introduced in our recent paper [3] by drawing inspiration both from the *interior capacitary radius* used by Maz'ya and Shubin in [18] and from the *inner cubic diameter* (see [16, Definition 14.2.2]), which has been extensively used by Maz'ya. We postpone to the following section a more detailed discussion about the comparison between these capacitary variants of the inradius.

**Remark 1.1.** The extremal cases  $\gamma = 1$  and  $\gamma = 0$  deserve a comment: In the first case, every set would be  $(p, \gamma)$ -negligible, and thus the corresponding capacitary inradius would be  $+\infty$  for every open set. The case  $\gamma = 0$  is more interesting: In this case, the quantity  $R_{p,0}(\Omega)$  would give the radius of the largest ball contained

in  $\Omega$ , up to a set of zero  $p$ -capacity. However, this capacity inradius would not be strong enough to permit the lower bound

$$\left(\frac{1}{R_{p,0}(\Omega)}\right)^p \lesssim \lambda_p(\Omega),$$

see [3, Example A.1] for a counter-example.

On the contrary, when  $0 < \gamma < 1$ , one can prove that  $R_{p,\gamma}(\Omega)$  has the desired property: It is actually equivalent to  $\lambda_p(\Omega)$ . More precisely, in [3, Main Theorem], we proved the two-sided estimate

$$\sigma_{N,p} \gamma \left(\frac{1}{R_{p,\gamma}(\Omega)}\right)^p \leq \lambda_p(\Omega) \leq C_{N,p,\gamma} \left(\frac{1}{R_{p,\gamma}(\Omega)}\right)^p, \quad (3)$$

which holds for every  $0 < \gamma < 1$  and within the range  $1 \leq p \leq N$ . Actually, a related result holds for the sharp Poincaré-Sobolev constants (see [3, Theorem 6.1]),

$$\lambda_{p,q}(\Omega) = \inf_{\varphi \in C_0^\infty(\Omega)} \left\{ \int_{\Omega} |\nabla \varphi|^p dx : \|\varphi\|_{L^q(\Omega)} = 1 \right\},$$

each time  $q > p$  is a *subcritical* (in the sense of Sobolev embeddings) exponent.

**Remark 1.2** (The case  $p > N$ ). The regime  $p > N$  is less interesting. In this case, points have positive  $p$ -capacity and thus they are not removable sets. Accordingly, one can prove that (1) holds for every open set<sup>1</sup>. We also mention that in [3, Proposition 7.3], it is shown that  $R_{p,\gamma}(\Omega) = r_\Omega$ , when  $p > N$  and  $0 < \gamma \leq \gamma_0$ , for an explicit (and optimal) parameter  $\gamma_0$ . For these reasons, in this paper we will confine our discussion to the case  $p \leq N$ .

**1.2. Description of the results.** In the first part of the paper, we want to consider some variants of  $R_{p,\gamma}$  and discuss their mutual relations. We will focus especially on two different kinds of variants:

- (A) changing the *metric*, i.e. replacing balls in the definition of  $R_{p,\gamma}$  with a family of rescaled copies of a more general fixed “shape”;
- (B) changing the *capacity*, i.e. replacing the relative  $p$ -capacity with other notions of  $p$ -capacity.

The first question may look a bit academic, but actually it is not: Indeed, the first appearance of the capacity inradius can be traced back in some works by Maz’ya from the 1970s, as summarized in his books [16, 17]. In these works, the capacity inradius is defined by using *cubes* in place of *balls*, which is precisely the *inner cubic diameter* mentioned above (see [17, Definition 10.2.2] and [16, Definition 14.2.2]). Crucially, this is exploited in order to derive necessary and sufficient conditions for the validity of Poincaré and Poincaré-Sobolev inequalities on open subsets of  $\mathbb{R}^N$ . The use of cubes has the advantage that they can tile the space, differently from balls. On the other hand, by using balls, it is simpler to prove a two-sided estimate like (3) valid for every  $0 < \gamma < 1$ , as shown in [18] (for  $p = 2 < N$ ) and in our paper [3] (general case). Thus, it is natural to inquire to which extent the shape of the chosen “ball” can affect this type of estimates. It should also be remarked that

<sup>1</sup>This result has been proved and re-proved various times, with slightly different proofs and different estimates of the constant  $C_{N,p}$ . However, it seems correct to attribute it to Maz’ya, see for example [17, Theorem 11.4.1].

this kind of generalization has already been considered in [16, Chapter 18] (for the case  $p = 2$ ).

As for point (B), we remark that we do not perversely consider any possible variant one could imagine: Rather, our interest is to consider a couple of alternative notions already existing in the literature. The first one is what we call the *Maz'ya-Shubin capacitary inradius*, used in [18], a paper which is very much inspired [3]. By referring to Definition 4.6 for its precise form, we point out that the main difference with  $R_{p,\gamma}$  is the use of the following *absolute  $p$ -capacity*

$$\text{cap}_p(\Sigma) = \inf_{\varphi \in C_0^\infty(\mathbb{R}^N)} \left\{ \int_{\mathbb{R}^N} |\nabla \varphi|^p dx : \varphi \geq 1 \text{ on } \Sigma \right\}.$$

As already explained in the introduction of [3], the main drawback of this notion is that it does not permit to consider the limit case  $p = N$ .

The second variant is the *Gallagher capacitary inradius*, recently considered in [10, 11]. Apart from the use of yet another notion of  $p$ -capacity (which permits to include the case  $p = N$ , as well), the main difference is that this inradius is *not* defined in terms of a negligibility condition *à la* Molchanov. Rather, it can be regarded as a sort of “endpoint”, i.e. it coincides with the limit of  $R_{p,\gamma}$  as the negligibility parameter  $\gamma$  goes to 0 (see Proposition 4.15). This has already been observed by Gallagher in [11]. Thus, even if this inradius can be used to characterize the validity of the Poincaré inequality, it *cannot* be directly used to provide two-sided estimates like (3) (more precisely, the lower bound fails, see Example 4.17).

The second part of the paper is focused on answering the following natural question: *Under which conditions on the open set can the capacitary inradius  $R_{p,\gamma}(\Omega)$  be compared with the classical inradius  $r_\Omega$ ?* In Section 5, we will show how a simple *measure density condition* on the open sets imply that

$$r_\Omega \leq R_{p,\gamma}(\Omega) \lesssim r_\Omega, \quad \text{for every } 0 \leq \gamma < 1.$$

Thus, the capacitary inradius  $R_{p,\gamma}(\Omega)$  turns out to be equivalent to the usual inradius for these sets. This is the content of Theorem 5.2 below. Our measure density condition amounts to requiring

$$\theta_{\Omega,r_0}^*(t) := \inf \left\{ \left( \frac{r_0}{r} \right)^t \frac{|B_r(x) \setminus \Omega|}{|B_r(x)|} : x \in \partial\Omega, 0 < r \leq r_0 \right\} > 0,$$

for some  $r_0 > 0$  and  $t \geq 0$ . Such a condition is not the sharpest possible assumption on the open sets in order to have such an equivalence. However, we think that the result is quite interesting for two reasons: The condition is sufficiently general to encompass a large class of open sets (we admit power-like cusps of arbitrary order for their boundaries), and, at the same time, it is easy to check. We refer to Lemma 5.13 for a large class of open sets which satisfy it. Moreover, the equivalence between  $R_{p,\gamma}(\Omega)$  and  $r_\Omega$  comes with an explicit estimate.

When combined with the lower bound in (3), this estimate in turn permits to infer the validity of (1) for this class of sets, with a precise control on the constant involved. At the same price, we can get the same result also for the Poincaré-Sobolev constants  $\lambda_{p,q}$ , see Corollary 5.6. As a remarkable consequence, we can obtain a *Buser-type inequality* for open sets satisfying the previous measure density condition. More precisely, we get

$$\lambda(\Omega) \leq C \left( h(\Omega) \right)^2, \tag{4}$$

where  $C > 0$  is constant depending only on  $N$  and the measure density index  $\theta_{\Omega, r_0}^*(t)$  (in an explicit way), and:

- $\lambda(\Omega)$  is the bottom of the spectrum of the Dirichlet-Laplacian on  $\Omega$ , that is<sup>2</sup>

$$\lambda(\Omega) := \lambda_2(\Omega) = \inf_{\varphi \in C_0^\infty(\Omega)} \left\{ \int_{\Omega} |\nabla \varphi|^2 dx : \|\varphi\|_{L^2(\Omega)} = 1 \right\};$$

- $h(\Omega)$  is the *Cheeger constant* of  $\Omega$ , defined by

$$h(\Omega) = \inf \left\{ \frac{\mathcal{H}^{N-1}(\partial E)}{|E|} : E \Subset \Omega \text{ open set with smooth boundary} \right\}.$$

We refer to Corollary 5.9 for the precise statement.

We conclude by recalling that for general open sets it is not possible to have an estimate like (4); we refer to [16, Chapter 4, Section 3] for a counter-example and to [2, 5, 19] for some positive results, under suitable topological/geometric assumptions on the open sets. Inequality (4) is sometimes also called the *reverse Cheeger's inequality* (see for example [8, 9, 14] and [15] for this result in the context of Riemannian manifolds). The motivation for this terminology is easily understood, once we recall the following *Cheeger inequality*:

$$\left( \frac{h(\Omega)}{2} \right)^2 \leq \lambda(\Omega), \quad (5)$$

see [16, equation (4.2.5)], which holds for *every* open set  $\Omega \subseteq \mathbb{R}^N$ .

**1.3. Plan of the paper.** In Section 2, we settle the notation and state some basic facts which will be repeatedly used throughout the paper. In Section 3, we compare our capacity inradius with the variant obtained by considering a more general class of test “shapes”, other than balls. An interesting open problem is presented as well. Section 4 is still devoted to comparison estimates with other notions of capacity inradius, this time obtained by considering different capacities. These variants includes the one considered by Maz'ya and Shubin in [18] and the one considered by Gallagher in [10, 11]. This section is complemented with examples and remarks, aimed at clarifying some substantial differences between  $R_{p,\gamma}$  and the inradius considered in [10, 11]. In the final section, i.e. Section 5, we state and prove our main result about the comparison between the capacity inradius and the classical one under a measure density condition (see Theorem 5.2). We also briefly discuss some consequences which can be drawn from it.

**2. Preliminaries.** Unless differently stated, in this paper we will always take the dimension to be  $N \geq 2$ . For a point  $x_0 \in \mathbb{R}^N$  and a positive real number  $R$ , we will indicate by  $B_R(x_0)$  the  $N$ -dimensional open ball, centered at  $x_0$  and with radius  $R$ , i.e.

$$B_R(x_0) = \left\{ x \in \mathbb{R}^N : |x - x_0| < R \right\}.$$

When the center  $x_0$  coincides with the origin, we will simply write  $B_R$ . We indicate by  $\omega_N$  the volume of the  $N$ -dimensional ball having radius 1. We also denote by

---

<sup>2</sup>At the risk of being inconsistent, for the special case  $p = 2$  we prefer to use a distinguished notation for the sharp Poincaré inequality. Indeed, experts in spectral theory could be misled by the notation  $\lambda_2(\Omega)$ .

$Q_R(x_0)$  the  $N$ -dimensional open hypercube centered at  $x_0$ , given by

$$Q_R(x_0) = \prod_{i=1}^N (x_0^i - R, x_0^i + R), \quad \text{where } x_0 = (x_0^1, \dots, x_0^N).$$

Given two non-empty sets  $\Sigma, K \subseteq \mathbb{R}^N$  that are, respectively, closed and compact, we will indicate

$$\text{dist}(\Sigma, K) = \min_{x \in \Sigma, y \in K} |x - y|.$$

Next, we recall the definition of  $p$ -capacity we wish to work with.

**Definition 2.1.** Let  $1 \leq p < \infty$ . For every  $E \subseteq \mathbb{R}^N$  open set and every  $\Sigma \subseteq E$  compact set, we define the  $p$ -capacity of  $\Sigma$  relative to  $E$  as follows

$$\text{cap}_p(\Sigma; E) = \inf_{\varphi \in C_0^\infty(E)} \left\{ \int_E |\nabla \varphi|^p dx : \varphi \geq 1 \text{ on } \Sigma \right\}.$$

It is intended that  $\text{cap}_p(\emptyset; E) = 0$ . By using standard approximation methods, it is not difficult to see that this infimum is unchanged, if we replace  $C_0^\infty(E)$  by the space of Lipschitz functions, compactly supported in  $E$ .

We refer to [16] for a comprehensive study on the properties of the various notions of  $p$ -capacity used in this paper. We also mention the paper [13] for a succinct presentation of the subject, in the case of the relative  $p$ -capacity introduced above.

From its definition, it is easy to see that we have the monotonicity relations

$$\text{cap}_p(\Sigma_0; E) \leq \text{cap}_p(\Sigma_1; E), \quad \text{if } \Sigma_0 \subseteq \Sigma_1 \Subset E, \tag{6}$$

and

$$\text{cap}_p(\Sigma; E_1) \leq \text{cap}_p(\Sigma; E_0), \quad \text{if } \Sigma \Subset E_0 \subseteq E_1,$$

which will be used repeatedly.

We also recall the following explicit formula for the  $p$ -capacity of concentric balls: They can be found, for example, in [16, page 148]. We limit ourselves to the case  $1 \leq p \leq N$ , which will be the most interesting one: For  $0 < r < R$  we have

$$\text{cap}_1(\overline{B_r}; B_R) = N \omega_N r^{N-1}, \tag{7}$$

$$\text{cap}_p(\overline{B_r}; B_R) = N \omega_N \left( \frac{N-p}{p-1} \right)^{p-1} \frac{r^{N-p}}{\left( 1 - \left( \frac{r}{R} \right)^{\frac{N-p}{p-1}} \right)^{p-1}}, \quad \text{if } 1 < p < N, \tag{8}$$

and

$$\text{cap}_N(\overline{B_r}; B_R) = N \omega_N \left( \log \left( \frac{R}{r} \right) \right)^{1-N}. \tag{9}$$

**Remark 2.2.** We explicitly remark that formula (7) holds in the case  $p = N = 1$ , as well. In this case, it reads as

$$\text{cap}_1(\overline{B_r}; B_R) = 2,$$

with  $B_r = (-r, r) \subseteq \mathbb{R}$  and  $B_R = (-R, R) \subseteq \mathbb{R}$ . This is not stated in [16, page 148], but it can be easily obtained as follows: For every  $\varphi \in C_0^\infty(B_R)$  such that  $\varphi \geq 1$  on  $B_r$ , we have

$$\int_{B_R} |\varphi'| dx \geq 2 \|\varphi\|_{L^\infty(B_R)} \geq 2,$$

thanks to the one-dimensional Sobolev inequality (see for example [4, Theorem 3.6.6]). Owing to the arbitrariness of  $\varphi$ , we get  $\text{cap}_1(\overline{B_r}; B_R) \geq 2$ . On the other hand, by using as test functions a sequence of smooth approximations of the characteristic function of  $(-r, r)$ , we get the reverse inequality, as well. The details are standard, and thus omitted.

The next simple result is well-known (see for example [16, Corollary 2.3.4]), we recall its proof for the reader's convenience. It will be crucially exploited in the sequel.

**Lemma 2.3.** *Let  $E \subseteq \mathbb{R}^N$  be an open set and let  $\Sigma \subseteq E$  be a compact set. For every  $1 \leq p < \infty$ , we have*

$$|\Sigma| \lambda_p(E) \leq \text{cap}_p(\Sigma; E).$$

*Proof.* We can suppose that  $\lambda_p(E) > 0$ , otherwise there is nothing to prove. Let  $\varphi \in C_0^\infty(E)$  be such that  $\varphi \geq 1$  on  $\Sigma$ . Then, we have

$$\lambda_p(E) |\Sigma| \leq \lambda_p(E) \int_{\Sigma} |\varphi|^p dx \leq \lambda_p(E) \int_E |\varphi|^p dx \leq \int_E |\nabla \varphi|^p dx.$$

Taking the infimum over  $\varphi$ , we get the claim.  $\square$

The following property immediately follows from the explicit expressions (7), (8), and (9).

**Lemma 2.4.** *Let  $1 \leq p \leq N$ . For every  $\alpha > 1$ , the function*

$$f(r) = \text{cap}_p(\overline{B_r(x_0)}; B_{\alpha r}(x_0)), \quad \text{for } r > 0.$$

*is non-decreasing.*

**3. Different metrics.** Following [16, Definition 18.1], we want to consider a variant of our capacity inradius  $R_{p,\gamma}$ , where we replace balls with more general “shapes”. We first need to define the kind of “shapes” we wish to consider.

We say that an open bounded set  $\mathcal{K} \subseteq \mathbb{R}^N$  is a *standard body* if it is starshaped with respect to the ball  $B_1$ . Such a set  $\mathcal{K}$  can be described as follows:

$$\mathcal{K} = \{x = r\omega \in \mathbb{R}^N : \omega \in \mathbb{S}^{N-1}, 0 \leq r < r(\omega)\},$$

for a Lipschitz function  $r : \mathbb{S}^{N-1} \rightarrow (0, \infty)$ , the so-called *radial function of  $\mathcal{K}$* , given by

$$r(\omega) := \sup\{r \geq 0 : r\omega \in \mathcal{K}\}, \quad \text{for every } \omega \in \mathbb{S}^{N-1},$$

see [16, Lemma 1.1.8]. Moreover, we have

$$r(\omega)\omega \in \partial\mathcal{K} \quad \text{and} \quad 1 \leq r(\omega) \leq R_{\mathcal{K}}, \quad \text{for every } \omega \in \mathbb{S}^{N-1},$$

where we set

$$R_{\mathcal{K}} := \max_{x \in \partial\mathcal{K}} |x| = \max_{\omega \in \mathbb{S}^{N-1}} r(\omega).$$

Observe that we used that  $B_1 \subseteq \mathcal{K}$ , by our assumption.

For every  $r > 0$  and for every  $x_0 \in \mathbb{R}^N$ , we denote by  $\mathcal{K}_r(x_0)$  the open set obtained from  $\mathcal{K}$  by combining a homothety (centered at 0) with coefficient  $r$  and a translation by the vector  $x_0$ , that is

$$\mathcal{K}_r(x_0) := \left\{ x \in \mathbb{R}^N : \frac{x - x_0}{r} \in \mathcal{K} \right\}.$$

When the “center”  $x_0$  coincides with the origin, we will simply write  $\mathcal{K}_r$  in place of  $\mathcal{K}_r(x_0)$ . For  $r = 1$ , we get back the original shape, and thus we will write  $\mathcal{K}$  in place of  $\mathcal{K}_1$ .

This family of sets will replace the balls  $B_r(x_0)$  in the capacity inradius we are going to define. Indeed, we have the following definition, which is essentially contained in [16, Chapter 18]. In light of Remark 1.1, we exclude the extremal case  $\gamma = 1$ , here as well.

**Definition 3.1.** Let  $1 \leq p < \infty$  and  $0 \leq \gamma < 1$ . A compact set  $\Sigma \subseteq \overline{\mathcal{K}_r(x_0)}$  is said to be  $(p, \gamma)$ -negligible relative to  $\mathcal{K}$  if

$$\text{cap}_p(\Sigma; \mathcal{K}_{2r}(x_0)) \leq \gamma \text{cap}_p(\overline{\mathcal{K}_r(x_0)}; \mathcal{K}_{2r}(x_0)).$$

For an open set  $\Omega \subseteq \mathbb{R}^N$ , we define its *capacity inradius relative to  $\mathcal{K}$*  as

$$R_{p,\gamma}(\Omega; \mathcal{K}) := \sup \left\{ r > 0 : \begin{array}{l} \exists x_0 \in \mathbb{R}^N \text{ such that } \overline{\mathcal{K}_r(x_0)} \setminus \Omega \\ \text{is } (p, \gamma)\text{-negligible relative to } \mathcal{K} \end{array} \right\}.$$

A couple of comments are in order concerning the previous definition.

**Remark 3.2.** A particularly interesting instance is when  $\mathcal{K}$  coincides with the unit ball of a norm on  $\mathbb{R}^N$ . For example, by choosing  $\mathcal{K} = B_1$ , the Euclidean ball, we get our initial definition back. On the other hand, by choosing the hypercube

$$\mathcal{K} = Q_1(0) = (-1, 1)^N,$$

i.e. the unit ball of the norm  $\|\cdot\|_{\ell^\infty}$ , the quantity  $R_{p,\gamma}(\Omega; \mathcal{K})$  coincides with the so-called *inner cubic diameter*, extensively used by Maz’ya in his works, as recalled in the introduction.

**Remark 3.3.** We point out that the requirement on the standard body  $\mathcal{K}$  to be starshaped with respect to a ball is needed in order to guarantee that  $\mathcal{K}_r(x_0)$  is compactly contained in  $\mathcal{K}_{2r}(x_0)$ . In this way, the quantity  $\text{cap}_p(\overline{\mathcal{K}_r(x_0)}; \mathcal{K}_{2r}(x_0))$  is well-defined.

**Lemma 3.4.** Let  $1 \leq p < \infty$ , for every non-empty compact set  $\Sigma \subseteq B_\varrho(x_0)$  we have

$$\text{cap}_p(\Sigma; B_\varrho(x_0)) \leq \left( \frac{\varrho}{\text{dist}(\Sigma, \partial B_\varrho(x_0))} \frac{1}{\lambda_p(\mathcal{K})^{\frac{1}{p}}} + 1 \right)^p \text{cap}_p(\Sigma; \mathcal{K}_\varrho(x_0)). \quad (10)$$

*Proof.* We first observe that if  $\Sigma \subseteq B_\varrho(x_0)$ , then we have  $\Sigma \subseteq \mathcal{K}_\varrho(x_0)$ , as well. This simply follows from the fact that

$$B_\varrho(x_0) \subseteq \mathcal{K}_\varrho(x_0),$$

thanks to the fact that the standard body  $\mathcal{K}$  is starshaped with respect to  $B_1$ .

We set for simplicity  $d = \text{dist}(\Sigma, \partial B_\varrho(x_0))$ , and then we have  $\Sigma \subseteq B_{\varrho-d}(x_0)$ . For every  $0 < \delta < d$ , we define the cut-off function

$$\eta_\delta(x) := \min \left\{ \left( \frac{(\varrho - \delta) - |x - x_0|}{d - \delta} \right)_+, 1 \right\}.$$

In particular,  $\eta_\delta$  is a Lipschitz function with compact support in  $B_\varrho(x_0)$ , and

$$\eta_\delta = 1 \quad \text{on } \overline{B_{\varrho-d}(x_0)}, \quad |\nabla \eta_\delta| \leq \frac{1}{d - \delta}, \quad \eta_\delta = 0 \quad \text{on } \mathbb{R}^N \setminus B_{\varrho-\delta}(x_0).$$

We take  $u \in C_0^\infty(\mathcal{K}_\varrho(x_0))$  such that  $u \geq 1$  on  $\Sigma$ . Then, by the definition of relative capacity and Minkowski's inequality, we have

$$\begin{aligned} \left(\text{cap}_p(\Sigma; B_\varrho(x_0))\right)^{\frac{1}{p}} &\leq \|\nabla(u \eta_\delta)\|_{L^p(B_\varrho(x_0))} \\ &\leq \frac{1}{d-\delta} \|u\|_{L^p(\mathcal{K}_\varrho(x_0))} + \|\nabla u\|_{L^p(\mathcal{K}_\varrho(x_0))} \\ &\leq \left(\frac{1}{d-\delta} \frac{1}{\lambda_p(\mathcal{K}_\varrho(x_0))^{\frac{1}{p}}} + 1\right) \|\nabla u\|_{L^p(\mathcal{K}_\varrho(x_0))}, \end{aligned}$$

where in the last line we used the Poincaré inequality on the bounded set  $\mathcal{K}_\varrho(x_0)$ . By sending  $\delta$  to 0 and by the arbitrariness of  $u$ , we then obtain the claimed estimate.  $\square$

The previous result has a sort of converse.

**Lemma 3.5.** *Let  $1 \leq p < \infty$ . For every non-empty compact set  $\Sigma \subseteq \mathcal{K}_\varrho(x_0)$ , we have*

$$\text{cap}_p(\Sigma; \mathcal{K}_\varrho(x_0)) \leq \left(\frac{\varrho}{\text{dist}(\Sigma, \partial\mathcal{K}_\varrho(x_0))} \frac{R_{\mathcal{K}}}{\lambda_p(B_1)^{\frac{1}{p}}} + 1\right)^p \text{cap}_p(\Sigma; B_{R_{\mathcal{K}}\varrho}(x_0)).$$

*Proof.* We first observe that if  $\Sigma \subseteq \mathcal{K}_\varrho(x_0)$ , then we have  $\Sigma \subseteq B_{R_{\mathcal{K}}\varrho}(x_0)$  as well. This simply follows from the fact that

$$\mathcal{K}_\varrho(x_0) \subseteq B_{R_{\mathcal{K}}\varrho}(x_0),$$

just by recalling the definition of  $R_{\mathcal{K}}$ .

Let  $u \in C_0^\infty(B_{R_{\mathcal{K}}\varrho}(x_0))$  be such that  $u \geq 1$  on  $\Sigma$ . For every  $0 < \delta < d := \text{dist}(\Sigma, \partial\mathcal{K}_\varrho(x_0))$ , we introduce the cut-off function  $\xi_\delta$  given by

$$\xi_\delta(x) = \min \left\{ \frac{1}{d-\delta} (\text{dist}(x; \partial\mathcal{K}_\varrho(x_0)) - \delta)_+, 1 \right\}, \quad \text{for } x \in \mathcal{K}_\varrho(x_0).$$

This is a Lipschitz function with compact support in  $\mathcal{K}_\varrho(x_0)$  such that  $\xi_\delta = 1$  on  $\Sigma$ . Thus, by the definition of relative capacity and Minkowski's inequality, we obtain

$$\begin{aligned} \left(\text{cap}_p(\Sigma; \mathcal{K}_\varrho(x_0))\right)^{\frac{1}{p}} &\leq \|\nabla(u \xi_\delta)\|_{L^p(\mathcal{K}_\varrho(x_0))} \\ &\leq \frac{1}{d-\delta} \|u\|_{L^p(B_{R_{\mathcal{K}}\varrho}(x_0))} + \|\nabla u\|_{L^p(B_{R_{\mathcal{K}}\varrho}(x_0))} \\ &\leq \left(\frac{\varrho}{d-\delta} \frac{R_{\mathcal{K}}}{\lambda_p(B_1)^{\frac{1}{p}}} + 1\right) \|\nabla u\|_{L^p(B_{R_{\mathcal{K}}\varrho}(x_0))}, \end{aligned}$$

where in the last line we used Poincaré inequality for  $B_{R_{\mathcal{K}}\varrho}(x_0)$ . By the arbitrariness of  $u$  and  $\delta$ , we then have the desired estimate.  $\square$

We have the following result, which compares the quantities  $R_{p,\gamma}(\Omega; \mathcal{K})$  and  $R_{p,\gamma}(\Omega)$ . This is a generalization of [1, Proposition 2.2.5], contained in the Ph.D. thesis of the second author.

**Proposition 3.6.** *Let  $1 \leq p \leq N$  and let  $\Omega \subseteq \mathbb{R}^N$  be an open set. For every standard body  $\mathcal{K} \subseteq \mathbb{R}^N$ , there exist two constants  $0 < c \leq 1 \leq d$ , both depending only on  $N$ ,  $p$ , and  $\mathcal{K}$ , such that we have*

$$R_{p,c,\gamma}(\Omega; \mathcal{K}) \leq R_{p,\gamma}(\Omega), \quad \text{for every } 0 \leq \gamma < 1, \quad (11)$$

and

$$R_{p,\gamma}(\Omega) \leq R_{\mathcal{K}} R_{p,d,\gamma}(\Omega; \mathcal{K}), \quad \text{for every } 0 \leq \gamma < \frac{1}{d}, \quad (12)$$

where  $R_{\mathcal{K}} = \max_{x \in \partial \mathcal{K}} |x|$ .

*Proof.* We prove these two estimates separately.

*Proof of (11).* We can suppose that  $R_{p,\gamma}(\Omega) < +\infty$ , otherwise the leftmost inequality in (11) is trivial. We set

$$c = c(N, p, \mathcal{K}) := \frac{\text{cap}_p(\overline{B_1}; B_2)}{\text{cap}_p(\overline{\mathcal{K}}; \mathcal{K}_2)} \left( \frac{2}{\lambda_p(\mathcal{K})^{\frac{1}{p}}} + 1 \right)^{-p}, \quad (13)$$

and observe that  $c \leq 1$ , by Lemma 3.4 with  $x_0 = 0$ ,  $\Sigma = \overline{B_1}$ , and  $\varrho = 2$  in conjunction with (6). Let  $r > R_{p,\gamma}(\Omega)$ . By the definition of capacity inradius, we then have

$$\text{cap}_p(\overline{B_r(x_0)} \setminus \Omega; B_{2r}(x_0)) > \gamma \text{cap}_p(\overline{B_r(x_0)}; B_{2r}(x_0)), \quad \text{for every } x_0 \in \mathbb{R}^N, \quad (14)$$

By using (10) with  $\Sigma = \overline{B_r(x_0)} \setminus \Omega$  and  $\varrho = 2r$ , we also get

$$\begin{aligned} \text{cap}_p(\overline{B_r(x_0)} \setminus \Omega; B_{2r}(x_0)) &\leq \left( \frac{2}{\lambda_p(\mathcal{K})^{\frac{1}{p}}} + 1 \right)^p \text{cap}_p(\overline{B_r(x_0)} \setminus \Omega; \mathcal{K}_{2r}(x_0)), \\ &\leq \left( \frac{2}{\lambda_p(\mathcal{K})^{\frac{1}{p}}} + 1 \right)^p \text{cap}_p(\overline{\mathcal{K}_r(x_0)} \setminus \Omega; \mathcal{K}_{2r}(x_0)). \end{aligned}$$

Observe that we also used (6) thanks to the fact that  $\overline{B_r(x_0)} \setminus \Omega \subseteq \overline{\mathcal{K}_r(x_0)} \setminus \Omega$ . This, combined with (14) gives that

$$\gamma \left( \frac{2}{\lambda_p(\mathcal{K})^{\frac{1}{p}}} + 1 \right)^{-p} \text{cap}_p(\overline{B_r(x_0)}; B_{2r}(x_0)) \leq \text{cap}_p(\overline{\mathcal{K}_r(x_0)} \setminus \Omega; \mathcal{K}_{2r}(x_0)).$$

On the other hand, by the scaling properties of the relative capacity, we have

$$\text{cap}_p(\overline{B_r(x_0)}; B_{2r}(x_0)) = \frac{\text{cap}_p(\overline{B_1}; B_2)}{\text{cap}_p(\overline{\mathcal{K}}; \mathcal{K}_2)} \text{cap}_p(\overline{\mathcal{K}_r(x_0)}; \mathcal{K}_{2r}(x_0)).$$

Thus, by recalling the definition (13) of  $c$ , we have obtained

$$c \cdot \gamma \text{cap}_p(\overline{\mathcal{K}_r(x_0)}; \mathcal{K}_{2r}(x_0)) < \text{cap}_p(\overline{\mathcal{K}_r(x_0)} \setminus \Omega; \mathcal{K}_{2r}(x_0)).$$

By the definition of capacity inradius relative to  $\mathcal{K}$ , this shows that we have  $r > R_{p,c,\gamma}(\Omega; \mathcal{K})$ . By the arbitrariness of  $r$ , we can finally obtain (11).

*Proof of (12).* Let us choose

$$d = d(N, p, \mathcal{K}) = \max \left\{ 1, R_{\mathcal{K}}^{N-p} \frac{\text{cap}_p(\overline{B_1}; B_2)}{\text{cap}_p(\overline{\mathcal{K}}; \mathcal{K}_2)} \left( \frac{2}{\text{dist}(\overline{\mathcal{K}}, \partial \mathcal{K}_2)} \frac{R_{\mathcal{K}}}{\lambda_p(B_1)^{\frac{1}{p}}} + 1 \right)^p \right\},$$

and take  $0 \leq \gamma < 1/d$ . We can suppose that  $R_{p,d,\gamma}(\Omega; \mathcal{K}) < +\infty$ . We take any  $r > R_{p,d,\gamma}(\Omega; \mathcal{K})$ , so that

$$\text{cap}_p(\overline{\mathcal{K}_r(x_0)} \setminus \Omega; \mathcal{K}_{2r}(x_0)) > d \cdot \gamma \text{cap}_p(\overline{\mathcal{K}_r(x_0)}; \mathcal{K}_{2r}(x_0)), \quad \text{for every } x_0 \in \mathbb{R}^N. \quad (15)$$

For the leftmost term, by using Lemma 3.5 with  $\Sigma = \overline{\mathcal{K}_r(x_0)} \setminus \Omega$  and  $\varrho = 2r$ , we have<sup>3</sup>

$$\begin{aligned} \text{cap}_p(\overline{\mathcal{K}_r(x_0)} \setminus \Omega; \mathcal{K}_{2r}(x_0)) &\leq \left( \frac{2}{\text{dist}(\overline{\mathcal{K}}, \partial\mathcal{K}_2)} \frac{R_{\mathcal{K}}}{\lambda_p(B_1)^{\frac{1}{p}}} + 1 \right)^p \\ &\quad \times \text{cap}_p(\overline{\mathcal{K}_r(x_0)} \setminus \Omega; B_{2R_{\mathcal{K}}r}(x_0)) \\ &\leq \left( \frac{2}{\text{dist}(\overline{\mathcal{K}}, \partial\mathcal{K}_2)} \frac{R_{\mathcal{K}}}{\lambda_p(B_1)^{\frac{1}{p}}} + 1 \right)^p \\ &\quad \times \text{cap}_p(\overline{B_{R_{\mathcal{K}}r}(x_0)} \setminus \Omega; B_{2R_{\mathcal{K}}r}(x_0)). \end{aligned} \quad (16)$$

The second inequality follows again from (6) since we have  $\mathcal{K}_r(x_0) \subseteq B_{R_{\mathcal{K}}r}(x_0)$ . Observe that by scaling

$$\begin{aligned} \text{cap}_p(\overline{\mathcal{K}_r(x_0)}; \mathcal{K}_{2r}(x_0)) &= r^{N-p} \frac{\text{cap}_p(\overline{\mathcal{K}}; \mathcal{K}_2)}{\text{cap}_p(\overline{B_1}, B_2)} \text{cap}_p(\overline{B_1}, B_2) \\ &= \left( \frac{1}{R_{\mathcal{K}}} \right)^{N-p} \frac{\text{cap}_p(\overline{\mathcal{K}}; \mathcal{K}_2)}{\text{cap}_p(\overline{B_1}, B_2)} \text{cap}_p(\overline{B_{R_{\mathcal{K}}r}(x_0)}, B_{2R_{\mathcal{K}}r}(x_0)), \end{aligned}$$

then, by recalling the definition of  $d$ , in particular we get

$$\text{cap}_p(\overline{\mathcal{K}_r(x_0)}; \mathcal{K}_{2r}(x_0)) \geq \frac{1}{d} \left( \frac{2R_{\mathcal{K}}}{\lambda_p(B_1)^{\frac{1}{p}}} + 1 \right)^p \text{cap}_p(\overline{B_{R_{\mathcal{K}}r}(x_0)}, B_{2R_{\mathcal{K}}r}(x_0)).$$

Together with (15) and (16), this implies that for every  $x_0 \in \mathbb{R}^N$ ,

$$\text{cap}_p(\overline{B_{R_{\mathcal{K}}r}(x_0)} \setminus \Omega; B_{2R_{\mathcal{K}}r}(x_0)) > d \cdot \gamma \text{cap}_p(\overline{B_{R_{\mathcal{K}}r}(x_0)}; B_{2R_{\mathcal{K}}r}(x_0)).$$

Therefore, by the definition of capacity inradius, we have

$$R_{\mathcal{K}} r \geq R_{p,\gamma}(\Omega).$$

By the arbitrariness of  $r > R_{p,d,\gamma}(\Omega; \mathcal{K})$ , we obtain (12).  $\square$

**Problem.** *Prove that there exist two constants  $a = a(N, p, \mathcal{K}, \gamma)$ ,  $b = b(N, p, \mathcal{K}, \gamma) > 0$  such that*

$$a R_{p,\gamma}(\Omega; \mathcal{K}) \leq R_{p,\gamma}(\Omega) \leq b R_{p,\gamma}(\Omega; \mathcal{K}), \quad \text{for every } 0 < \gamma < 1.$$

*This would permit to obtain a two-sided estimate on  $\lambda_p(\Omega)$ , and, more generally, on  $\lambda_{p,q}(\Omega)$  with  $q > p$  subcritical in terms on  $R_{p,\gamma}(\Omega; \mathcal{K})$ , no matter the shape  $\mathcal{K}$  and the negligibility parameter  $\gamma$ . For the particular case  $\mathcal{K} = (-1, 1)^N$ , such a result would extend [16, Theorem 15.4.1] (in the case of first-order Sobolev spaces) by dropping the restriction  $\gamma \leq \gamma_0(N, p)$  taken there (see [16, equation (14.1.2)]).*

**4. Different capacities.** In this section, rather than changing the basic “shape” used to compute the capacity, we will vary the concept of capacity itself, used to define the capacity inradius. In particular, we want to compare our definition of  $R_{p,\gamma}$  with the ones contained in [11] and [18].

<sup>3</sup>We use that

$$\text{dist}(\overline{\mathcal{K}_r(x_0)} \setminus \Omega; \partial\mathcal{K}_{2r}(x_0)) \geq \text{dist}(\overline{\mathcal{K}_r(x_0)}; \partial\mathcal{K}_{2r}(x_0)) = r \text{dist}(\overline{\mathcal{K}}; \partial\mathcal{K}_2).$$

**4.1. Maz’ya-Shubin inradius.** We start with the capacitary inradius defined in [18]. To this aim, we need at first to recall the following notion of  $p$ -capacity.

**Definition 4.1.** Let  $1 \leq p < N$  or  $p = N = 1$ . For every  $\Sigma \subseteq \mathbb{R}^N$  compact set, we define its *absolute homogeneous  $p$ -capacity* as

$$\text{cap}_p(\Sigma) = \inf_{\varphi \in C_0^\infty(\mathbb{R}^N)} \left\{ \int_{\mathbb{R}^N} |\nabla \varphi|^p dx : \varphi \geq 1 \text{ on } \Sigma \right\}.$$

Here as well, it is intended that  $\text{cap}_p(\emptyset) = 0$ . Observe that this coincides with  $\text{cap}_p(\Sigma; \mathbb{R}^N)$  in our previous notation, i.e. this is the  $p$ -capacity of  $\Sigma$  relative to the whole space  $\mathbb{R}^N$ .

**Remark 4.2.** The restriction  $p < N$  is unavoidable in the previous definition, at least for  $N \geq 2$ . Indeed, in this situation we have

$$\text{cap}_p(\Sigma) = 0, \quad \text{for every } p \geq N \text{ and every } \Sigma \subseteq \mathbb{R}^N \text{ compact set.}$$

For  $p > N$ , this follows by a simple scaling argument. The borderline case  $p = N \geq 2$  is slightly more delicate, because of the scale invariance of the  $N$ -Dirichlet integral (see for example [16, pages 148–149]).

We quantitatively compare the absolute capacity with the relative one. To this aim, we need to recall the definition of the sharp constant in Sobolev’s inequality for  $p < N$ , i.e.

$$\mathcal{S}_{N,p} = \sup_{\varphi \in C_0^\infty(\mathbb{R}^N)} \left\{ \left( \int_{\mathbb{R}^N} |\varphi|^{p^*} dx \right)^{\frac{p}{p^*}} : \|\nabla \varphi\|_{L^p(\mathbb{R}^N)} = 1 \right\}, \quad p^* = \frac{Np}{N-p}.$$

**Lemma 4.3** (Absolute vs. relative). *Let  $1 \leq p < N$ , and let  $\Sigma \subseteq \mathbb{R}^N$  be a non-empty compact set. If  $\Sigma \Subset E$  with  $E \subseteq \mathbb{R}^N$  an open bounded set, we have*

$$\text{cap}_p(\Sigma) \leq \text{cap}_p(\Sigma; E) \leq 2^{p-1} \left( 1 + \frac{|E|^{\frac{p}{N}}}{(\text{dist}(\Sigma, \partial E))^p} \mathcal{S}_{N,p} \right) \text{cap}_p(\Sigma).$$

*Proof.* The leftmost inequality trivially follows from the fact that, in the definition of  $\text{cap}_p(\Sigma)$ , we perform the infimum on a larger class of functions. For the rightmost one, let us take  $\varphi \in C_0^\infty(\mathbb{R}^N)$  such that  $\varphi \geq 1$  on  $\Sigma$ . For every  $0 < \delta < \text{dist}(\Sigma, \partial E)$ , we take the following Lipschitz cut-off function with compact support in  $E$ , i.e.

$$\eta_\delta(x) = \min \left\{ \frac{(\text{dist}(x, \partial E) - \delta)_+}{\text{dist}(\Sigma, \partial E) - \delta}, 1 \right\}, \quad \text{for } x \in E. \tag{17}$$

Observe that

$$0 \leq \eta_\delta \leq 1, \quad \eta_\delta \equiv 1 \text{ on } \Sigma, \quad |\nabla \eta_\delta| \leq \frac{1}{\text{dist}(\Sigma, \partial E) - \delta}.$$

The function  $\varphi \eta_\delta$  is feasible for the minimization problem which defines  $\text{cap}_p(\Sigma; E)$ . Thus, we obtain

$$\begin{aligned} \text{cap}_p(\Sigma; E) &\leq 2^{p-1} \int_E |\nabla \varphi|^p \eta_\delta^p dx + 2^{p-1} \int_E |\nabla \eta_\delta|^p |\varphi|^p dx \\ &\leq 2^{p-1} \int_{\mathbb{R}^N} |\nabla \varphi|^p dx + \frac{2^{p-1}}{(\text{dist}(\Sigma, \partial E) - \delta)^p} \int_E |\varphi|^p dx. \end{aligned}$$

To control the last term, we combine Hölder’s and Sobolev’s inequalities as follows:

$$\int_E |\varphi|^p dx \leq |E|^{1-\frac{p}{p^*}} \left( \int_E |\varphi|^{p^*} dx \right)^{\frac{p}{p^*}} \leq |E|^{1-\frac{p}{p^*}} \mathcal{S}_{N,p} \int_{\mathbb{R}^N} |\nabla \varphi|^p dx.$$

In conclusion, we obtain

$$\text{cap}_p(\Sigma; E) \leq 2^{p-1} \left( 1 + \frac{|E|^{\frac{p}{N}}}{(\text{dist}(\Sigma, \partial E) - \delta)^p} \mathcal{S}_{N,p} \right) \int_{\mathbb{R}^N} |\nabla \varphi|^p dx.$$

By the arbitrariness of  $\delta$  and  $\varphi$ , we get the conclusion.  $\square$

**Remark 4.4.** As one can easily see by inspecting the previous proof, the boundedness assumption on  $E$  is not really needed. The same proof works under the assumption  $|E| < +\infty$ .

For  $N \geq 2$ , we see that the rightmost inequality of Lemma 4.3 gets spoiled as  $p$  goes to  $N$ . Indeed, we recall that

$$\mathcal{S}_{N,p} \sim \frac{1}{(N-p)^{p-1}}, \quad \text{as } p \nearrow N,$$

see for example [22]. This is in accordance with Remark 4.2, thus the equivalence of the two capacities *must* cease to be true in this limit case.

The case  $p = N = 1$  is however exceptional: As simple and useless as it may seem, it deserves a separate statement.

**Lemma 4.5.** *Let  $\Sigma \subseteq \mathbb{R}$  be a non-empty compact set. If  $\Sigma \Subset E$  with  $E \subseteq \mathbb{R}$  an open bounded set, we have*

$$\text{cap}_1(\Sigma) \leq \text{cap}_1(\Sigma; E) \leq \left( 1 + \frac{|E|}{\text{dist}(\Sigma, \partial E)} \frac{1}{2} \right) \text{cap}_1(\Sigma).$$

*Proof.* We can proceed as in the proof of Lemma 4.3. The only difference is in the estimate of

$$\int_E |\varphi| dx \leq |E| \|\varphi\|_{L^\infty(\mathbb{R})} \leq \frac{|E|}{2} \int_{\mathbb{R}} |\varphi'| dx.$$

This is enough to conclude.  $\square$

**Definition 4.6.** Let  $1 \leq p < N$  or  $p = N = 1$ , and let  $0 \leq \gamma < 1$ . For an open set  $\Omega \subseteq \mathbb{R}^N$ , we define its *Maz'ya-Shubin capacity inradius* by

$$R_{p,\gamma}^{\text{MS}}(\Omega) := \sup \left\{ r > 0 : \exists x_0 \in \mathbb{R}^N \text{ s. t. } \text{cap}_p(\overline{B_r(x_0)} \setminus \Omega) \leq \gamma \text{cap}_p(\overline{B_r(x_0)}) \right\}.$$

Observe that the only difference with our capacity inradius  $R_{p,\gamma}(\Omega)$  is the use of the absolute homogeneous  $p$ -capacity in formulating the  $(p, \gamma)$ -negligibility condition, in place of the relative one.

**Proposition 4.7.** *Let  $1 \leq p < N$  or  $p = N = 1$ . Then, there exist two constants  $\alpha = \alpha(N, p) \leq 1$  and  $\beta = \beta(N, p) \geq 1$  such that, for every open set  $\Omega \subseteq \mathbb{R}^N$ , we have*

$$R_{p,\alpha,\gamma}^{\text{MS}}(\Omega) \leq R_{p,\gamma}(\Omega), \quad \text{for every } 0 \leq \gamma < 1,$$

and

$$R_{p,\gamma}(\Omega) \leq R_{p,\beta,\gamma}^{\text{MS}}(\Omega), \quad \text{for every } 0 \leq \gamma < \frac{1}{\beta}.$$

Moreover, for  $N \geq 2$  we have

$$\lim_{p \nearrow N} \alpha = 0 \quad \text{and} \quad \lim_{p \nearrow N} \beta = +\infty.$$

*Proof.* We can confine ourselves to consider the case  $N \geq 2$ ; the case  $N = 1$  can be handled in exactly the same manner by using Lemma 4.5 in place of Lemma 4.3.

In order to prove the first inequality, we take  $0 \leq \gamma < 1$ , and we can suppose that  $R_{p,\gamma}(\Omega) < +\infty$ . Let  $r > R_{p,\gamma}(\Omega)$ . This implies that

$$\text{cap}_p(\overline{B_r(x_0)} \setminus \Omega; B_{2r}(x_0)) > \gamma \text{cap}_p(\overline{B_r(x_0)}; B_{2r}(x_0)), \quad \text{for every } x_0 \in \mathbb{R}^N.$$

By using Lemma 4.3, this in particular entails that

$$\gamma \text{cap}_p(\overline{B_r(x_0)}) < 2^{p-1} \left(1 + (2\omega_N)^{\frac{p}{N}} \mathcal{S}_{N,p}\right) \text{cap}_p(\overline{B_r(x_0)} \setminus \Omega), \quad \text{for every } x_0 \in \mathbb{R}^N.$$

Thus, by the arbitrariness of  $r$  and Definition 4.6, we can infer that

$$R_{p,\alpha\cdot\gamma}^{\text{MS}}(\Omega) \leq R_{p,\gamma}(\Omega),$$

where  $\alpha$  is given by

$$\alpha := 2^{1-p} \left(1 + (2\omega_N)^{\frac{p}{N}} \mathcal{S}_{N,p}\right)^{-1}.$$

Observe that  $\alpha \leq 1$ , and for  $N \geq 2$  it converges to 0, as  $p$  goes to  $N$ .

In order to prove the second inequality, we set

$$\beta := \frac{\text{cap}_p(\overline{B_1}; B_2)}{\text{cap}_p(\overline{B_1})} = \left(1 - \left(\frac{1}{2}\right)^{\frac{N-p}{p-1}}\right)^{1-p}, \quad \text{if } 1 < p < N,$$

and  $\beta = 1$  in the case  $p = 1$ . We take  $0 \leq \gamma < 1/\beta$  and suppose that  $R_{p,\beta\cdot\gamma}^{\text{MS}}(\Omega) < +\infty$ , otherwise there is nothing to prove. We take any  $r > R_{p,\beta\cdot\gamma}^{\text{MS}}(\Omega)$ . Then, thanks to Lemma 4.3 and the definition of  $\beta$ , we have

$$\begin{aligned} \text{cap}_p(\overline{B_r(x_0)} \setminus \Omega; B_{2r}(x_0)) &\geq \text{cap}_p(\overline{B_r(x_0)} \setminus \Omega) \\ &> \beta \gamma \text{cap}_p(\overline{B_r(x_0)}) \\ &= \gamma \text{cap}_p(\overline{B_r(x_0)}; B_{2r}(x_0)), \end{aligned}$$

for every  $x_0 \in \mathbb{R}^N$ , where the second inequality comes from the Definition 4.6. In particular, we have that  $r > R_{p,\gamma}(\Omega)$ . By the arbitrariness of  $r$ , we eventually conclude that  $R_{p,\beta\cdot\gamma}^{\text{MS}}(\Omega) \geq R_{p,\gamma}(\Omega)$ .  $\square$

**Problem.** For  $1 \leq p < N$ , prove that there exist two constants  $c_1 = c_1(N, p, \gamma)$ ,  $c_2 = c_2(N, p, \gamma) > 0$  such that

$$c_1 R_{p,\gamma}^{\text{MS}}(\Omega) \leq R_{p,\gamma}(\Omega) \leq c_2 R_{p,\gamma}^{\text{MS}}(\Omega), \quad \text{for every } 0 < \gamma < 1.$$

For  $p = 2$ , this can be simply obtained by combining (3) with the two-sided estimate by Maz'ya and Shubin, i.e. [18, Theorem 1.1]. For a general  $p$ , one could proceed similarly. One then needs a generalization of their result, i.e. a two-sided estimate on  $\lambda_p(\Omega)$  in terms of  $R_{p,\gamma}^{\text{MS}}(\Omega)$ .

**4.2. Gallagher inradius.** First, we need the following alternative notion of absolute capacity.

**Definition 4.8.** Let  $1 \leq p < \infty$ . For every  $\Sigma \subseteq \mathbb{R}^N$  compact set, we define its *absolute inhomogeneous  $p$ -capacity* as

$$\mathcal{C}_p(\Sigma) = \inf_{\varphi \in C_0^\infty(\mathbb{R}^N)} \left\{ \int_{\mathbb{R}^N} |\nabla \varphi|^p dx + \int_{\mathbb{R}^N} |\varphi|^p dx : \varphi \geq 1 \text{ on } \Sigma \right\}.$$

As usual, we have that  $\mathcal{C}_p(\emptyset) = 0$ .

Observe that the two norms appearing in the minimization problem scale differently. Accordingly, this quantity does not enjoy a scaling relation<sup>4</sup>. Nevertheless, we have the following simple relation for the  $p$ -capacity of rescaled sets.

**Lemma 4.9.** *Let  $1 \leq p < \infty$ . For every  $\Sigma \subseteq \mathbb{R}^N$  compact set and every  $t > 0$ , we have*

$$\min\{t^{N-p}, t^N\} \mathcal{C}_p(\Sigma) \leq \mathcal{C}_p(t\Sigma) \leq \max\{t^{N-p}, t^N\} \mathcal{C}_p(\Sigma).$$

*Proof.* We can suppose that  $\Sigma \neq \emptyset$ , otherwise there is nothing to prove. Let  $\varphi \in C_0^\infty(\mathbb{R}^N)$  be such that  $\varphi \geq 1$  on  $\Sigma$ . Then, the rescaled function  $\varphi_t(x) = \varphi(x/t)$  belongs to  $C_0^\infty(\mathbb{R}^N)$  and is such that  $\varphi_t \geq 1$  on  $t\Sigma$ . Thus, with a change of variable, we get

$$\begin{aligned} \mathcal{C}_p(t\Sigma) &\leq t^{N-p} \int_{\mathbb{R}^N} |\nabla\varphi|^p dx + t^N \int_{\mathbb{R}^N} |\varphi|^p dx \\ &\leq \max\{t^{N-p}, t^N\} \left[ \int_{\mathbb{R}^N} |\nabla\varphi|^p dx + \int_{\mathbb{R}^N} |\varphi|^p dx \right]. \end{aligned}$$

By taking the infimum over  $\varphi$ , we get the upper bound.

For the lower bound, we proceed in a similar way. Let  $\varphi \in C_0^\infty(\mathbb{R}^N)$  be such that  $\varphi \geq 1$  on  $t\Sigma$ . Then, the rescaled function  $\varphi_t(x) = \varphi(tx)$  belongs to  $C_0^\infty(\mathbb{R}^N)$  and is such that  $\varphi_t \geq 1$  on  $\Sigma$ . Similarly as before, we get

$$\begin{aligned} \mathcal{C}_p(\Sigma) &\leq t^{p-N} \int_{\mathbb{R}^N} |\nabla\varphi|^p dx + t^{-N} \int_{\mathbb{R}^N} |\varphi|^p dx \\ &\leq \max\{t^{p-N}, t^{-N}\} \left[ \int_{\mathbb{R}^N} |\nabla\varphi|^p dx + \int_{\mathbb{R}^N} |\varphi|^p dx \right]. \end{aligned}$$

By the arbitrariness of  $\varphi$ , we conclude.  $\square$

The comparison between this  $p$ -capacity and the relative one is contained, for example, in [16, Proposition 13.1.1/2]. We repeat its simple proof so as to make precise the constants appearing in the estimate.

**Lemma 4.10.** *Let  $1 \leq p < \infty$  and let  $\Sigma \subseteq \mathbb{R}^N$  be a non-empty compact set. If  $\Sigma \Subset E$  with  $E \subseteq \mathbb{R}^N$  an open bounded set, we have*

$$\frac{\lambda_p(E)}{1 + \lambda_p(E)} \mathcal{C}_p(\Sigma) \leq \text{cap}_p(\Sigma; E) \leq 2^{p-1} \max \left\{ 1, \frac{1}{(\text{dist}(\Sigma, \partial E))^p} \right\} \mathcal{C}_p(\Sigma).$$

*Proof.* We take  $\varphi \in C_0^\infty(E)$  such that  $\varphi \geq 1$  on  $\Sigma$ . Thus, by the definition of  $\mathcal{C}_p(\Sigma)$  and the Poincaré inequality, we get

$$\mathcal{C}_p(\Sigma) \leq \int_{\mathbb{R}^N} |\nabla\varphi|^p dx + \int_{\mathbb{R}^N} |\varphi|^p dx \leq \left( 1 + \frac{1}{\lambda_p(E)} \right) \int_{\mathbb{R}^N} |\nabla\varphi|^p dx.$$

By the arbitrariness of  $\varphi$ , we get the first inequality.

For the second inequality, we take  $\varphi \in C_0^\infty(\mathbb{R}^N)$  such that  $\varphi \geq 1$  on  $\Sigma$  and the cut-off function  $\eta_\delta$  is defined as in (17). By using  $\varphi\eta_\delta$ , we get

$$\text{cap}_p(\Sigma; E) \leq 2^{p-1} \int_{\mathbb{R}^N} |\nabla\varphi|^p \eta_\delta^p dx + 2^{p-1} \int_{\mathbb{R}^N} |\nabla\eta_\delta|^p |\varphi|^p dx.$$

By using the properties of  $\eta_\delta$  and recalling the definition of  $\mathcal{C}_p(\Sigma)$ , we easily get the conclusion.  $\square$

<sup>4</sup>This explains the name *inhomogeneous*.

**Remark 4.11.** An inspection of the previous proof reveals again that the boundedness assumption on  $E$  is not really needed. It would be sufficient to suppose for example that  $E$  supports the  $p$ -Poincaré inequality. This condition is somehow necessary, as well: Take for example  $E = \mathbb{R}^N$  and  $\Sigma = \overline{B_r}$ . Then, we have

$$\text{cap}_p(\overline{B_r}; \mathbb{R}^N) = \text{cap}_p(\overline{B_r}).$$

For  $p \geq N \geq 2$ , we have already said that this is zero for every  $r > 0$ . For  $1 \leq p < N$ , we have

$$\text{cap}_p(\overline{B_r}) = N \omega_N \left( \frac{N-p}{p-1} \right)^{p-1} r^{N-p},$$

it is sufficient to take the limit as  $R$  goes to  $\infty$  in (7) and (8). On the other hand, by proceeding as in the proof of Lemma 2.3, we easily see that

$$\mathcal{C}_p(\overline{B_r}) \geq |\overline{B_r}| = \omega_N r^N.$$

Thus, in any case we get that

$$\lim_{r \rightarrow +\infty} \frac{\text{cap}_p(\overline{B_r})}{\mathcal{C}_p(\overline{B_r})} = 0.$$

Observe that the very same example works also in the borderline case  $p = N = 1$ , in light of Remark 2.2.

Based on the previous  $p$ -capacity, in [10, 11] the following capacity inradius is introduced. The definition is inspired by a related quantity defined by Souplet in [21, Section 2] using the Lebesgue measure in place of  $p$ -capacity.

**Definition 4.12.** Let  $1 \leq p < \infty$ . For an open set  $\Omega \subseteq \mathbb{R}^N$ , we define its *Gallagher capacity inradius* by

$$R_p^G(\Omega) := \sup \left\{ r > 0 : \forall \varepsilon > 0, \exists x_0 \in \mathbb{R}^N \text{ such that } \mathcal{C}_p(\overline{B_r(x_0)} \setminus \Omega) < \varepsilon \right\}.$$

Observe that we have also included the case  $p = 1$ , which was not considered in [11].

Since this definition is based on an inhomogeneous concept of  $p$ -capacity, it is not clear whether  $R_p^G$  enjoys some scaling rule or not. This is actually the case, as shown in the following lemma.

**Lemma 4.13.** *Let  $1 \leq p < \infty$  and let  $\Omega \subseteq \mathbb{R}^N$  be an open set. Then, for every  $t > 0$  we have*

$$R_p^G(t\Omega) = t R_p^G(\Omega).$$

*Proof.* By the definition of supremum, for every  $\delta > 0$  there exists  $R_p^G(t\Omega) - \delta < r_\delta < R_p^G(t\Omega)$ . Thus, for every  $\varepsilon > 0$  there exists a point  $x_0 \in \mathbb{R}^N$  such that

$$\mathcal{C}_p(\overline{B_{r_\delta}(x_0)} \setminus (t\Omega)) < \varepsilon.$$

Observe that

$$\overline{B_{r_\delta}(x_0)} \setminus (t\Omega) = \overline{t B_{r_\delta/t}(x_0/t)} \setminus (t\Omega) = t \left( \overline{B_{r_\delta/t}(x_0/t)} \setminus \Omega \right).$$

Thus, in particular we have

$$\mathcal{C}_p \left( \overline{B_{r_\delta/t}(x_0/t)} \setminus \Omega \right) < \frac{1}{\min\{t^N, t^{N-p}\}} \varepsilon,$$

thanks to Lemma 4.9. By the arbitrariness of  $\varepsilon > 0$ , this entails that

$$\frac{R_p^G(t\Omega) - \delta}{t} < \frac{r_\delta}{t} \leq R_p^G(\Omega).$$

Finally, by taking the limit as  $\delta$  goes to 0, we obtain

$$\frac{R_p^G(t\Omega)}{t} \leq R_p^G(\Omega).$$

To prove the reverse inequality, we can proceed similarly. For every  $\delta > 0$ , there exists  $R_p^G(\Omega) - \delta < r_\delta < R_p^G(\Omega)$ . Thus, for every  $\varepsilon > 0$  there exists a point  $x_0 \in \mathbb{R}^N$  such that

$$\mathcal{C}_p(\overline{B_{r_\delta}(x_0)} \setminus \Omega) < \varepsilon.$$

Observe that

$$\overline{B_{r_\delta}(x_0)} \setminus \Omega = \frac{1}{t} t \left( \overline{B_{r_\delta}(x_0)} \setminus \Omega \right) = \frac{1}{t} \left( \overline{B_{tr_\delta}(tx_0)} \setminus (t\Omega) \right).$$

Thus, this time we have

$$\min \left\{ \left( \frac{1}{t} \right)^{N-p}, \left( \frac{1}{t} \right)^N \right\} \mathcal{C}_p \left( \overline{B_{tr_\delta}(tx_0)} \setminus (t\Omega) \right) < \varepsilon,$$

again by Lemma 4.9. By the arbitrariness of  $\varepsilon > 0$ , this entails that

$$t(R_p^G(\Omega) - \delta) < tr_\delta \leq R_p^G(t\Omega).$$

By taking the limit as  $\delta$  goes to 0, we conclude. □

The capacitary inradius  $R_p^G$  admits the following equivalent definition in terms of the relative  $p$ -capacity. The following is a particular case of [11, Lemma 2.12].

**Lemma 4.14.** *Let  $1 \leq p < \infty$ . For every open set  $\Omega \subseteq \mathbb{R}^N$ , we have*

$$R_p^G(\Omega) = \sup \left\{ r > 0 : \forall \varepsilon > 0, \exists x_0 \in \mathbb{R}^N \text{ s. t. } \text{cap}_p \left( \overline{B_r(x_0)} \setminus \Omega; B_{2r}(x_0) \right) < \varepsilon \right\}.$$

*Proof.* We suppose that  $R_p^G(\Omega) < +\infty$ , and in the case  $R_p^G(\Omega) = +\infty$ , the argument below can be easily adapted. For every  $\delta > 0$ , by definition there exists  $R_p^G(\Omega) - \delta < r_\delta < R_p^G(\Omega)$  such that, for every  $\varepsilon > 0$ , we have

$$\mathcal{C}_p(\overline{B_{r_\delta}(x_0)} \setminus \Omega) < \varepsilon,$$

for some  $x_0 \in \mathbb{R}^N$ . In particular, from Lemma 4.10 with  $\Sigma = \overline{B_{r_\delta}(x_0)} \setminus \Omega$  and  $E = B_{2r_\delta}(x_0)$ , we get

$$\text{cap}_p \left( \overline{B_{r_\delta}(x_0)} \setminus \Omega; B_{2r_\delta}(x_0) \right) < \varepsilon 2^{p-1} \max \left\{ 1, \frac{1}{r_\delta^p} \right\}.$$

By the arbitrariness of  $\varepsilon > 0$ , this implies that

$$\sup \left\{ r > 0 : \forall \varepsilon > 0, \exists x_0 \in \mathbb{R}^N \text{ such that } \text{cap}_p \left( \overline{B_r(x_0)} \setminus \Omega; B_{2r}(x_0) \right) < \varepsilon \right\} \geq r_\delta.$$

By recalling that  $R_p^G(\Omega) - \delta < r_\delta < R_p^G(\Omega)$ , the arbitrariness of  $\delta$  implies that

$$\sup \left\{ r > 0 : \forall \varepsilon > 0, \exists x_0 \in \mathbb{R}^N \text{ s. t. } \text{cap}_p \left( \overline{B_r(x_0)} \setminus \Omega; B_{2r}(x_0) \right) < \varepsilon \right\} \geq R_p^G(\Omega).$$

The reverse inequality can be proved in a similar way by again using Lemma 4.10. This concludes the proof. □

About the comparison between  $R_{p,\gamma}$  and  $R_p^G$ , we have the following result, which is essentially contained in [11] up to some technical adjustments in the proof.

**Proposition 4.15.** *Let  $1 \leq p \leq N$ . Then, for every open set  $\Omega \subseteq \mathbb{R}^N$ , we have*

$$R_p^G(\Omega) = \inf_{0 < \gamma < 1} R_{p,\gamma}(\Omega) = \lim_{\gamma \searrow 0} R_{p,\gamma}(\Omega).$$

*Proof.* The fact that

$$\inf_{0 < \gamma < 1} R_{p,\gamma}(\Omega) = \lim_{\gamma \searrow 0} R_{p,\gamma}(\Omega),$$

simply follows from the monotonicity of  $\gamma \mapsto R_{p,\gamma}(\Omega)$ . Thus, it is sufficient to prove the first identity. We then divide the proof in two parts.

*Part 1.* Here, we show that

$$R_{p,\gamma}(\Omega) \geq R_p^G(\Omega), \quad \text{for every } \gamma \in (0, 1), \tag{18}$$

by proceeding as in the proof of [11, Lemma 2.16]. Let us fix  $\gamma \in (0, 1)$ . We can assume that  $R_p^G(\Omega) < +\infty$ , and in the case  $R_p^G(\Omega) = +\infty$  the reader will immediately see how to adapt the argument below. By the definition of supremum, for every  $\delta > 0$  there exists  $R_p^G(\Omega) - \delta < r_\delta < R_p^G(\Omega)$  with the following property: For every  $\varepsilon > 0$ , there exists a point  $x_0 \in \mathbb{R}^N$  such that

$$\text{cap}_p(\overline{B_{r_\delta}(x_0)} \setminus \Omega; B_{2r_\delta}(x_0)) < \varepsilon.$$

We also used the equivalent formulation of  $R_p^G(\Omega)$  contained in Lemma 4.14. Thus, if we use the property above with the choice

$$\varepsilon = \gamma \text{cap}_p(\overline{B_{r_\delta}}; B_{2r_\delta}),$$

we get existence of a center  $x_0$  such that

$$\text{cap}_p(\overline{B_{r_\delta}(x_0)} \setminus \Omega; B_{2r_\delta}(x_0)) < \gamma \text{cap}_p(\overline{B_{r_\delta}(x_0)}; B_{2r_\delta}(x_0)).$$

This shows that the radius  $r_\delta$  is admissible for the definition of  $R_{p,\gamma}(\Omega)$ . Thus, we obtain

$$R_{p,\gamma}(\Omega) \geq r_\delta > R_p^G(\Omega) - \delta.$$

By the arbitrariness of  $\delta > 0$ , this in turn implies (18).

*Part 2.* In order to prove that

$$\lim_{\gamma \searrow 0} R_{p,\gamma}(\Omega) \leq R_p^G(\Omega), \tag{19}$$

we need to distinguish two possibilities<sup>5</sup>:

- (i) either  $R_{p,\gamma}(\Omega) = +\infty$  for all  $0 < \gamma < 1$ ,
- (ii) or  $R_{p,\gamma}(\Omega) < +\infty$  for all  $0 < \gamma < 1$ .

If alternative (i) occurs, we claim that we must have  $R_p^G(\Omega) = +\infty$ , as well. This would establish (19). Indeed, let us fix a radius  $r > 0$ . Then, for every  $\varepsilon > 0$  we set

$$\gamma_\varepsilon = \min \left\{ \frac{1}{2}, \frac{\varepsilon}{2 \text{cap}_p(\overline{B_r}; B_{2r})} \right\} = \min \left\{ \frac{1}{2}, \frac{\varepsilon}{2 r^{N-p} \text{cap}_p(\overline{B_1}; B_2)} \right\}.$$

Since  $R_{p,\gamma_\varepsilon}(\Omega) = +\infty$ , there exists a point  $x_0 \in \mathbb{R}^N$  such that

$$\text{cap}_p(\overline{B_r(x_0)} \setminus \Omega; B_{2r}(x_0)) \leq \gamma_\varepsilon \text{cap}_p(\overline{B_r(x_0)}; B_{2r}(x_0)) < \varepsilon.$$

According to Lemma 4.14, this shows that  $r \leq R_p^G(\Omega)$ . By the arbitrariness of  $r > 0$ , we obtain  $R_p^G(\Omega) = +\infty$ , as claimed.

<sup>5</sup>We recall that the condition  $R_{p,\gamma}(\Omega) < +\infty$  does not depend on the particular  $\gamma \in (0, 1)$ , thanks to (3).

We now assume alternative (ii): Observe that we also have  $R_p^G(\Omega) < +\infty$  by *Part 1*. We need to show that for every  $0 < \delta \ll 1$ , there exists  $0 < \gamma_\delta < 1$  such that

$$R_{p,\gamma_\delta}(\Omega) \leq R_p^G(\Omega) + \delta. \quad (20)$$

This would be sufficient to get (19). We preliminarily observe that we can suppose that

$$R_p^G(\Omega) < R_{p,1/2}(\Omega).$$

Otherwise, in light of *Part 1*, we would have  $R_p^G(\Omega) = R_{p,1/2}(\Omega)$ , and thus (20) would follow with  $\gamma_\delta \equiv 1/2$ .

We then choose any  $0 < \delta < \delta_0 := R_{p,1/2}(\Omega) - R_p^G(\Omega)$  so that we still have

$$R_p^G(\Omega) + \delta < R_{p,1/2}(\Omega).$$

Observe that since  $R_p^G(\Omega) + \delta > R_p^G(\Omega)$ , there exists an  $\varepsilon_\delta > 0$  such that

$$\text{cap}_p \left( \overline{B_{R_p^G(\Omega)+\delta}(x_0)} \setminus \Omega; B_{2(R_p^G(\Omega)+\delta)}(x_0) \right) \geq \varepsilon_\delta, \quad \text{for every } x_0 \in \mathbb{R}^N,$$

thanks to the equivalent formulation of Lemma 4.14. Thus, from Lemma 2.4, for every  $r > R_p^G(\Omega) + \delta$ , we get

$$\text{cap}_p(\overline{B_r(x_0)} \setminus \Omega; B_{2r}(x_0)) \geq \varepsilon_\delta, \quad \text{for every } x_0 \in \mathbb{R}^N,$$

In particular, for every radius  $r$  such that

$$R_p^G(\Omega) + \delta < r \leq R_{p,1/2}(\Omega),$$

we also have

$$\varepsilon_\delta \frac{\text{cap}_p(\overline{B_r(x_0)}; B_{2r}(x_0))}{\left(R_{p,1/2}(\Omega)\right)^{N-p} \text{cap}_p(\overline{B_1}; B_2)} \leq \text{cap}_p(\overline{B_r(x_0)} \setminus \Omega; B_{2r}(x_0)),$$

thanks to the scaling properties of the relative  $p$ -capacity and the restriction  $r \leq R_{p,1/2}(\Omega)$ . In particular, if we define

$$\gamma_\delta = \min \left\{ \frac{1}{4}, \frac{\varepsilon_\delta}{2} \frac{1}{\left(R_{p,1/2}(\Omega)\right)^{N-p} \text{cap}_p(\overline{B_1}; B_2)} \right\},$$

from the previous estimate, we get

$$\gamma_\delta \text{cap}_p(\overline{B_r(x_0)}; B_{2r}(x_0)) < \text{cap}_p(\overline{B_r(x_0)} \setminus \Omega; B_{2r}(x_0)).$$

This holds for every  $R_p^G(\Omega) + \delta < r \leq R_{p,1/2}(\Omega)$  and every  $x_0 \in \mathbb{R}^N$ . Thus, we have two possibilities:

$$\text{either } R_{p,\gamma_\delta}(\Omega) \leq R_p^G(\Omega) + \delta \quad \text{or} \quad R_{p,\gamma_\delta}(\Omega) > R_{p,1/2}(\Omega).$$

Since  $\gamma_\delta \leq 1/4$  by construction and  $\gamma \mapsto R_{p,\gamma}(\Omega)$  is non-decreasing, the second alternative can not occur. Thus, we finally get (20).  $\square$

**Remark 4.16.** We observe that for a general open set, it may happen that

$$R_{p,0}(\Omega) < \lim_{\gamma \searrow 0} R_{p,\gamma}(\Omega).$$

By Proposition 4.15, it is sufficient to construct an open set such that  $R_{p,0}(\Omega) < R_p^G(\Omega)$ ; this can be done as in [11, Examples].

We conclude this section by noticing that while we have (see [11, Theorem 1.3])

$$\lambda_p(\Omega) \leq \lambda_p(B_1) \left( \frac{1}{R_p^G(\Omega)} \right)^p,$$

it is *not* possible to have a lower bound of the type

$$C_{N,p} \left( \frac{1}{R_p^G(\Omega)} \right)^p \leq \lambda_p(\Omega), \quad \text{for every open set } \Omega \subseteq \mathbb{R}^N. \quad (21)$$

This is the content of the next example.

**Example 4.17.** Let  $0 < \delta < 1/4$  be fixed. We introduce the periodically perforated set

$$\Omega_\delta = \mathbb{R}^N \setminus \bigcup_{\mathbf{i} \in \mathbb{Z}^N} \overline{B_\delta(\mathbf{i})},$$

as in [3, Example A.1]. Take  $r > \sqrt{N}/2 + 1/4$ . Let  $x_0 \in \mathbb{R}^N$  and let  $\mathbf{i}_0 \in \mathbb{Z}^N$  be such that

$$|x_0 - \mathbf{i}_0| = \text{dist}(x_0, \mathbb{Z}^N).$$

For every  $y \in B_\delta(\mathbf{i}_0)$ , by using the triangle inequality, we have

$$|y - x_0| \leq |y - \mathbf{i}_0| + |\mathbf{i}_0 - x_0| < \delta + \frac{\sqrt{N}}{2} < \frac{1}{4} + \frac{\sqrt{N}}{2} < r,$$

that is,

$$B_\delta(\mathbf{i}_0) \subseteq B_r(x_0).$$

This entails that

$$|\overline{B_r(x_0)} \setminus \Omega_\delta| \geq |B_\delta(\mathbf{i}_0)| = \omega_N \delta^N, \quad \text{for every } x_0 \in \mathbb{R}^N. \quad (22)$$

Then, by using Lemma 4.10 in combination with Lemma 2.3, we get

$$\begin{aligned} \mathcal{C}_p(\overline{B_r(x_0)} \setminus \Omega_\delta) &\geq 2^{1-p} \frac{r^p}{r^p + 1} \text{cap}_p(\overline{B_r(x_0)} \setminus \Omega_\delta; B_{2r}(x_0)) \\ &\geq 2^{1-p} \frac{r^p}{r^p + 1} \lambda_p(B_{2r}(x_0)) |\overline{B_r(x_0)} \setminus \Omega_\delta|. \end{aligned}$$

By using that  $\lambda_p(B_{2r}(x_0)) = (2r)^{-p} \lambda_p(B_1)$  and (22), we get

$$\mathcal{C}_p(\overline{B_r(x_0)} \setminus \Omega_\delta) \geq \varepsilon_0, \quad \text{where } \varepsilon_0 = \frac{2}{4^p} \frac{\omega_N}{r^p + 1} \lambda_p(B_1) \delta^N.$$

By the arbitrariness of  $x_0 \in \mathbb{R}^N$ , this shows that we must have  $r > R_p^G(\Omega_\delta)$ . In turn, by the arbitrariness of  $r$ , we also have

$$R_p^G(\Omega_\delta) \leq \frac{\sqrt{N}}{2} + \frac{1}{4}, \quad \text{for every } 0 < \delta < \frac{1}{4}.$$

On the other hand, by using the same computations of [3, Example A.1], we have that

$$\lim_{\delta \searrow 0} \lambda_p(\Omega_\delta) = 0.$$

Thus, an estimate like (21) cannot be true.

**5. Capacitary inradius and measure density condition.**

**5.1. A two-sided estimate.** In this section, we want to compare our capacity inradius with the classical inradius under a simple measure density condition on the open sets. Namely, we will consider open sets  $\Omega \subsetneq \mathbb{R}^N$  such that the *measure density index*

$$\theta_{\Omega, r_0}^*(t) := \inf \left\{ \left( \frac{r_0}{r} \right)^t \frac{|B_r(x) \setminus \Omega|}{|B_r(x)|} : x \in \partial\Omega, 0 < r \leq r_0 \right\},$$

is positive for some  $r_0 > 0$  and  $t \geq 0$ . We notice that, by definition, we have

$$\theta_{\Omega, r_0}^*(t) \leq \frac{|B_{r_0}(x) \setminus \Omega|}{|B_{r_0}(x)|} \leq 1.$$

The next simple result gives a quantitative equivalence of this index with a slightly different one, which will be more handy.

**Proposition 5.1.** *Let  $\Omega \subsetneq \mathbb{R}^N$  be an open set and let  $t \geq 0$ ,  $r_0 > 0$ . We set*

$$\theta_{\Omega, r_0}(t) = \inf \left\{ \left( \frac{r_0}{r} \right)^t \frac{|B_r(x) \setminus \Omega|}{|B_r(x)|} : x \in \mathbb{R}^N \setminus \Omega, 0 < r \leq r_0 \right\}. \quad (23)$$

Then, we have

$$\theta_{\Omega, r_0}(t) \leq \theta_{\Omega, r_0}^*(t) \leq 2^{N+t} \theta_{\Omega, r_0}(t).$$

*Proof.* This is essentially a particular case of [12, Theorem 3.1]. We reproduce the proof to keep track of the relevant constant.

The leftmost inequality follows from the fact that  $\partial\Omega \subseteq \mathbb{R}^N \setminus \Omega$ . In order to prove the rightmost inequality, we can assume that

$$\theta_{\Omega, r_0}^*(t) > 0, \quad (24)$$

otherwise there is nothing to prove. We fix a point  $x \in \mathbb{R}^N \setminus \Omega$  and a radius  $0 < r \leq r_0$ . We set

$$d_{\Omega}(x) = \min_{y \in \partial\Omega} |x - y|,$$

and distinguish two cases for  $0 < \varepsilon < 1$ .

**Case  $d_{\Omega}(x) \geq (1 - \varepsilon)r$ .** By observing that  $B_{(1-\varepsilon)r}(x) \subseteq \mathbb{R}^N \setminus \Omega$ , we have

$$\begin{aligned} |B_r(x) \setminus \Omega| &\geq |B_{(1-\varepsilon)r}(x) \setminus \Omega| \\ &= |B_{(1-\varepsilon)r}(x)| \\ &= \omega_N (1 - \varepsilon)^N r^N \\ &\geq \theta_{\Omega, r_0}^*(t) \left( \frac{r}{r_0} \right)^t \omega_N (1 - \varepsilon)^N r^N, \end{aligned} \quad (25)$$

where in the last inequality we used the trivial fact that  $\theta_{\Omega, r_0}^*(t) \leq 1$  and  $r \leq r_0$ .

**Case  $d_{\Omega}(x) < (1 - \varepsilon)r$ .** In this case, it must result that  $B_{(1-\varepsilon)r}(x) \cap \partial\Omega \neq \emptyset$ . Given  $y \in B_{(1-\varepsilon)r}(x) \cap \partial\Omega$ , we set  $\delta := r - |x - y|$ . In particular, by construction we have

$$\varepsilon r < \delta \leq r \quad \text{and} \quad B_{\delta}(y) \subseteq B_r(x).$$

By using (24), we then infer that

$$\begin{aligned} |B_r(x) \setminus \Omega| &\geq |B_\delta(y) \setminus \Omega| \\ &\geq \theta_{\Omega, r_0}^*(t) \left(\frac{\delta}{r_0}\right)^t \omega_N \delta^N \\ &> \theta_{\Omega, r_0}^*(t) \left(\frac{r}{r_0}\right)^t \omega_N \varepsilon^{N+t} r^N. \end{aligned} \quad (26)$$

If we now combine (25) and (26), we eventually obtain

$$\left(\frac{r_0}{r}\right)^t \frac{|B_r(x) \setminus \Omega|}{|B_r(x)|} \geq \min\{(1-\varepsilon)^N, \varepsilon^{N+t}\} \theta_{\Omega, r_0}^*(t).$$

By the arbitrariness of both  $x \in \mathbb{R}^N \setminus \Omega$  and  $0 < r \leq r_0$ , we get

$$\theta_{\Omega, r_0}(t) \geq \min\{(1-\varepsilon)^N, \varepsilon^{N+t}\} \theta_{\Omega, r_0}^*(t).$$

The choice  $\varepsilon = 1/2$  leads to the desired conclusion.  $\square$

We now come to the main result of this paper.

**Theorem 5.2.** *Let  $1 \leq p \leq N$  and let  $\Omega \subsetneq \mathbb{R}^N$  be an open set such that  $\theta_{\Omega, r_0}^*(t) > 0$ , for some  $r_0 > 0$  and  $t \geq 0$ . Then, there exists an explicit parameter  $0 < \gamma_0 = \gamma_0(N, p, \theta_{\Omega, r_0}^*(t), t, r_0/r_\Omega) < 1$  such that if we set*

$$\mathcal{C} = \begin{cases} 6\sqrt{N}, & \text{if } 0 \leq \gamma < \gamma_0, \\ 6\sqrt{N} \left(\frac{2C_{N,p,\gamma}}{\gamma_0 \sigma_{N,p}}\right)^{\frac{1}{p}}, & \text{if } \gamma_0 \leq \gamma < 1, \end{cases}$$

with  $C_{N,p,\gamma}$  and  $\sigma_{N,p}$  as in (3), then we have

$$r_\Omega \leq R_{p,\gamma}(\Omega) \leq \mathcal{C} r_\Omega, \quad \text{for every } 0 \leq \gamma < 1. \quad (27)$$

*Proof.* The leftmost inequality in (27) follows from the definition of capacitary inradius, so we focus on the other estimate. Without loss of generality, we can assume that  $r_\Omega < +\infty$ . We divide the proof into two parts: We first construct  $\gamma_0$  and show that the estimate is true in the range  $0 \leq \gamma < \gamma_0$ ; then, we show how this is sufficient to get the claimed estimate for  $\gamma \geq \gamma_0$ , as well.

*Part 1: Case  $\gamma < \gamma_0$ .* We set

$$\alpha := \frac{r_0}{r_\Omega} \quad \text{and} \quad A := 2\sqrt{N} (2 + \min\{1, \alpha\}), \quad (28)$$

in particular  $A \leq 6\sqrt{N}$ . We take an open ball  $B_r(x_0)$  having radius  $r > A r_\Omega$ . We set<sup>6</sup>

$$\ell = r_\Omega (2 + \min\{1, \alpha\}) \quad \text{and} \quad m = \left\lfloor \frac{r/\sqrt{N}}{\ell} \right\rfloor, \quad (29)$$

and then

$$Q_{m\ell}(x_0) \subseteq Q_{r/\sqrt{N}}(x_0) \subseteq B_r(x_0).$$

<sup>6</sup>For every  $\delta \in \mathbb{R}$ , we denote by  $[\delta]$  its integer part, defined by

$$[\delta] = \max\{n \in \mathbb{Z} : \delta \geq n\}.$$




FIGURE 1. A cube  $Q_\ell(\mathbf{p}_i)$ , containing a concentric ball with radius  $2r_\Omega$ . The black dot indicates a point outside  $\Omega$ . By construction, at this point, we can center a ball which is feasible for the measure density index  $\theta_{\Omega, r_0}(t)$ .

Moreover, the cube  $Q_{m\ell}(x_0)$  can be tiled (up to a Lebesgue negligible set) with  $m^N$  disjoint cubes having edge length  $2\ell$ ; let us refer to these cubes as  $\{Q_\ell(\mathbf{p}_i)\}_{i=1}^{m^N}$ . By construction, the cube  $Q_\ell(\mathbf{p}_i)$  contains the ball  $B_{2r_\Omega}(\mathbf{p}_i)$ . Thanks to the definition of inradius, there exists in particular a point  $x_i \in B_{2r_\Omega}(\mathbf{p}_i) \setminus \Omega$ . By recalling the definition of  $\ell$ , we have that

$$B_{\ell-2r_\Omega}(x_i) \subseteq Q_\ell(\mathbf{p}_i).$$

Observe that  $\ell - 2r_\Omega = r_\Omega \min\{1, \alpha\} \leq r_0$ . Thus, in particular, we have that each ball  $B_{\ell-2r_\Omega}(x_i)$  is feasible for the definition of the modified measure density index  $\theta_{\Omega, r_0}(t)$  defined by (23) (see Figure 1). Accordingly, we get

$$\begin{aligned} |B_{\ell-2r_\Omega}(x_i) \setminus \Omega| &\geq \theta_{\Omega, r_0}(t) \omega_N (\ell - 2r_\Omega)^N \left(\frac{\ell - 2r_\Omega}{r_0}\right)^t \\ &= \theta_{\Omega, r_0}(t) \omega_N \min\{1, \alpha^{N+t}\} r_\Omega^N \left(\frac{r_\Omega}{r_0}\right)^t \\ &= \theta_{\Omega, r_0}(t) \omega_N \min\{\alpha^{-t}, \alpha^N\} r_\Omega^N, \end{aligned}$$

By taking into account all the contributions of these balls, we can estimate

$$\begin{aligned} |B_r(x_0) \setminus \Omega| &\geq \sum_{i=1}^{m^N} |Q_\ell(\mathbf{p}_i) \setminus \Omega| \\ &\geq \sum_{i=1}^{m^N} |B_{\ell-2r_\Omega}(x_i) \setminus \Omega| \\ &\geq m^N \theta_{\Omega, r_0}(t) \omega_N \min\{\alpha^{-t}, \alpha^N\} r_\Omega^N. \end{aligned} \tag{30}$$

We claim that the latter can be further estimated from below by a term of the type  $cr^N$ . Indeed, by recalling that we are assuming  $r > Ar_\Omega$  and using both (28) and

(29), the number of cubes  $m$  is such that

$$m = \left\lceil \frac{r/\sqrt{N}}{\ell} \right\rceil \geq \frac{r/\sqrt{N}}{\ell} - 1 > \frac{r/\sqrt{N}}{\ell} - \frac{\sqrt{N}}{A} \frac{r/\sqrt{N}}{r_\Omega} = \frac{r}{r_\Omega} \frac{1}{2\sqrt{N}} \frac{1}{2 + \min\{1, \alpha\}}.$$

We use this information in (30) to obtain

$$|B_r(x_0) \setminus \Omega| \geq \left( \frac{1}{2\sqrt{N}} \frac{\min\{\alpha^{-t/N}, \alpha\}}{2 + \min\{1, \alpha\}} \right)^N \theta_{\Omega, r_0}(t) \omega_N r^N.$$

By further using Proposition 5.1, we can replace  $\theta_{\Omega, r_0}(t)$  with  $\theta_{\Omega, r_0}^*(t)$  up to a modification of the constant. Namely, we can infer

$$|\overline{B_r(x_0)} \setminus \Omega| \geq \left( \frac{1}{4 \cdot 2^{t/N} \sqrt{N}} \frac{\min\{\alpha^{-t/N}, \alpha\}}{2 + \min\{1, \alpha\}} \right)^N \theta_{\Omega, r_0}^*(t) \omega_N r^N. \quad (31)$$

On the other hand, by using Lemma 2.3 with  $\Sigma = \overline{B_r(x_0)} \setminus \Omega$  and  $E = B_{2r}(x_0)$ , the measure of  $\overline{B_r(x_0)} \setminus \Omega$  bounds from below its relative  $p$ -capacity. More precisely, we have

$$\text{cap}_p(\overline{B_r(x_0)} \setminus \Omega; B_{2r}(x_0)) \geq \frac{\lambda_p(B_2)}{r^p} |\overline{B_r(x_0)} \setminus \Omega|.$$

On account of the lower bound (31), this yields

$$\begin{aligned} & \text{cap}_p(\overline{B_r(x_0)} \setminus \Omega; B_{2r}(x_0)) \\ & \geq \lambda_p(B_2) |B_1| \left( \frac{1}{4 \cdot 2^{t/N} \sqrt{N}} \frac{\min\{\alpha^{-t/N}, \alpha\}}{2 + \min\{1, \alpha\}} \right)^N \theta_{\Omega, r_0}^*(t) r^{N-p}. \end{aligned}$$

On the other hand, if  $B_r(x_0)$  is  $(p, \gamma)$ -negligible, we would have

$$\gamma r^{N-p} \text{cap}_p(\overline{B_1}; B_2) \geq \text{cap}_p(\overline{B_r(x_0)} \setminus \Omega; B_{2r}(x_0)).$$

By joining the last two estimates and canceling out the term  $r^{N-p}$  on both sides, we eventually obtain that it must result<sup>7</sup> in

$$\gamma \geq \frac{\lambda_p(B_2) |B_1|}{\text{cap}_p(\overline{B_1}; B_2)} \left( \frac{1}{4 \cdot 2^{t/N} \sqrt{N}} \frac{\min\{\alpha^{-t/N}, \alpha\}}{2 + \min\{1, \alpha\}} \right)^N \theta_{\Omega, r_0}^*(t) =: \gamma_0. \quad (32)$$

This shows that every ball with radius  $r > A r_\Omega$  cannot be  $(p, \gamma)$ -negligible for  $\gamma < \gamma_0$ . On account of the definition (2) of capacity inradius, we obtain

$$R_{p, \gamma}(\Omega) \leq A r_\Omega, \quad \text{for every } 0 \leq \gamma < \gamma_0.$$

As already observed, we have in particular  $A \leq 6\sqrt{N}$ , and thus our claim follows.

*Part 2: Case  $\gamma \geq \gamma_0$ .* We can exploit the two-sided estimate (3). Indeed, by using it with  $\gamma \geq \gamma_0$  on the right-hand side and with  $\gamma = \gamma_0/2$  on the left-hand side, in particular, we get

$$\sigma_{N,p} \frac{\gamma_0}{2} \left( \frac{1}{R_{p, \gamma_0/2}(\Omega)} \right)^p \leq \lambda_p(\Omega) \leq C_{N,p,\gamma} \left( \frac{1}{R_{p,\gamma}(\Omega)} \right)^p.$$

<sup>7</sup>Observe that  $\gamma_0 < 1$ , since, by Lemma 2.3 with  $E = B_2$  and  $\Sigma = \overline{B_1}$ , we have

$$\frac{\lambda_p(B_2) |B_1|}{\text{cap}_p(\overline{B_1}; B_2)} \leq 1.$$

Thus, we also obtain

$$R_{p,\gamma}(\Omega) \leq \left( \frac{2C_{N,p,\gamma}}{\sigma_{N,p}\gamma_0} \right)^{\frac{1}{p}} R_{p,\gamma_0/2}(\Omega).$$

By using the result of the first part, we now conclude.  $\square$

**Remark 5.3** (The case  $p > N$ ). This case is simpler, and one can prove that  $R_{p,\gamma}(\Omega)$  and  $r_\Omega$  are equivalent for every open set. This is clearly due to the fact that for  $p > N$ , the only set with zero  $p$ -capacity is the empty set. We refer to [3, Section 7] for the discussion of this case.

**Remark 5.4.** From the explicit expression (32) of the parameter  $\gamma_0$ , we see that we have

$$\gamma_0 \rightarrow 0 \quad \text{if} \quad \theta_{\Omega,r_0}^*(t) \rightarrow 0,$$

and also

$$\gamma_0 \rightarrow 0 \quad \text{if either} \quad \alpha = \frac{r_0}{r_\Omega} \rightarrow +\infty \quad \text{or} \quad \alpha = \frac{r_0}{r_\Omega} \rightarrow 0.$$

The case  $t = 0$  is special: In this case, it is easily seen from (32) that  $\alpha^{-t/N} = 1$ , and thus  $\gamma_0$  stays bounded away from 0, even if the ratio  $r_0/r_\Omega$  diverges. We point out that for  $t = 0$ , the measure density condition becomes

$$\theta_{\Omega,r_0}^*(0) := \inf \left\{ \frac{|B_r(x) \setminus \Omega|}{|B_r(x)|} : x \in \partial\Omega, 0 < r \leq r_0 \right\} > 0,$$

which sounds like a uniform cone condition (but it is actually weaker than this, see [12, Theorem 4.1]).

On the other hand, from the previous theorem we have that

$$r_\Omega \leq \lim_{\gamma \searrow 0} R_{p,\gamma}(\Omega) \leq 6\sqrt{N}r_\Omega,$$

for every open set with a positive measure density index  $\theta_{\Omega,r_0}^*(t)$ . In particular, for the Gallagher capacity inradius, from Proposition 4.15 we get the following corollary.

**Corollary 5.5.** *Let  $1 \leq p \leq N$  and let  $\Omega \subsetneq \mathbb{R}^N$  be an open set such that  $\theta_{\Omega,r_0}^*(t) > 0$  for some  $r_0 > 0$  and  $t \geq 0$ . Then,*

$$R_p^G(\Omega) \leq 6\sqrt{N}r_\Omega.$$

**5.2. Some consequences.** We now highlight some interesting consequences which can be drawn from Theorem 5.2. We start with the following two-sided estimate on the sharp Poincaré–Sobolev constants.

**Corollary 5.6.** *Let  $1 \leq p \leq N$ , and let  $q \geq p$  be such that*

$$\begin{cases} q < p^*, & \text{if } 1 \leq p < N, \\ q < \infty, & \text{if } p = N. \end{cases}$$

*Let  $\Omega \subseteq \mathbb{R}^N$  be an open set such that  $\theta_{\Omega,r_0}^*(t) > 0$  for some  $r_0 > 0$  and  $t \geq 0$ . Then, we have*

$$\lambda_{p,q}(\Omega) > 0 \quad \iff \quad r_\Omega < +\infty.$$

*Moreover, if we have*

$$t \leq t_0, \quad \theta_{\Omega,r_0}^*(t) \geq \theta > 0 \quad \text{and} \quad 0 < \kappa_1 \leq \frac{r_0}{r_\Omega} \leq \kappa_2,$$

then there holds

$$\frac{c}{r_\Omega^{\frac{p-N+N\frac{p}{q}}{q}}} \leq \lambda_{p,q}(\Omega) \leq \frac{\lambda_{p,q}(B_1)}{r_\Omega^{\frac{p-N+N\frac{p}{q}}{q}}}, \tag{33}$$

for some  $c = c(N, p, q, t_0, \theta, \kappa_1, \kappa_2) > 0$ . Finally, if  $t_0 = 0$ , the previous constant can be taken independent of  $\kappa_2$ .

*Proof.* We directly prove the two-sided estimate (33). The rightmost inequality trivially follows from the monotonicity of  $\lambda_{p,q}$  with respect to the set inclusion and its scaling properties.

For the leftmost one, we use that

$$\gamma \sigma_{N,p,q} \left( \frac{1}{R_{p,\gamma}(\Omega)} \right)^{p-N+N\frac{p}{q}} \leq \lambda_{p,q}(\Omega), \quad \text{for every } 0 < \gamma < 1,$$

thanks to [2, Main Theorem & Theorem 6.1]. We choose  $\gamma = \gamma_0/2$ , with  $\gamma_0$  as in Theorem 5.2, and use (27) to estimate  $R_{p,\gamma_0/2}(\Omega)$  from above with  $r_\Omega$ . This gives

$$\frac{\gamma_0 \sigma_{N,p,q}}{2} \left( \frac{1}{6\sqrt{N}} \right)^{p-N+N\frac{p}{q}} \frac{1}{r_\Omega^{\frac{p-N+N\frac{p}{q}}{q}}} \leq \lambda_{p,q}(\Omega).$$

By recalling the explicit expression (32) of  $\gamma_0$ , we get the desired estimate with a constant  $c$  only depending on the claimed quantities.  $\square$

**Remark 5.7.** We remark that the previous result extends [21, Proposition 2.1]. Apart from admitting a general subcritical exponent  $q \geq p$ , we provide a quantitative lower bound on the relevant sharp Poincaré-Sobolev constant. Moreover, we notice that our measure density condition is much weaker than the cone condition required in [21] (see Lemma 5.13 below).

**Remark 5.8.** In the case  $p > N$ , the previous result holds for *every* open set, and it has been obtained by Maz’ya, see [17, Theorem 11.4.1]. For different proofs, see also [2, Theorem 1.3], [6, Corollary 5.9], [20, Theorem 1.4.1], and [23, Theorem 1.1].

In turn, from Corollary 5.6 we can obtain the equivalence between different sharp Poincaré constants. For example, if for  $p = 2$  we use the distinguished notation

$$\lambda(\Omega) = \lambda_2(\Omega) = \inf_{\varphi \in C_0^\infty(\Omega)} \left\{ \int_\Omega |\nabla \varphi|^2 dx : \|\varphi\|_{L^2(\Omega)} = 1 \right\},$$

and we recall the definition of Cheeger’s constant

$$h(\Omega) = \inf \left\{ \frac{\mathcal{H}^{N-1}(\partial E)}{|E|} : E \Subset \Omega \text{ open set with smooth boundary} \right\},$$

we can get at first the Buser-type inequality announced in the introduction.

**Corollary 5.9** (Buser-type inequality). *Let  $\Omega \subseteq \mathbb{R}^N$  be an open set such that  $\theta_{\Omega,r_0}^*(t) > 0$ , for some  $r_0 > 0$  and  $t \geq 0$ . Let us suppose that*

$$t \leq t_0, \quad \theta_{\Omega,r_0}^*(t) \geq \theta > 0 \quad \text{and} \quad 0 < \kappa_1 \leq \frac{r_0}{r_\Omega} \leq \kappa_2.$$

Then, we have

$$\lambda(\Omega) \leq C \left( h(\Omega) \right)^2,$$

for some  $C = C(N, t_0, \theta, \kappa_1, \kappa_2) > 0$ . Finally, if  $t_0 = 0$ , the previous constant can be taken independent of  $\kappa_2$ .

*Proof.* The condition on  $r_0/r_\Omega$  implies that  $r_\Omega < +\infty$  in particular. We recall that  $\lambda_{1,1}(\Omega) = h(\Omega)$  (see [16, Theorem 2.1.3] for this fact). Thus, from Corollary 5.6 with  $p = q = 1$ , we get

$$h(\Omega) \geq \frac{c}{r_\Omega},$$

with the constant  $c$  depending only on  $N, t_0, \theta, \kappa_1$ , and  $\kappa_2$ . By combining this lower bound with the classical estimate

$$\lambda(\Omega) \leq \frac{\lambda(B_1)}{r_\Omega^2},$$

recalled in the introduction, we obtain the claimed inequality with

$$C = \frac{\lambda(B_1)}{c^2}.$$

This concludes the proof.  $\square$

More generally, we recall that we have the following Cheeger–type inequality:

$$\left(\frac{p}{q}\right)^q \left(\lambda_p(\Omega)\right)^{\frac{q}{p}} \leq \lambda_q(\Omega), \quad \text{for } q > p,$$

which generalizes (5), see for example [7, Proposition 2.2]. With a similar argument as above, we may “reverse” this estimate and infer the following equivalence between sharp Poincaré constants.

**Corollary 5.10** (Generalized Buser–type inequality). *Let  $\Omega \subseteq \mathbb{R}^N$  be an open set such that  $\theta_{\Omega, r_0}^*(t) > 0$  for some  $r_0 > 0$  and  $t \geq 0$ . Then, for every  $1 \leq p < q \leq N$  we have*

$$\lambda_p(\Omega) > 0 \quad \iff \quad \lambda_q(\Omega) > 0.$$

Moreover, if we have

$$t \leq t_0, \quad \theta_{\Omega, r_0}^*(t) \geq \theta > 0 \quad \text{and} \quad 0 < \kappa_1 \leq \frac{r_0}{r_\Omega} \leq \kappa_2,$$

then

$$\lambda_q(\Omega) \leq C \left(\lambda_p(\Omega)\right)^{\frac{q}{p}},$$

for some  $C = C(N, p, q, t_0, \theta, \kappa_1, \kappa_2) > 0$ . Finally, if  $t_0 = 0$ , the previous constant can be taken independent of  $\kappa_2$ .

**5.3. Example: Uniform exterior funnel condition.** In this final subsection, we give a class of open sets which satisfy the measure density condition previously exposed.

**Definition 5.11.** We fix an exponent  $0 < \beta \leq 1$ , two positive real numbers  $\delta$  and  $h$ , and a direction  $\omega \in \mathbb{S}^{N-1}$ . We refer to  $\beta$ –funnel of axis  $\omega$  with height  $h$  and opening  $\delta$  as the following set

$$\mathcal{F}_\beta(\omega; \delta, h) = \{x \in \mathbb{R}^N : \delta |x - \langle x, \omega \rangle \omega|^\beta \leq \langle x, \omega \rangle \leq h\}.$$

We then say that an open set  $\Omega \subsetneq \mathbb{R}^N$  satisfies a *uniform exterior  $\beta$ –funnel condition with height  $h_0$  and opening  $\delta$*  if for every  $x \in \partial\Omega$  there exists a direction  $\omega_x \in \mathbb{S}^{N-1}$  with the property that

$$x + \mathcal{F}_\beta(\omega_x; \delta, h_0) \subseteq \mathbb{R}^N \setminus \Omega.$$




FIGURE 2. In the bold line, the profile of a  $\beta$ -funnel, with  $\beta = 2/5$ , opening  $\delta = 1/2$ , and height  $h_0 = 1/2$ . The circle in the dashed line has radius  $r_0$  given by (34).

**Remark 5.12.** One can easily verify that  $\mathcal{F}_\beta(\omega; \delta, h) \subseteq \mathcal{F}_\beta(\omega; \delta, h_0)$  for every  $0 < h \leq h_0$ . Thus, for an open set  $\Omega \subseteq \mathbb{R}^N$  satisfying the uniform exterior  $\beta$ -funnel condition with height  $h_0$  and opening  $\delta$ , we actually have that for every  $x \in \partial\Omega$ , there exists a direction  $\omega_x \in \mathbb{S}^{N-1}$  such that

$$x + \mathcal{F}_\beta(\omega_x; \delta, h) \subseteq \mathbb{R}^N \setminus \Omega, \quad \text{for every } 0 < h \leq h_0.$$

We have the following technical result, which illustrates concrete cases of applicability of the results of the previous subsection.

**Lemma 5.13.** *Let  $\Omega \subsetneq \mathbb{R}^N$  be an open set satisfying a uniform exterior  $\beta$ -funnel condition with height  $h_0$  and opening  $\delta$ . Then, for*

$$r_0 = \sqrt{h_0^2 + \left(\frac{h_0}{\delta}\right)^{\frac{2}{\beta}}} \quad \text{and} \quad t = (N-1) \left(\frac{1}{\beta} - 1\right), \quad (34)$$

we have

$$\theta_{\Omega, r_0}^*(t) \geq c_{N, \beta} \left( \sqrt{1 + \left(\frac{h_0^{1-\beta}}{\delta}\right)^{\frac{2}{\beta}}} \right)^{-N} \left(\frac{h_0^{1-\beta}}{\delta}\right)^{\frac{N-1}{\beta}}, \quad (35)$$

for an explicit positive constant  $c_{N, \beta}$ .

*Proof.* For every  $h \in (0, h_0]$ , we define the radius

$$r = \psi(h) := \sqrt{h^2 + \left(\frac{h}{\delta}\right)^{\frac{2}{\beta}}}.$$

Observe that this establishes a bijection between the intervals  $(0, h_0]$  and  $(0, r_0]$ , where  $r_0 := \psi(h_0)$  is the same as in (34). Thus, for every  $0 < r \leq r_0$ , we have

$$\mathcal{F}_\beta(\omega; \delta, \psi^{-1}(r)) \subseteq B_r(0), \quad \text{for every } \omega \in \mathbb{S}^{N-1}.$$

In particular, for every  $x \in \partial\Omega$  and every  $0 < r \leq r_0$ , by definition and by Remark 5.12, we have

$$|B_r(x) \setminus \Omega| \geq |(x + \mathcal{F}_\beta(\omega_x; \delta, \psi^{-1}(r))) \setminus \Omega| = |\mathcal{F}_\beta(\mathbf{e}_N; \delta, \psi^{-1}(r))|. \quad (36)$$

The last volume can be computed by using Fubini's theorem:

$$\begin{aligned} |\mathcal{F}_\beta(\mathbf{e}_N; \delta, \psi^{-1}(r))| &= \frac{\omega_{N-1}}{\delta^{\frac{N-1}{\beta}}} \int_0^{\psi^{-1}(r)} \tau^{\frac{N-1}{\beta}} d\tau \\ &= \frac{\omega_{N-1}}{\delta^{\frac{N-1}{\beta}}} \frac{\beta}{N-1+\beta} (\psi^{-1}(r))^{\frac{N-1+\beta}{\beta}}. \end{aligned} \quad (37)$$

In order to give a lower bound for the last term, we observe that

$$\psi(h) = \sqrt{h^2 + \left(\frac{h}{\delta}\right)^{\frac{2}{\beta}}} \leq h \sqrt{1 + \left(\frac{h_0^{1-\beta}}{\delta}\right)^{\frac{2}{\beta}}}, \quad \text{for } 0 < h \leq h_0,$$

so that

$$h \leq \psi^{-1} \left( h \sqrt{1 + \left(\frac{h_0^{1-\beta}}{\delta}\right)^{\frac{2}{\beta}}} \right), \quad \text{for every } 0 < h \leq h_0.$$

By introducing

$$s = h \sqrt{1 + \left(\frac{h_0^{1-\beta}}{\delta}\right)^{\frac{2}{\beta}}} \in \left( 0, h_0 \sqrt{1 + \left(\frac{h_0^{1-\beta}}{\delta}\right)^{\frac{2}{\beta}}} \right] = (0, \psi(h_0)] = (0, r_0],$$

in particular, we get

$$\psi^{-1}(s) \geq s \frac{1}{\sqrt{1 + \left(\frac{h_0^{1-\beta}}{\delta}\right)^{\frac{2}{\beta}}}}, \quad \text{for every } s \in (0, r_0].$$

For every  $0 < r \leq r_0$ , we get from (36) and (37) that

$$\begin{aligned} \left(\frac{r_0}{r}\right)^{(N-1)\left(\frac{1}{\beta}-1\right)} \frac{|B_r(x) \setminus \Omega|}{|B_r(x)|} &\geq \frac{c_{N,\beta}}{\delta^{\frac{N-1}{\beta}}} \left( \sqrt{h_0^2 + \left(\frac{h_0}{\delta}\right)^{\frac{2}{\beta}}} \right)^{(N-1)\left(\frac{1}{\beta}-1\right)} \\ &\quad \times \left( \sqrt{1 + \left(\frac{h_0^{1-\beta}}{\delta}\right)^{\frac{2}{\beta}}} \right)^{\frac{1-N-\beta}{\beta}}, \end{aligned}$$

where

$$c_{N,\beta} = \frac{\omega_{N-1}}{\omega_N} \frac{\beta}{N-1+\beta}.$$

We now use that

$$\sqrt{h_0^2 + \left(\frac{h_0}{\delta}\right)^{\frac{2}{\beta}}} = h_0 \sqrt{1 + \left(\frac{h_0^{1-\beta}}{\delta}\right)^{\frac{2}{\beta}}},$$

and thus, from the previous estimate we get

$$\theta_{\Omega, r_0}^*(t) \geq c_{N,\beta} \left( \sqrt{1 + \left(\frac{h_0^{1-\beta}}{\delta}\right)^{\frac{2}{\beta}}} \right)^{-N} \left( \frac{h_0^{1-\beta}}{\delta} \right)^{\frac{N-1}{\beta}}.$$

This concludes the proof.  $\square$

**Remark 5.14.** We observe that for  $\beta = 1$ , the uniform exterior  $\beta$ -funnel condition boils down to a uniform exterior *cone* condition. In this case, in the previous lemma we have  $t = 0$ , and we get the condition  $\theta_{\Omega, r_0}^*(0) > 0$ , which amounts to the classical measure density condition

$$\inf \left\{ \frac{|B_r(x) \setminus \Omega|}{|B_r(x)|} : x \in \partial\Omega, 0 < r \leq r_0 \right\} > 0.$$

In this case, the lower bound in (35) depends only on the opening of the cone  $\delta$ , as it is natural. More precisely, we get

$$\theta_{\Omega, r_0}^*(0) \geq c_N \frac{\delta}{(\delta^2 + 1)^{\frac{N}{2}}}.$$

**Acknowledgments.** We thank an anonymous referee for his comments on a preliminary version of the paper. We are grateful to Vladimir Bobkov for drawing our attention to the papers [10, 11].

F. B. is a member of the *Gruppo Nazionale per l'Analisi Matematica, la Probabilità e le loro Applicazioni* (GNAMPA) of the Istituto Nazionale di Alta Matematica (INdAM) and partially supported by the “INdAM - GNAMPA Project *Ottimizzazione Spettrale, Geometrica e Funzionale*”, CUP E5324001950001. L. B. has been financially supported by the *Fondo di Ateneo per la Ricerca FAR 2023* and the *Fondo per l'Incentivazione alla Ricerca Dipartimentale FIRD 2024* of the University of Ferrara.

This paper has been finalized during the XXXIV CONVEGNO NAZIONALE DI CALCOLO DELLE VARIAZIONI and the workshop CALCULUS OF VARIATIONS - OLD PROBLEMS AND NEW DIRECTIONS, respectively held in Riccione and Edinburgh in February 2025. The hosting institutions and the organizers of both the events are gratefully acknowledged.

## REFERENCES

- [1] F. Bozzola, *Topological and Capacitary Methods for Poincaré Inequalities*, Ph.D. Thesis, Università di Parma, 2025, available at <https://cvgmt.sns.it/paper/7018/>.
- [2] F. Bozzola and L. Brasco, The role of topology and capacity in some bounds for principal frequencies, *J. Geom. Anal.*, **34** (2024), Paper No. 299.
- [3] F. Bozzola and L. Brasco, *Capacitary inradius and Poincaré-Sobolev inequalities*, *ESAIM Control Optim. Calc. Var.*, **31** (2025), Paper No. 26, 38 pp.
- [4] L. Brasco, *Handbook of Calculus of Variations for Absolute Beginners*, Unitext, 163, La Mat. per il 3+2, Springer, Cham, 2025.
- [5] L. Brasco, *On principal frequencies and isoperimetric ratios in convex sets*, *Ann. Fac. Sci. Toulouse Math. (6)*, **29** (2020), 977-1005.
- [6] L. Brasco, F. Prinari and A. Zagati, *Sobolev embeddings and distance functions*, *Adv. Calc. Var.*, **17** (2024), 1365-1398.
- [7] L. Briani, G. Buttazzo and F. Prinari, *On a class of Cheeger inequalities*, *Ann. Mat. Pura Appl. (4)*, **202** (2023), 657-678.
- [8] P. Buser, *A note on the isoperimetric constant*, *Ann. Sci. École Norm. Sup. (4)*, **15** (1982), 213-230.
- [9] P. Buser, *On Cheeger's inequality  $\lambda_1 \geq h^2/4$* , *Geometry of the Laplace Operator (Proc. Sympos. Pure Math., Univ. Hawaii, Honolulu, Hawaii, 1979)*, *Proc. Sympos. Pure Math.*, XXXVI. American Mathematical Society, Providence, RI, (1980), 29-77.
- [10] A.-K. Gallagher, *On the Poincaré Inequality on Open Sets in  $\mathbb{R}^n$* , *Comput. Methods Funct. Theory*, **25** (2025), 213-238.
- [11] A.-K. Gallagher, *Equivalence between validity of the  $p$ -Poincaré inequality and finiteness of the strict  $p$ -capacitary inradius*, preprint, 2024. [arXiv:2404.19207v4](https://arxiv.org/abs/2404.19207v4).

- [12] P. Górká and P. Lefelbajn, [Cone property and measure density condition](#), *Math. Inequal. Appl.*, **27** (2024), 311-325.
- [13] T. Horiuchi, [On the relative  \$p\$ -capacity](#), *J. Math. Soc. Japan*, **43** (1991), 605-617.
- [14] M. Ledoux, [Spectral gap, logarithmic Sobolev constant, and geometric bounds](#), *Surv. Differ. Geom.*, **9**, Int. Press, Somerville, MA, (2004), 219-240.
- [15] M. Ledoux, [A simple analytic proof of an inequality by P. Buser](#), *Proc. Amer. Math. Soc.*, **121** (1994), 951-959.
- [16] V. Maz'ya, *Sobolev Spaces with Applications to Elliptic Partial Differential Equations*, Second, Revised and Augmented Edition, Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], **342**, Springer, Heidelberg, 2011.
- [17] V. Maz'ya, *Sobolev Spaces*, Translated from the Russian by T. O. Shaposhnikova. Springer Ser. Soviet Math. Springer-Verlag, Berlin, 1985.
- [18] V. Maz'ya and M. Shubin, [Can one see the fundamental frequency of a drum?](#), *Lett. Math. Phys.*, **74** (2005), 135-151.
- [19] E. Parini, [Reverse Cheeger inequality for planar convex sets](#), *J. Convex Anal.*, **24** (2017), 107-122.
- [20] G. Poliquin, [Principal frequency of the  \$p\$ -Laplacian and the inradius of Euclidean domains](#), *J. Topol. Anal.*, **7** (2015), 505-511.
- [21] P. Souplet, [Geometry of unbounded domains, Poincaré inequalities and stability in semilinear parabolic equations](#), *Comm. Partial Differential Equations*, **24** (1999), 951-973.
- [22] G. Talenti, [Best constant in Sobolev inequality](#), *Ann. Mat. Pura Appl. (4)*, **110** (1976), 353-372.
- [23] A. Vitolo,  [\$H^{1,p}\$ -eigenvalues and  \$L^\infty\$ -estimates in quasicylindrical domains](#), *Commun. Pure Appl. Anal.*, **10** (2011), 1315-1329.

Received March 2025; revised June 2025; early access June 2025.