



Asthma and climate change: unveiling hidden vulnerabilities

Francesco Maria Galassi^{1,2} · Elena Varotto³ · Mauro Vaccarezza⁴ · Domenico Ribatti⁵

Received: 2 September 2025 / Accepted: 21 October 2025
© The Author(s) 2025

Keywords Asthma · Climate change · Air pollution · Pollen · Respiratory mucosa · Child

Climate change is now firmly established as a determinant of respiratory health, with asthma among the conditions most sensitive to its shifting contours [1]. Rising temperatures and elevated atmospheric carbon dioxide drive longer, more intense pollen seasons [2], while extreme events such as wildfires, droughts, and storms degrade air quality. Importantly, air quality and, consequently, respiratory health are also compromised by anthropogenic combustion processes (e.g., fossil fuel burning, vehicular traffic, industrial emissions), which generate pollutants directly responsible for both climate change and respiratory morbidity [3]. The result is a discernible rise in asthma exacerbations, particularly among children and women, whose vulnerability reflects both biological and social factors. This vulnerability is not uniform across the life course: male children have higher asthma prevalence in childhood, whereas after puberty women exhibit higher prevalence and severity (the “sex reversal”), likely mediated by hormonal, immunologic, and gender-related factors [4]. Recent syntheses further underscore these converging risks in the climate–allergy–asthma nexus, which is the interconnected effects of meteorological shifts, allergens, and airway inflammation [5, 16]. Recent analyses have traced clear associations between meteorological instability—oscillations in temperature, extremes of humidity, and erratic

wind patterns—and surges in paediatric asthma admissions, highlighting how sudden shifts in climate parameters can precipitate acute respiratory crises [5]. For instance, modelling suggests that climate-related increases in ozone could raise summertime paediatric asthma emergency visits by ~7.3% in U.S. settings [6]. Moreover, under a 2 °C warming scenario, approximately 34,500 additional paediatric asthma cases per year are projected globally, approaching ~90,000 under higher warming trajectories [7]. The cumulative effect of these exposures weighs most heavily upon children, who possess fewer physiological reserves, narrower airways, and developing lungs that are more readily harmed by air pollutants and aeroallergens, thereby magnifying both frequency and severity of asthma exacerbations in this vulnerable group. Alongside epidemiological insights, there is scope for methodological renewal. Precision-cut lung slices (PCLS), air–liquid interface models (ALI), and lung-on-a-chip systems offer novel means of interrogating how pollutants and bioaerosols, modulated by climate change, interact with airway epithelia [2]. PCLS preserve the structural and multicellular complexity of human lung tissue, enabling *ex vivo* modelling of airway physiology and injury–repair dynamics [9]. ALI cultures allow primary airway epithelium to differentiate and recapitulate *in vivo*-like exposure to airborne stimuli [10]. Lung-on-a-chip platforms reproduce key features of the pulmonary microenvironment under dynamic flow, supporting mechanistic studies of pollutant and allergen effects [11]. While powerful, their routine use remains limited by cost, technical complexity, and the need for specialised infrastructure. There is, however, a further dimension that has attracted scant notice. Beyond heat and allergens lies the insidious role of climate-amplified chemical pollutants in undermining epithelial barrier integrity. Airborne particulates, volatile organic compounds, phthalates, and polycyclic aromatic hydrocarbons have each been implicated in erosion of mucosal defences [2, 5]. Equally important are gaseous pollutants such as nitrogen dioxide (NO₂), sulfur dioxide (SO₂), carbon monoxide (CO), and ozone (O₃), all associated

✉ Francesco Maria Galassi
francesco.galassi@biol.uni.lodz.pl

¹ Department of Anthropology, University of Lodz, Łódź, Poland

² School of Biomedicine, The University of Adelaide, Adelaide, SA, Australia

³ College of Humanities, Arts and Social Sciences, Flinders University, Adelaide, SA, Australia

⁴ Curtin Medical School, Faculty of Health Sciences, Curtin University, Bentley, WA, Australia

⁵ Department of Translational Biomedicine and Neuroscience, University of Bari Medical School, Bari, Italy

with asthma incidence and exacerbations [3]. The “epithelial barrier hypothesis”, here applied to paediatric asthma in the context of climate change, posits that environmental stressors disrupt epithelial integrity, increase permeability, and promote chronic inflammation, a framework that now warrants systematic mechanistic and epidemiological investigation to elucidate how pollutants, aeroallergens, and climate-related extremes converge on barrier dysfunction [12]. In conclusion, strategies to mitigate climate-related asthma risk must extend beyond allergen calendars and emergency preparedness. Primary prevention measures include exclusive breastfeeding, associated with reduced risk of wheezing and asthma in predisposed children [13]; comprehensive home-based interventions to reduce allergens, dampness, and mould [14]; and rural/ “farm-like” lifestyle exposures that enhance early-life microbial diversity and appear protective against allergic sensitisation and asthma [15]. At the population level, transitions to clean energy, reductions in fossil-fuel combustion, and urban greening with low-allergenic vegetation offer co-benefits for climate mitigation and respiratory health [3, 16]. Research integrating barrier biology with advanced *in vitro* modelling could illuminate critical early-life tipping points where preventive interventions might prove most effective. Policy frameworks, likewise, should reckon not only with the obvious meteorological threats, but also with the more subtle barrier-compromising pollutants whose influence is magnified by our changing climate.

Declarations

Conflict of interest The authors have no conflict of interest.

Ethics approval Not applicable.

Human and animal rights Not applicable.

Informed consent Not applicable.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article’s Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article’s Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Goshua A, Sampath V, Efobi JA, Nadeau K (2023) The role of climate change in asthma. *Adv Exp Med Biol* 1426:25–41. https://doi.org/10.1007/978-3-031-32259-4_2
- D’Amato G, Chong-Neto HJ, Monge Ortega OP, Vitale C, Ansotegui I, Rosario N et al (2020) The effects of climate change on respiratory allergy and asthma induced by pollen and mold allergens. *Allergy* 75:2219–2228. <https://doi.org/10.1111/all.14476>
- Guarnieri M, Balmes JR (2014) Outdoor air pollution and asthma. *Lancet* 383(9928):1581–1592. [https://doi.org/10.1016/S0140-6736\(14\)60617-6](https://doi.org/10.1016/S0140-6736(14)60617-6)
- Shah R, Newcomb DC (2018) Sex bias in asthma prevalence and pathogenesis. *Front Immunol* 9:2997. <https://doi.org/10.3389/fimmu.2018.02997>
- Kelly G, Idubor OI, Binney S et al (2023) The impact of climate change on asthma and allergic immunologic disease. *Curr Allergy Asthma Rep* 23:453–461. <https://doi.org/10.1007/s11882-0723-01093-y>
- Knowlton K, Rosenthal JE, Hogrefe C, Lynn B, Gaffin S, Goldberg R, Rosenzweig C, Civerolo K, Ku JY, Kinney PL (2004) Assessing ozone-related health impacts under a changing climate. *Environ Health Perspect* 112(15):1557–1563. <https://doi.org/10.1289/ehp.7163>
- Bignier C, Havet L, Brisoux M, Omeiche C, Misra S, Gonsard A, Drummond D (2025) Climate change and children’s respiratory health. *Paediatr Respir Rev* 53:64–73. <https://doi.org/10.1016/j.prrv.2024.07.002>
- Burbank AJ (2025) Climate change and the future of allergies and asthma. *Curr Allergy Asthma Rep* 25(1):20. <https://doi.org/10.1007/s11882-025-01201-0>
- Alsafadi HN, Staab-Weijnitz CA, Lehmann M et al (2017) An *ex vivo* model to induce early fibrosis-like changes in human precision-cut lung slices. *Am J Physiol Lung Cell Mol Physiol* 312(6):L896–L902. <https://doi.org/10.1152/ajplung.00084.2017>
- Pezzulo AA, Starner TD, Scheetz TE et al (2011) The air-liquid interface and use of primary cell cultures are important to recapitulate the transcriptional profile of *in vivo* airway epithelia. *Am J Physiol Lung Cell Mol Physiol* 300(1):L25–L31. <https://doi.org/10.1152/ajplung.00256.2010>
- Huh D, Matthews BD, Mammoto A, Montoya-Zavala M, Hsin HY, Ingber DE (2010) Reconstituting organ-level lung functions on a chip. *Science* 328(5986):1662–1668. <https://doi.org/10.1126/science.1188302>
- Akdis CA (2021) Does the epithelial barrier hypothesis explain the increase in allergy, autoimmunity and other chronic conditions? *Nat Rev Immunol* 21(11):739–751. <https://doi.org/10.1038/s41577-021-00538-7>
- Kramer MS, Chalmers B, Hodnett ED et al (2001) Promotion of breastfeeding intervention trial (PROBIT): a randomized trial in the Republic of Belarus. *JAMA* 285(4):413–420. <https://doi.org/10.1001/jama.285.4.413>
- Murray CS, Foden P, Sumner H, Shepley E, Custovic A, Simpson A (2017) Preventing severe asthma exacerbations in children: a randomized trial of mite-impermeable bedcovers. *Am J Respir Crit Care Med* 196(2):150–158. <https://doi.org/10.1164/rccm.201609-1966OC>
- Stein MM, Hrusch CL, Gozdz J et al (2016) Innate immunity and asthma risk in Amish and Hutterite farm children. *N Engl J Med* 375(5):411–421. <https://doi.org/10.1056/NEJMoa1508749>
- D’Amato G, Annesi-Maesano I, Biagioni B, Lancia A, Cecchi L, D’Ovidio MC, D’Amato M (2023) New developments in climate change, air pollution, pollen allergy, and interaction with SARS-CoV-2. *Atmosphere* 14(5):848. <https://doi.org/10.3390/atmos14050848>

Publisher’s Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.