

Observation of an Isoscalar Resonance with Exotic $J^{PC} = 1^{-+}$ Quantum Numbers in $J/\psi \rightarrow \gamma\eta\eta'$

M. Ablikim,¹ M. N. Achasov,^{10,b} P. Adlarson,⁶⁸ S. Ahmed,¹⁴ M. Albrecht,⁴ R. Aliberti,²⁸ A. Amoroso,^{67a,67c} M. R. An,³² Q. An,^{64,50} X. H. Bai,⁵⁸ Y. Bai,⁴⁹ O. Bakina,²⁹ R. Baldini Ferroli,^{23a} I. Balossino,^{24a} Y. Ban,^{39,g} V. Batozskaya,^{1,37} D. Becker,²⁸ K. Begzsuren,²⁶ N. Berger,²⁸ M. Bertani,^{23a} D. Bettoni,^{24a} F. Bianchi,^{67a,67c} J. Bloms,⁶¹ A. Bortone,^{67a,67c} I. Boyko,²⁹ R. A. Briere,⁵ H. Cai,⁶⁹ X. Cai,^{1,50} A. Calcaterra,^{23a} G. F. Cao,^{1,55} N. Cao,^{1,55} S. A. Cetin,^{54a} J. F. Chang,^{1,50} W. L. Chang,^{1,55} G. Chelkov,^{29,a} C. Chen,³⁶ G. Chen,¹ H. S. Chen,^{1,55} M. L. Chen,^{1,50} S. J. Chen,³⁵ T. Chen,¹ X. R. Chen,²⁵ X. T. Chen,¹ Y. B. Chen,^{1,50} Z. J. Chen,^{20,h} W. S. Cheng,^{67c} G. Cibinetto,^{24a} F. Cossio,^{67c} J. J. Cui,⁴² X. F. Cui,³⁶ H. L. Dai,^{1,50} J. P. Dai,⁷¹ X. C. Dai,^{1,55} A. Dbeyssi,¹⁴ R. E. de Boer,⁴ D. Dedovich,²⁹ Z. Y. Deng,¹ A. Denig,²⁸ I. Denysenko,²⁹ M. Destefanis,^{67a,67c} F. De Mori,^{67a,67c} Y. Ding,³³ C. Dong,³⁶ J. Dong,^{1,50} L. Y. Dong,^{1,55} M. Y. Dong,^{1,50,55} X. Dong,⁶⁹ S. X. Du,⁷³ P. Egorov,^{29,a} Y. L. Fan,⁶⁹ J. Fang,^{1,50} S. S. Fang,^{1,55} Y. Fang,¹ R. Farinelli,^{24a} L. Fava,^{67b,67c} F. Feldbauer,⁴ G. Felici,^{23a} C. Q. Feng,^{64,50} J. H. Feng,⁵¹ M. Fritsch,⁴ C. D. Fu,¹ Y. Gao,^{39,g} Y. Gao,^{64,50} I. Garzia,^{24a,24b} P. T. Ge,⁶⁹ C. Geng,⁵¹ E. M. Gersabeck,⁵⁹ A. Gilman,⁶² K. Goetzen,¹¹ L. Gong,³³ W. X. Gong,^{1,50} W. Gradl,²⁸ M. Greco,^{67a,67c} M. H. Gu,^{1,50} Y. T. Gu,⁷⁴ C. Y. Guan,^{1,55} A. Q. Guo,²⁵ A. Q. Guo,²² L. B. Guo,³⁴ R. P. Guo,⁴¹ Y. P. Guo,^{9,f} A. Guskov,^{29,a} T. T. Han,⁴² W. Y. Han,³² X. Q. Hao,¹⁵ F. A. Harris,⁵⁷ K. K. He,⁴⁷ K. L. He,^{1,55} F. H. Heinsius,⁴ C. H. Heinz,²⁸ Y. K. Heng,^{1,50,55} C. Herold,⁵² M. Himmelreich,^{11,d} T. Holtmann,⁴ G. Y. Hou,^{1,55} Y. R. Hou,⁵⁵ Z. L. Hou,¹ H. M. Hu,^{1,55} J. F. Hu,^{48,i} T. Hu,^{1,50,55} Y. Hu,¹ G. S. Huang,^{64,50} L. Q. Huang,⁶⁵ X. T. Huang,⁴² Y. P. Huang,¹ Z. Huang,^{39,g} T. Hussain,⁶⁶ N. Hüsken,^{22,28} W. Ikegami Andersson,⁶⁸ W. Imoehl,²² M. Irshad,^{64,50} S. Jaeger,⁴ S. Janchiv,²⁶ Q. Ji,¹ Q. P. Ji,¹⁵ X. B. Ji,^{1,55} X. L. Ji,^{1,50} Y. Y. Ji,⁴² H. B. Jiang,⁴² S. S. Jiang,³² X. S. Jiang,^{1,50,55} J. B. Jiao,⁴² Z. Jiao,¹⁸ S. Jin,³⁵ Y. Jin,⁵⁸ M. Q. Jing,^{1,55} T. Johansson,⁶⁸ N. Kalantar-Nayestanaki,⁵⁶ X. S. Kang,³³ R. Kappert,⁵⁶ M. Kavatsyuk,⁵⁶ B. C. Ke,⁷³ I. K. Keshk,⁴ A. Khokkaz,⁶¹ P. Kiese,²⁸ R. Kiuchi,¹ R. Kliemt,¹¹ L. Koch,³⁰ O. B. Kolcu,^{54a} B. Kopf,⁴ M. Kuemmel,⁴ M. Kuessner,⁴ A. Kupsc,^{37,68} M. G. Kurth,^{1,55} W. Kühn,³⁰ J. J. Lane,⁵⁹ J. S. Lange,³⁰ P. Larin,¹⁴ A. Lavanaia,²¹ L. Lavezzi,^{67a,67c} Z. H. Lei,^{64,50} H. Leithoff,²⁸ M. Lellmann,²⁸ T. Lenz,²⁸ C. Li,⁴⁰ C. Li,³⁶ C. H. Li,³² Cheng Li,^{64,50} D. M. Li,⁷³ F. Li,^{1,50} G. Li,¹ H. Li,^{64,50} H. Li,⁴⁴ H. B. Li,^{1,55} H. J. Li,¹⁵ H. N. Li,^{48,i} J. L. Li,⁴² J. Q. Li,⁴ J. S. Li,⁵¹ Ke Li,¹ L. J. Li,¹ L. K. Li,¹ Lei Li,³ M. H. Li,³⁶ P. R. Li,^{31,j,k} S. Y. Li,⁵³ T. Li,⁴² W. D. Li,^{1,55} W. G. Li,¹ X. H. Li,^{64,50} X. L. Li,⁴² Xiaoyu Li,^{1,55} Z. Y. Li,⁵¹ H. Liang,^{64,50} H. Liang,²⁷ H. Liang,^{1,55} Y. F. Liang,⁴⁶ Y. T. Liang,²⁵ G. R. Liao,¹² L. Z. Liao,^{1,55} J. Libby,²¹ A. Limphirat,⁵² C. X. Lin,⁵¹ D. X. Lin,²⁵ T. Lin,¹ B. J. Liu,¹ C. X. Liu,¹ D. Liu,^{14,64} F. H. Liu,⁴⁵ Fang Liu,¹ Feng Liu,⁶ G. M. Liu,^{48,i} H. B. Liu,⁷⁴ H. M. Liu,^{1,55} Huanhuan Liu,¹ Huihui Liu,¹⁶ J. B. Liu,^{64,50} J. L. Liu,⁶⁵ J. Y. Liu,^{1,55} K. Liu,¹ K. Y. Liu,³³ Ke Liu,¹⁷ L. Liu,^{64,50} M. H. Liu,^{9,f} P. L. Liu,¹ Q. Liu,⁵⁵ S. B. Liu,^{64,50} T. Liu,^{1,55} T. Liu,^{9,f} W. M. Liu,^{64,50} X. Liu,^{31,j,k} Y. Liu,^{31,j,k} Y. B. Liu,³⁶ Z. A. Liu,^{1,50,55} Z. Q. Liu,⁴² X. C. Lou,^{1,50,55} F. X. Lu,⁵¹ H. J. Lu,¹⁸ J. D. Lu,^{1,55} J. G. Lu,^{1,50} X. L. Lu,¹ Y. Lu,¹ Y. P. Lu,^{1,50} Z. H. Lu,¹ C. L. Luo,³⁴ M. X. Luo,⁷² T. Luo,^{9,f} X. L. Luo,^{1,50} X. R. Lyu,⁵⁵ Y. F. Lyu,³⁶ F. C. Ma,³³ H. L. Ma,¹ L. L. Ma,⁴² M. M. Ma,^{1,55} Q. M. Ma,¹ R. Q. Ma,^{1,55} R. T. Ma,⁵⁵ X. X. Ma,^{1,55} X. Y. Ma,^{1,50} Y. Ma,^{39,g} F. E. Maas,¹⁴ M. Maggiora,^{67a,67c} S. Maldaner,⁴ S. Malde,⁶² Q. A. Malik,⁶⁶ A. Mangoni,^{23b} Y. J. Mao,^{39,g} Z. P. Mao,¹ S. Marcello,^{67a,67c} Z. X. Meng,⁵⁸ J. G. Messchendorp,⁵⁶ G. Mezzadri,^{24a} H. Miao,¹ T. J. Min,³⁵ R. E. Mitchell,²² X. H. Mo,^{1,50,55} N. Yu. Muchnoi,^{10,b} H. Muramatsu,⁶⁰ S. Nakhoul,^{11,d} Y. Nefedov,²⁹ F. Nerling,^{11,d} I. B. Nikolaev,^{10,b} Z. Ning,^{1,50} S. Nisar,^{8,1} S. L. Olsen,⁵⁵ Q. Ouyang,^{1,50,55} S. Pacetti,^{23b,23c} X. Pan,^{9,f} Y. Pan,⁵⁹ A. Pathak,¹ A. Pathak,²⁷ P. Patteri,^{23a} M. Pelizaeus,⁴ H. P. Peng,^{64,50} K. Peters,^{11,d} J. Pettersson,⁶⁸ J. L. Ping,³⁴ R. G. Ping,^{1,55} S. Plura,²⁸ S. Pogodin,²⁹ R. Poling,⁶⁰ V. Prasad,^{64,50} H. Qi,^{64,50} H. R. Qi,⁵³ M. Qi,³⁵ T. Y. Qi,^{9,f} S. Qian,^{1,50} W. B. Qian,⁵⁵ Z. Qian,⁵¹ C. F. Qiao,⁵⁵ J. J. Qin,⁶⁵ L. Q. Qin,¹² X. P. Qin,^{9,f} X. S. Qin,⁴² Z. H. Qin,^{1,50} J. F. Qiu,¹ S. Q. Qu,³⁶ K. H. Rashid,⁶⁶ K. Ravindran,²¹ C. F. Redmer,²⁸ K. J. Ren,³² A. Rivetti,^{67c} V. Rodin,⁵⁶ M. Rolo,^{67c} G. Rong,^{1,55} Ch. Rosner,¹⁴ M. Rump,⁶¹ H. S. Sang,⁶⁴ A. Sarantsev,^{29,c} Y. Schelhaas,²⁸ C. Schnier,⁴ K. Schoenning,⁶⁸ M. Scodreggio,^{24a,24b} W. Shan,¹⁹ X. Y. Shan,^{64,50} J. F. Shangguan,⁴⁷ L. G. Shao,^{1,55} M. Shao,^{64,50} C. P. Shen,^{9,f} H. F. Shen,^{1,55} X. Y. Shen,^{1,55} B.-A. Shi,⁵⁵ H. C. Shi,^{64,50} R. S. Shi,^{1,55} X. Shi,^{1,50} X. D. Shi,^{64,50} J. J. Song,¹⁵ W. M. Song,^{27,1} Y. X. Song,^{39,g} S. Sosio,^{67a,67c} S. Spataro,^{67a,67c} F. Stielor,²⁸ K. X. Su,⁶⁹ P. P. Su,⁴⁷ Y.-J. Su,⁵⁵ G. X. Sun,¹ H. K. Sun,¹ J. F. Sun,¹⁵ L. Sun,⁶⁹ S. S. Sun,^{1,55} T. Sun,^{1,55} W. Y. Sun,²⁷ X. Sun,^{20,h} Y. J. Sun,^{64,50} Y. Z. Sun,¹ Z. T. Sun,⁴² Y. H. Tan,⁶⁹ Y. X. Tan,^{64,50} C. J. Tang,⁴⁶ G. Y. Tang,¹ J. Tang,⁵¹ Q. T. Tao,^{20,h} J. X. Teng,^{64,50} V. Thoren,⁶⁸ W. H. Tian,⁴⁴ Y. T. Tian,²⁵ I. Uman,^{54b} B. Wang,¹ D. Y. Wang,^{39,g} H. J. Wang,^{31,j,k} H. P. Wang,^{1,55} K. Wang,^{1,50} L. L. Wang,¹ M. Wang,⁴² M. Z. Wang,^{39,g} Meng Wang,^{1,55} S. Wang,^{9,f} T. J. Wang,³⁶ W. Wang,⁵¹ W. H. Wang,⁶⁹ W. P. Wang,^{64,50} X. Wang,^{39,g} X. F. Wang,^{31,j,k} X. L. Wang,^{9,f} Y. Wang,⁵¹ Y. D. Wang,³⁸ Y. F. Wang,^{1,50,55}

Y. Q. Wang,¹ Y. Y. Wang,^{31,j,k} Z. Wang,^{1,50} Z. Y. Wang,¹ Ziyi Wang,⁵⁵ Zongyuan Wang,^{1,55} D. H. Wei,¹² F. Weidner,⁶¹ S. P. Wen,¹ D. J. White,⁵⁹ U. Wiedner,⁴ G. Wilkinson,⁶² M. Wolke,⁶⁸ L. Wollenberg,⁴ J. F. Wu,^{1,55} L. H. Wu,¹ L. J. Wu,^{1,55} X. Wu,^{9,f} X. H. Wu,²⁷ Z. Wu,^{1,50} L. Xia,^{64,50} T. Xiang,^{39,g} H. Xiao,^{9,f} S. Y. Xiao,¹ Z. J. Xiao,³⁴ X. H. Xie,^{39,g} Y. G. Xie,^{1,50} Y. H. Xie,⁶ T. Y. Xing,^{1,55} C. F. Xu,¹ C. J. Xu,⁵¹ G. F. Xu,¹ Q. J. Xu,¹³ W. Xu,^{1,55} X. P. Xu,⁴⁷ Y. C. Xu,⁵⁵ F. Yan,^{9,f} L. Yan,^{9,f} W. B. Yan,^{64,50} W. C. Yan,⁷³ H. J. Yang,^{43,e} H. X. Yang,¹ L. Yang,⁴⁴ S. L. Yang,⁵⁵ Y. X. Yang,^{1,55} Y. X. Yang,¹² Yifan Yang,^{1,55} Zhi Yang,²⁵ M. Ye,^{1,50} M. H. Ye,⁷ J. H. Yin,¹ Z. Y. You,⁵¹ B. X. Yu,^{1,50,55} C. X. Yu,³⁶ G. Yu,^{1,55} J. S. Yu,^{20,h} T. Yu,⁶⁵ C. Z. Yuan,^{1,55} L. Yuan,² S. C. Yuan,¹ Y. Yuan,¹ Z. Y. Yuan,⁵¹ C. X. Yue,³² A. A. Zafar,⁶⁶ X. Zeng Zeng,⁶ Y. Zeng,^{20,h} A. Q. Zhang,¹ B. L. Zhang,¹ B. X. Zhang,¹ G. Y. Zhang,¹⁵ H. Zhang,⁶⁴ H. H. Zhang,²⁷ H. H. Zhang,⁵¹ H. Y. Zhang,^{1,50} J. L. Zhang,⁷⁰ J. Q. Zhang,³⁴ J. W. Zhang,^{1,50,55} J. Y. Zhang,¹ J. Z. Zhang,^{1,55} Jianyu Zhang,^{1,55} Jiawei Zhang,^{1,55} L. M. Zhang,⁵³ L. Q. Zhang,⁵¹ Lei Zhang,³⁵ P. Zhang,¹ Shulei Zhang,^{20,h} X. D. Zhang,³⁸ X. M. Zhang,¹ X. Y. Zhang,⁴⁷ X. Y. Zhang,⁴² Y. Zhang,⁶² Y. T. Zhang,⁷³ Y. H. Zhang,^{1,50} Yan Zhang,^{64,50} Yao Zhang,¹ Z. H. Zhang,¹ Z. Y. Zhang,³⁶ Z. Y. Zhang,⁶⁹ G. Zhao,¹ J. Zhao,³² J. Y. Zhao,^{1,55} J. Z. Zhao,^{1,50} Lei Zhao,^{64,50} Ling Zhao,¹ M. G. Zhao,³⁶ Q. Zhao,¹ S. J. Zhao,⁷³ Y. B. Zhao,^{1,50} Y. X. Zhao,²⁵ Z. G. Zhao,^{64,50} A. Zhemchugov,^{29,a} B. Zheng,⁶⁵ J. P. Zheng,^{1,50} Y. H. Zheng,⁵⁵ B. Zhong,³⁴ C. Zhong,⁶⁵ L. P. Zhou,^{1,55} Q. Zhou,^{1,55} X. Zhou,⁶⁹ X. K. Zhou,⁵⁵ X. R. Zhou,^{64,50} X. Y. Zhou,³² A. N. Zhu,^{1,55} J. Zhu,³⁶ K. Zhu,¹ K. J. Zhu,^{1,50,55} S. H. Zhu,⁶³ T. J. Zhu,⁷⁰ W. J. Zhu,³⁶ W. J. Zhu,^{9,f} Y. C. Zhu,^{64,50} Z. A. Zhu,^{1,55} B. S. Zou,¹ and J. H. Zou¹

(BESIII Collaboration)

- ¹*Institute of High Energy Physics, Beijing 100049, People's Republic of China*
²*Beihang University, Beijing 100191, People's Republic of China*
³*Beijing Institute of Petrochemical Technology, Beijing 102617, People's Republic of China*
⁴*Bochum Ruhr-University, D-44780 Bochum, Germany*
⁵*Carnegie Mellon University, Pittsburgh, Pennsylvania 15213, USA*
⁶*Central China Normal University, Wuhan 430079, People's Republic of China*
⁷*China Center of Advanced Science and Technology, Beijing 100190, People's Republic of China*
⁸*COMSATS University Islamabad, Lahore Campus, Defence Road, Off Raiwind Road, 54000 Lahore, Pakistan*
⁹*Fudan University, Shanghai 200443, People's Republic of China*
¹⁰*G.I. Budker Institute of Nuclear Physics SB RAS (BINP), Novosibirsk 630090, Russia*
¹¹*GSI Helmholtzcentre for Heavy Ion Research GmbH, D-64291 Darmstadt, Germany*
¹²*Guangxi Normal University, Guilin 541004, People's Republic of China*
¹³*Hangzhou Normal University, Hangzhou 310036, People's Republic of China*
¹⁴*Helmholtz Institute Mainz, Staudinger Weg 18, D-55099 Mainz, Germany*
¹⁵*Henan Normal University, Xinxiang 453007, People's Republic of China*
¹⁶*Henan University of Science and Technology, Luoyang 471003, People's Republic of China*
¹⁷*Henan University of Technology, Zhengzhou 450001, People's Republic of China*
¹⁸*Huangshan College, Huangshan 245000, People's Republic of China*
¹⁹*Hunan Normal University, Changsha 410081, People's Republic of China*
²⁰*Hunan University, Changsha 410082, People's Republic of China*
²¹*Indian Institute of Technology Madras, Chennai 600036, India*
²²*Indiana University, Bloomington, Indiana 47405, USA*
^{23a}*INFN Laboratori Nazionali di Frascati, INFN Laboratori Nazionali di Frascati, I-00044 Frascati, Italy*
^{23b}*INFN Sezione di Perugia, I-06100 Perugia, Italy*
^{23c}*University of Perugia, I-06100 Perugia, Italy*
^{24a}*INFN Sezione di Ferrara, INFN Sezione di Ferrara, I-44122 Ferrara, Italy*
^{24b}*University of Ferrara, I-44122 Ferrara, Italy*
²⁵*Institute of Modern Physics, Lanzhou 730000, People's Republic of China*
²⁶*Institute of Physics and Technology, Peace Ave. 54B, Ulaanbaatar 13330, Mongolia*
²⁷*Jilin University, Changchun 130012, People's Republic of China*
²⁸*Johannes Gutenberg University of Mainz, Johann-Joachim-Becher-Weg 45, D-55099 Mainz, Germany*
²⁹*Joint Institute for Nuclear Research, 141980 Dubna, Moscow region, Russia*
³⁰*Justus-Liebig-Universitaet Giessen, II. Physikalisches Institut, Heinrich-Buff-Ring 16, D-35392 Giessen, Germany*
³¹*Lanzhou University, Lanzhou 730000, People's Republic of China*
³²*Liaoning Normal University, Dalian 116029, People's Republic of China*
³³*Liaoning University, Shenyang 110036, People's Republic of China*
³⁴*Nanjing Normal University, Nanjing 210023, People's Republic of China*
³⁵*Nanjing University, Nanjing 210093, People's Republic of China*

- ³⁶Nankai University, Tianjin 300071, People's Republic of China
³⁷National Centre for Nuclear Research, Warsaw 02-093, Poland
³⁸North China Electric Power University, Beijing 102206, People's Republic of China
³⁹Peking University, Beijing 100871, People's Republic of China
⁴⁰Qufu Normal University, Qufu 273165, People's Republic of China
⁴¹Shandong Normal University, Jinan 250014, People's Republic of China
⁴²Shandong University, Jinan 250100, People's Republic of China
⁴³Shanghai Jiao Tong University, Shanghai 200240, People's Republic of China
⁴⁴Shanxi Normal University, Linfen 041004, People's Republic of China
⁴⁵Shanxi University, Taiyuan 030006, People's Republic of China
⁴⁶Sichuan University, Chengdu 610064, People's Republic of China
⁴⁷Soochow University, Suzhou 215006, People's Republic of China
⁴⁸South China Normal University, Guangzhou 510006, People's Republic of China
⁴⁹Southeast University, Nanjing 211100, People's Republic of China
⁵⁰State Key Laboratory of Particle Detection and Electronics, Beijing 100049, Hefei 230026, People's Republic of China
⁵¹Sun Yat-Sen University, Guangzhou 510275, People's Republic of China
⁵²Suranaree University of Technology, University Avenue 111, Nakhon Ratchasima 30000, Thailand
⁵³Tsinghua University, Beijing 100084, People's Republic of China
^{54a}Turkish Accelerator Center Particle Factory Group, Istinye University, 34010 Istanbul, Turkey
^{54b}Near East University, Nicosia, North Cyprus, Mersin 10, Turkey
⁵⁵University of Chinese Academy of Sciences, Beijing 100049, People's Republic of China
⁵⁶University of Groningen, NL-9747 AA Groningen, Netherlands
⁵⁷University of Hawaii, Honolulu, Hawaii 96822, USA
⁵⁸University of Jinan, Jinan 250022, People's Republic of China
⁵⁹University of Manchester, Oxford Road, Manchester M13 9PL, United Kingdom
⁶⁰University of Minnesota, Minneapolis, Minnesota 55455, USA
⁶¹University of Muenster, Wilhelm-Klemm-Str. 9, 48149 Muenster, Germany
⁶²University of Oxford, Keble Rd, Oxford, United Kingdom OX13RH
⁶³University of Science and Technology Liaoning, Anshan 114051, People's Republic of China
⁶⁴University of Science and Technology of China, Hefei 230026, People's Republic of China
⁶⁵University of South China, Hengyang 421001, People's Republic of China
⁶⁶University of the Punjab, Lahore-54590, Pakistan
^{67a}University of Turin and INFN, University of Turin, I-10125 Turin, Italy
^{67b}University of Eastern Piedmont, I-15121 Alessandria, Italy
^{67c}INFN, I-10125 Turin, Italy
⁶⁸Uppsala University, Box 516, SE-75120 Uppsala, Sweden
⁶⁹Wuhan University, Wuhan 430072, People's Republic of China
⁷⁰Xinyang Normal University, Xinyang 464000, People's Republic of China
⁷¹Yunnan University, Kunming 650500, People's Republic of China
⁷²Zhejiang University, Hangzhou 310027, People's Republic of China
⁷³Zhengzhou University, Zhengzhou 450001, People's Republic of China
⁷⁴Guangxi University, Nanning 530004, People's Republic of China



(Received 3 February 2022; revised 3 May 2022; accepted 27 September 2022; published 31 October 2022)

Using a sample of $(10.09 \pm 0.04) \times 10^9$ J/ψ events collected with the BESIII detector operating at the BEPCII storage ring, a partial wave analysis of the decay $J/\psi \rightarrow \gamma\eta\eta'$ is performed. The first observation of an isoscalar state with exotic quantum numbers $J^{PC} = 1^{-+}$, denoted as $\eta_1(1855)$, is reported in the process $J/\psi \rightarrow \gamma\eta_1(1855)$ with $\eta_1(1855) \rightarrow \eta\eta'$. Its mass and width are measured to be $(1855 \pm 9_{-1}^{+6})$ MeV/ c^2 and $(188 \pm 18_{-8}^{+3})$ MeV, respectively, where the first uncertainties are statistical and the second are systematic, and its statistical significance is estimated to be larger than 19σ .

DOI: 10.1103/PhysRevLett.129.192002

Published by the American Physical Society under the terms of the Creative Commons Attribution 4.0 International license. Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI. Funded by SCOAP³.

The quark model describes a conventional meson as a bound state of a quark and an antiquark. However, due to the non-Abelian nature of QCD, bound states with gluonic degrees of freedom, such as glueballs and hybrids, are also expected. The clear identification of these QCD exotics

would validate and advance our quantitative understanding of QCD. Radiative decays of the J/ψ meson provide a gluon-rich environment and are therefore regarded as one of the most promising hunting grounds for gluonic excitations [1–4].

Hybrid mesons are $q\bar{q}$ states with explicit excitations of the gluon field. They were first proposed several decades ago [5–9], and have been the source of more recent lattice QCD (LQCD) [10–13] and phenomenological QCD studies [14–17]. Models and LQCD predict that the exotic $J^{PC} = 1^{-+}$ nonet of hybrid mesons is the lightest, with a mass around 1.7–2.1 GeV/ c^2 [10,13,18]. The predicted decay widths are model dependent; most hybrids are expected to be rather broad, but some can be as narrow as 100 MeV [19]. There are currently three 1^{-+} candidates: the $\pi_1(1400)$, $\pi_1(1600)$, and $\pi_1(2015)$ [20–23], which are all isovector states. Finding an isoscalar 1^{-+} hybrid state is critical for establishing the hybrid multiplet. Decaying to $\eta\eta'$ in a P wave is expected for an isoscalar 1^{-+} hybrid state [24–26].

In this Letter, a partial wave analysis (PWA) of the process $J/\psi \rightarrow \gamma\eta\eta'$ is performed. The first observation of an isoscalar state with exotic quantum numbers $J^{PC} = 1^{-+}$, denoted as $\eta_1(1855)$, is reported with high statistical significance in the decay chain $J/\psi \rightarrow \gamma\eta_1(1855) \rightarrow \gamma\eta\eta'$. In addition, a large $J/\psi \rightarrow \gamma f_0(1500) \rightarrow \gamma\eta\eta'$ component is observed, while $J/\psi \rightarrow \gamma f_0(1710) \rightarrow \gamma\eta\eta'$ is found to be insignificant. More details are presented in a companion paper [27]. The analysis is based on $(10.09 \pm 0.04) \times 10^9$ J/ψ events accumulated with the BESIII detector [28] operating at the BEPCII storage ring. A detailed description of the BESIII detector can be found in Ref. [29].

Candidate events for the process $J/\psi \rightarrow \gamma\eta\eta'$ are selected using the criteria described in Ref. [27]. The η is reconstructed via the decay channel $\gamma\gamma$ and the η' is reconstructed via $\eta' \rightarrow \gamma\pi^+\pi^-$ and $\eta' \rightarrow \eta\pi^+\pi^-$. Backgrounds are estimated by the η' mass sidebands, with details given in Ref. [27]. For $J/\psi \rightarrow \gamma\eta\eta'$, $\eta' \rightarrow \eta\pi^+\pi^-$, the selected sample contains a total of 4788 candidate events including 391 ± 9 background events, while for $J/\psi \rightarrow \gamma\eta\eta'$, $\eta' \rightarrow \gamma\pi^+\pi^-$, there are 10 544 total events including 1336 ± 21 background events.

Using the GPUPWA framework [30], a PWA is performed for the selected candidate events from the process $J/\psi \rightarrow \gamma\eta\eta'$ with $\eta' \rightarrow \gamma\pi^+\pi^-$ and $\eta' \rightarrow \eta\pi^+\pi^-$. Quasi-two-body decay amplitudes in the sequential decay processes $J/\psi \rightarrow \gamma X$, $X \rightarrow \eta\eta'$ and $J/\psi \rightarrow \eta X$, $X \rightarrow \gamma\eta'$ and $J/\psi \rightarrow \eta' X$, $X \rightarrow \gamma\eta$ are constructed using the covariant tensor amplitudes described in Ref. [31]. The resonance X is parametrized by a relativistic Breit-Wigner (BW) propagator with constant width. The complex coefficients of the amplitudes (relative magnitudes and phases) and resonance parameters (masses and widths) are determined by an unbinned maximum likelihood fit to the data. The joint probability for observing the N events in the data sample is

$$\mathcal{L} \equiv \prod_{i=1}^N \frac{|M(\xi_i)|^2 \epsilon(\xi_i) \Phi(\xi_i)}{\sigma'}, \quad (1)$$

where $\epsilon(\xi_i)$ is the detection efficiency, $\Phi(\xi_i)$ is the standard element of phase space, and $M(\xi_i) = \sum_X A_X(\xi_i)$ is the matrix element describing the decay processes from the J/ψ to the final state $\gamma\eta\eta'$. $A_X(\xi_i)$ is the amplitude corresponding to intermediate resonance X . Details of the likelihood function construction can be found in Ref. [27]. The free parameters are optimized using MINUIT [32]. To account for background, the background contribution to the likelihood function is estimated using η' sideband events and is subtracted from the total log-likelihood value [33]. The two decay channels $J/\psi \rightarrow \gamma\eta\eta'$, $\eta' \rightarrow \eta\pi^+\pi^-$ and $J/\psi \rightarrow \gamma\eta\eta'$, $\eta' \rightarrow \gamma\pi^+\pi^-$ are combined by adding their log-likelihood values; they share the same set of masses, widths, relative magnitudes, and phases.

The set of two-body amplitudes used in the PWA is determined in three steps. First, a ‘‘PDG-optimized’’ set of amplitudes is determined. To describe the $\eta\eta'$ spectrum, all kinematically allowed resonances with $J^{PC} = 0^{++}$, 2^{++} , and 4^{++} listed in the PDG [34–36] are considered. Similarly, to describe the $\gamma\eta^{(\prime)}$ spectrum, all resonances listed in the PDG with $J^{PC} = 1^{+-}$ and 1^{--} are considered. All possible combinations of these resonances are evaluated. The statistical significance for each resonance is determined by examining the probability of the change in log-likelihood values when including and excluding this resonance in the fits, where the probability is calculated under the χ^2 distribution hypothesis taking into account the change in the number of degrees of freedom. The masses and widths of the resonances near $\eta\eta'$ threshold [$f_0(1500)$, $f_2(1525)$, $f_2(1565)$, and $f_2(1640)$] as well as those with small fit fractions ($< 3\%$) are always fixed to the PDG [34] values. The mass and width of the $f_0(2330)$, which corresponds to a clear structure around 2.3 GeV/ c^2 in the $\eta\eta'$ mass spectrum, are free parameters. All other masses and widths are also free parameters in the fit. The final PDG-optimized set of amplitudes is the combination where each included resonance has a statistical significance larger than 5σ .

In the second step, a search is performed for additional resonances with $J^{PC} = 1^{-+}$, 0^{++} , 2^{++} , 4^{++} , $1_{\gamma\eta^{(\prime)}}^{+-}$, and $1_{\gamma\eta^{(\prime)}}^{--}$ by individually adding each possibility to the PDG-optimized solution and scanning over its mass and width. The significance of each additional resonance at each mass and width is evaluated. The result indicates that a significant 1^{-+} contribution ($> 7\sigma$) is needed around 1.9 GeV in the $\eta\eta'$ system. The significances for all other additional contributions are less than 5σ . Therefore, an η_1 state is included in the PWA.

In the third step, a baseline set of amplitudes is determined that includes the η_1 state with its mass and width as free

TABLE I. The masses, widths, $\mathcal{B}(J/\psi \rightarrow \gamma X \rightarrow \gamma\eta\eta')$ or $\mathcal{B}(J/\psi \rightarrow \eta' h_1 \rightarrow \gamma\eta\eta')$ (B.F.), and statistical significances (Sig.) for each component in the baseline set of amplitudes. The first uncertainties are statistical, and the second are systematic.

Resonance	M (MeV/ c^2)	Γ (MeV)	B.F.($\times 10^{-5}$)	Sig.
$f_0(1500)$	1506	112	$1.81 \pm 0.11^{+0.19}_{-0.13}$	$> 30\sigma$
$f_0(1810)$	1795	95	$0.11 \pm 0.01^{+0.04}_{-0.03}$	11.1σ
$f_0(2020)$	$2010 \pm 6^{+6}_{-4}$	$203 \pm 9^{+13}_{-11}$	$2.28 \pm 0.12^{+0.29}_{-0.20}$	24.6σ
$f_0(2330)$	$2312 \pm 7^{+7}_{-3}$	$65 \pm 10^{+3}_{-12}$	$0.10 \pm 0.02^{+0.01}_{-0.02}$	13.2σ
$\eta_1(1855)$	$1855 \pm 9^{+6}_{-1}$	$188 \pm 18^{+3}_{-8}$	$0.27 \pm 0.04^{+0.02}_{-0.04}$	21.4σ
$f_2(1565)$	1542	122	$0.32 \pm 0.05^{+0.12}_{-0.02}$	8.7σ
$f_2(2010)$	$2062 \pm 6^{+10}_{-7}$	$165 \pm 17^{+10}_{-5}$	$0.71 \pm 0.06^{+0.10}_{-0.06}$	13.4σ
$f_4(2050)$	2018	237	$0.06 \pm 0.01^{+0.03}_{-0.01}$	4.6σ
0^{++} PHSP	$1.44 \pm 0.15^{+0.10}_{-0.20}$	15.7σ
$h_1(1415)$	1416	90	$0.08 \pm 0.01^{+0.01}_{-0.02}$	10.2σ
$h_1(1595)$	1584	384	$0.16 \pm 0.02^{+0.03}_{-0.01}$	9.9σ

parameters. The statistical significances of all resonances in the PDG-optimized set are reevaluated in the presence of the η_1 state. Resonances with significance less than 5σ are removed. The resulting baseline set of amplitudes contains a significant contribution from an isoscalar state with exotic quantum numbers $J^{PC} = 1^{-+}$, denoted as $\eta_1(1855)$. Its statistical significance is 21.4σ , and its mass and width are

$(1855 \pm 9_{\text{stat}}) \text{ MeV}/c^2$ and $(188 \pm 18_{\text{stat}}) \text{ MeV}$, respectively. In addition, the baseline set of amplitudes includes four 0^{++} resonances [$f_0(1500)$, $f_0(1810)$, $f_0(2020)$, $f_0(2330)$], two 2^{++} resonances [$f_2(1565)$, $f_2(2010)$], a nonresonant contribution modeled by a $0^{++} \eta\eta'$ system uniformly distributed in phase space (PHSP), and two 1^{+-} resonances [$h_1(1415)$, $h_1(1595)$] in the $\gamma\eta$ system. In addition, a 4^{++} resonance $f_4(2050)$ with statistical significance 4.6σ is included.

The results of the PWA with the baseline set of amplitudes, including the masses and widths of the resonances, the product branching fractions $J/\psi \rightarrow \gamma X \rightarrow \gamma\eta\eta'$ or $J/\psi \rightarrow \eta^{(\prime)} X \rightarrow \gamma\eta\eta'$, and the statistical significances, are summarized in Table I. The measured masses and widths of the $f_0(2020)$ and $f_2(2010)$ are consistent with the PDG [34] average values. The measured mass of the $f_0(2330)$, which is unestablished in the PDG [34], is consistent with the results of Ref. [35], but our measured width is 79 MeV smaller (3.4σ).

Figure 1 shows the invariant mass distributions of $M(\eta\eta')$, $M(\gamma\eta)$, and $M(\gamma\eta')$ for the data (with background subtracted) and the PWA fit projections. Figure 1 also shows the $\cos\theta_\eta$ distribution, where θ_η is the angle of the η momentum in the $\eta\eta'$ (Jacob and Wick) helicity frame [37]. This angle carries information about the spin of the particle decaying to $\eta\eta'$. Figure 2 shows the Dalitz plots for the PWA fit projection, the selected data, and the background estimated from the η' sideband.




FIG. 1. Background-subtracted data (black points) and the PWA fit projections (lines) for (a),(b),(c) the invariant mass distributions of (a) $\eta\eta'$, (b) $\gamma\eta$, and (c) $\gamma\eta'$, and (d),(e) the distribution of $\cos\theta_\eta$, where θ_η is the angle of the η momentum in the $\eta\eta'$ (Jacob and Wick) helicity frame for (d) all $\eta\eta'$ masses and (e) $\eta\eta'$ masses between 1.7 and 2.0 GeV/c^2 . The red lines are the total fit projections from the baseline PWA. The blue lines are the total fit projections from a fit excluding the η_1 component. The dashed lines for the 1^{-+} , 0^{++} , 2^{++} , 4^{++} , and 1^{+-} contributions are the coherent sums of amplitudes for each J^{PC} . Note that the process $J/\psi \rightarrow \phi\eta'$, $\phi \rightarrow \eta\eta'$ is rejected, which leads to the depletion of events around $1.02 \text{ GeV}/c^2$ in $M(\gamma\eta)$.




FIG. 2. Dalitz plots for (a) the baseline PWA, (b) the selected data, and (c) background estimated from the η' sideband. Note that the process $J/\psi \rightarrow \phi\eta'$, $\phi \rightarrow \gamma\eta$ is rejected, which leads to the depletion of events in a band around $1 \text{ GeV}^2/c^4$ in $M^2(\gamma\eta)$.

Various checks are performed to validate the existence of the $\eta_1(1855)$. The fits are carried out by assigning all other possible J^{PC} ($J \leq 4$) to the $\eta_1(1855)$, and the log-likelihoods are worse by at least 235 units ($> 30\sigma$). To probe the significance of the BW phase motion, the BW parametrization of the $\eta_1(1855)$ in the baseline PWA is replaced with an amplitude whose magnitude matches that of a BW function but with constant phase (independent of s). This alternative fit has a log-likelihood 43 units (9.2σ) worse than the baseline fit.

To visualize the agreement between the PWA fit results and data, angular moments as a function of $M(\eta\eta')$ can be calculated for data (with background subtracted) and the PWA model. For events within a given region of $M(\eta\eta')$, the $\cos\theta_\eta$ distribution can be expressed as an expansion in terms of Legendre polynomials. The coefficients, which are called the unnormalized moments of the expansion, characterize the spin of the contributing $\eta\eta'$ resonances. The moment for the k th bin of $M(\eta\eta')$ is

$$\langle Y_l^0 \rangle \equiv \sum_{i=1}^{N_k} W_i Y_l^0(\cos\theta_\eta^i). \quad (2)$$

For data, N_k is the number of observed events in the k th bin of $M(\eta\eta')$ and W_i is a weight used to implement background subtraction. For the PWA model, N_k is the number of events in a PHSP MC sample and W_i is the intensity for each event calculated in the PWA model. Neglecting $\eta\eta'$ amplitudes with spin greater than 2, and ignoring the effects of symmetrization and the presence of resonance contributions in the $\gamma\eta$ and $\gamma\eta'$ subsystems, the moments are related to the spin-0 (S), spin-1 (P), and spin-2 (D) amplitudes by [38]

$$\sqrt{4\pi}\langle Y_0^0 \rangle = S^2 + P^2 + D^2, \quad (3)$$

$$\sqrt{4\pi}\langle Y_1^0 \rangle = 2SP \cos\phi_P + 4PD \cos(\phi_P - \phi_D), \quad (4)$$

$$\sqrt{4\pi}\langle Y_2^0 \rangle = \frac{2}{\sqrt{5}}P^2 + \frac{2\sqrt{5}}{7}D^2 + 2SD \cos\phi_D, \quad (5)$$

$$\sqrt{4\pi}\langle Y_4^0 \rangle = \frac{6}{7}D^2, \quad (6)$$

where ϕ_P and ϕ_D are the phases of the P wave and D wave relative to the S wave. Figure 3 shows the moments computed for the data and the PWA model, using Eq. (2), where good data and PWA consistency can be seen. The need for the $\eta_1(1855)$ P -wave component is apparent in the $\langle Y_1^0 \rangle$ moment [Fig. 3(b)].

For the branching fraction measurements, systematic uncertainties arising from the number of J/ψ events, the pion tracking, photon detection, kinematic fit, mass resolution of the η' , and the branching fractions of $\eta' \rightarrow \eta\pi^+\pi^-$, $\eta' \rightarrow \gamma\pi^+\pi^-$, and $\eta \rightarrow \gamma\gamma$ have been estimated to be 4.8% [27].

Uncertainty associated with the PWA affects both the branching fraction measurements and the resonance parameters. The sources of uncertainty include the background estimation, the resonance description, the resonance parameters, and additional resonances. The statistical significance of the $\eta_1(1855)$ is recalculated in each fit variation.

To estimate the uncertainty due to the background estimation, alternative fits are performed using different background normalization factors and different η' sideband regions. The statistical significance of the $\eta_1(1855)$ is always above 21.1σ . The changes in the branching fractions and resonance parameters are assigned as systematic uncertainties.

Uncertainty arising from the BW parametrization is estimated by replacing the constant width Γ_0 of the BW for the threshold state $f_0(1500)$ with a mass-dependent width as described in Ref. [27]. The significance of the $\eta_1(1855)$ in this case is 21.8σ .

In the baseline fit, the resonance parameters of the $f_0(1500)$, $f_0(1810)$, $f_2(1565)$, $f_4(2050)$, $h_1(1415)(\gamma\eta)$, and $h_1(1595)(\gamma\eta)$ are fixed to PDG [34] average values.




FIG. 3. The distributions of the unnormalized moments $\langle Y_L^0 \rangle$ ($L = 0, 1, 2$, and 4) for $J/\psi \rightarrow \gamma\eta\eta'$ as functions of the $\eta\eta'$ mass. Black dots with error bars represent the background-subtracted data weighted with angular moments; the red solid lines represent the baseline fit projections; and the blue dotted lines represent the projections from a fit excluding the η_1 component.

An alternative fit is performed where resonance parameters are allowed to vary within 1 standard deviation of the PDG values [34], and the changes in the results are taken as systematic uncertainties. The statistical significance of the $\eta_1(1855)$ in this case is 20.6σ .

Uncertainties arising from possible additional resonances are estimated by adding the $f_0(1710)$, $f_2(2220)$, $f_4(2300)$, $h_1(1595)(\gamma\eta')$, and $\rho(1900)(\gamma\eta')$, which are the most significant additional resonances for each possible J^{PC} , into the baseline fit individually. The resulting changes in the measurements are assigned as systematic uncertainties. In all cases, the significance of the $\eta_1(1855)$ remains larger than 19.0σ .

Assuming all of these sources are independent, the total systematic uncertainties are ${}_{-1}^{+6}$ MeV/ c^2 and ${}_{-8}^{+3}$ MeV for the mass and width of the $\eta_1(1855)$, respectively. For the branching fraction of the $\eta_1(1855)$, the total relative systematic uncertainty is determined to be ${}_{-13.1}^{+5.9}\%$. Tables VII and VIII of Ref. [27] summarize the systematic uncertainties.

The ratios $\mathcal{B}(f_0 \rightarrow \eta\eta')/\mathcal{B}(f_0 \rightarrow \pi\pi)$ can be calculated with the branching fractions measured in this analysis and previous measurements for $J/\psi \rightarrow \gamma\pi^+\pi^-$, $\gamma\pi^0\pi^0$ [39]. The ratio $\mathcal{B}[f_0(1500) \rightarrow \eta\eta']/\mathcal{B}[f_0(1500) \rightarrow \pi\pi]$ is determined to be $(8.96_{-2.87}^{+2.95}) \times 10^{-2}$, where the error is the combined systematic and statistical uncertainties. In comparison, the upper limit on $\mathcal{B}(f_0(1710) \rightarrow \eta\eta')/\mathcal{B}(f_0(1710) \rightarrow \pi\pi)$ at 90% confidence level is determined to be 1.61×10^{-3} . The suppressed decay rate of $f_0(1710)$ into $\eta\eta'$ is further discussed in Ref. [27].

In summary, a PWA of $J/\psi \rightarrow \gamma\eta\eta'$ has been performed based on $(10.09 \pm 0.04) \times 10^9$ J/ψ events collected with the BESIII detector. An isoscalar state with exotic quantum numbers $J^{PC} = 1^{-+}$, denoted as $\eta_1(1855)$, has been observed for the first time. The statistical significance of the resonance hypothesis is estimated to be larger than 19σ . The product branching fraction $\mathcal{B}[J/\psi \rightarrow \gamma\eta_1(1855)]\mathcal{B}[\eta_1(1855) \rightarrow \eta\eta']$ is measured to be $(2.70 \pm 0.41_{-0.35}^{+0.16}) \times 10^{-6}$. Its mass and width are measured to be $(1855 \pm 9_{-1}^{+6})$ MeV/ c^2 and $(188 \pm 18_{-8}^{+3})$ MeV, respectively. The first uncertainties are statistical and the second are systematic. The mass and width of the $\eta_1(1855)$ are consistent with LQCD calculations for the 1^{-+} hybrids [13]. The observation of the isoscalar $\eta_1(1855)$, combined with previous measurements of the isovector π_1 states, provides critical information about the 1^{-+} hybrid nonet. Further studies with more production mechanisms and decay modes will help clarify the nature of the $\eta_1(1855)$.

The BESIII collaboration thanks the staff of BEPCII and the IHEP computing center for their strong support. This work is supported in part by National Key R&D Program of China under Contracts No. 2020YFA0406300, No. 2020YFA0406400; National Natural Science Foundation of China (NSFC) under Contracts No. 11625523, No. 11635010, No. 11675183, No. 11735014, No. 11822506, No. 11835012, No. 11922511, No. 11935015, No. 11935016, No. 11935018, No. 11961141012, No. 12022510, No. 12025502, No. 12035009, No. 12035013,

No. 12061131003; the Chinese Academy of Sciences (CAS) Large-Scale Scientific Facility Program; the CAS Center for Excellence in Particle Physics (CCEPP); Joint Large-Scale Scientific Facility Funds of the NSFC and CAS under Contracts No. U1732263, No. U1832207; CAS Key Research Program of Frontier Sciences under Contract No. QYZDJ-SSW-SLH040; 100 Talents Program of CAS; INPAC and Shanghai Key Laboratory for Particle Physics and Cosmology; ERC under Contract No. 758462; European Union Horizon 2020 research and innovation programme under Contract No. Marie Skłodowska-Curie Grant Agreement No. 894790; German Research Foundation DFG under Contracts No. 443159800, Collaborative Research Center CRC 1044, GRK 214; Istituto Nazionale di Fisica Nucleare, Italy; Ministry of Development of Turkey under Contract No. DPT2006K-120470; National Science and Technology fund; Olle Engkvist Foundation under Contract No. 200-0605; STFC (United Kingdom); The Knut and Alice Wallenberg Foundation (Sweden) under Contract No. 2016.0157; The Royal Society, UK under Contracts No. DH140054, No. DH160214; The Swedish Research Council; U.S. Department of Energy under Contracts No. DE-FG02-05ER41374, No. DE-SC-0012069.

^aAlso at the Moscow Institute of Physics and Technology, Moscow 141700, Russia.

^bAlso at the Novosibirsk State University, Novosibirsk, 630090, Russia.

^cAlso at the NRC “Kurchatov Institute,” PNPI, 188300, Gatchina, Russia.

^dAlso at Goethe University Frankfurt, 60323 Frankfurt am Main, Germany.

^eAlso at Key Laboratory for Particle Physics, Astrophysics and Cosmology, Ministry of Education; Shanghai Key Laboratory for Particle Physics and Cosmology; Institute of Nuclear and Particle Physics, Shanghai 200240, People’s Republic of China.

^fAlso at Key Laboratory of Nuclear Physics and Ion-beam Application (MOE) and Institute of Modern Physics, Fudan University, Shanghai 200443, People’s Republic of China.

^gAlso at State Key Laboratory of Nuclear Physics and Technology, Peking University, Beijing 100871, People’s Republic of China.

^hAlso at School of Physics and Electronics, Hunan University, Changsha 410082, China.

ⁱAlso at Guangdong Provincial Key Laboratory of Nuclear Science, Institute of Quantum Matter, South China Normal University, Guangzhou 510006, China.

^jAlso at Frontiers Science Center for Rare Isotopes, Lanzhou University, Lanzhou 730000, People’s Republic of China.

^kAlso at Lanzhou Center for Theoretical Physics, Lanzhou University, Lanzhou 730000, People’s Republic of China.

^lAlso at the Department of Mathematical Sciences, IBA, Karachi, Pakistan.

[1] M. B. Cakir and G. R. Farrar, *Phys. Rev. D* **50**, 3268 (1994).

- [2] F. E. Close, G. R. Farrar, and Z.-p. Li, *Phys. Rev. D* **55**, 5749 (1997).
- [3] A. V. Sarantsev, I. Denisenko, U. Thoma, and E. Klempt, *Phys. Lett. B* **816**, 136227 (2021).
- [4] A. Rodas, A. Pilloni, M. Albaladejo, C. Fernández-Ramírez, V. Mathieu, and A. P. Szczepaniak (Joint Physics Analysis Center), *Eur. Phys. J. C* **82**, 80 (2022).
- [5] D. Horn and J. Mandula, *Phys. Rev. D* **17**, 898 (1978).
- [6] N. Isgur and J. E. Paton, *Phys. Rev. D* **31**, 2910 (1985).
- [7] M. S. Chanowitz and S. R. Sharpe, *Nucl. Phys.* **B222**, 211 (1983).
- [8] T. Barnes, F. E. Close, and F. de Viron, *Nucl. Phys.* **B224**, 241 (1983).
- [9] F. E. Close and P. R. Page, *Nucl. Phys.* **B443**, 233 (1995).
- [10] P. Lacock, C. Michael, P. Boyle, and P. Rowland (UKQCD Collaboration), *Phys. Lett. B* **401**, 308 (1997).
- [11] C. Bernard, J. E. Hetrick, T. A. DeGrand, M. Wingate, C. DeTar, C. McNeile, S. Gottlieb, U. M. Heller, K. Rummukainen, B. Sugar, and D. Toussaint (MILC Collaboration), *Phys. Rev. D* **56**, 7039 (1997).
- [12] J. J. Dudek, *Phys. Rev. D* **84**, 074023 (2011).
- [13] J. J. Dudek, R. G. Edwards, P. Guo, and C. E. Thomas (Hadron Spectrum Collaboration), *Phys. Rev. D* **88**, 094505 (2013).
- [14] A. P. Szczepaniak and E. S. Swanson, *Phys. Rev. D* **65**, 025012 (2001).
- [15] A. P. Szczepaniak and P. Krupinski, *Phys. Rev. D* **73**, 116002 (2006).
- [16] P. Guo, A. P. Szczepaniak, G. Galata, A. Vassallo, and E. Santopinto, *Phys. Rev. D* **78**, 056003 (2008).
- [17] S. D. Bass, M. Skurzok, and P. Moskal, *Phys. Rev. C* **98**, 025209 (2018).
- [18] C. A. Meyer and E. S. Swanson, *Prog. Part. Nucl. Phys.* **82**, 21 (2015).
- [19] P. R. Page, E. S. Swanson, and A. P. Szczepaniak, *Phys. Rev. D* **59**, 034016 (1999).
- [20] C. A. Meyer and Y. Van Haarlem, *Phys. Rev. C* **82**, 025208 (2010).
- [21] E. Klempt and A. Zaitsev, *Phys. Rep.* **454**, 1 (2007).
- [22] A. Rodas, A. Pilloni, M. Albaladejo, C. Fernández-Ramírez, A. Jackura, V. Mathieu, M. Mikhasenko, J. Nys, V. Pauk, B. Ketzer, and A. P. Szczepaniak (JPAC Collaboration), *Phys. Rev. Lett.* **122**, 042002 (2019).
- [23] A. J. Woss, J. J. Dudek, R. G. Edwards, C. E. Thomas, and D. J. Wilson (Hadron Spectrum Collaboration), *Phys. Rev. D* **103**, 054502 (2021).
- [24] H.-X. Chen, Z. X. Cai, P. Z. Huang, and S. L. Zhu, *Phys. Rev. D* **83**, 014006 (2011).
- [25] P.-Z. Huang, H.-X. Chen, and S.-L. Zhu, *Phys. Rev. D* **83**, 014021 (2011).
- [26] W. I. Eshraim, C. S. Fischer, F. Giacosa, and D. Parganlija, *Eur. Phys. J. Plus* **135**, 945 (2020).
- [27] M. Ablikim *et al.* (BESIII Collaboration), companion paper, *Phys. Rev. D* **106**, 072012 (2022).
- [28] M. Ablikim *et al.* (BESIII Collaboration), *Chin. Phys. C* **46**, 074001 (2022).
- [29] M. Ablikim *et al.* (BESIII Collaboration), *Nucl. Instrum. Methods Phys. Res., Sect. A* **614**, 345 (2010).
- [30] N. Berger, B. Liu, and J. Wang, *J. Phys.* **219**, 042031 (2010).

- [31] B. S. Zou and D. V. Bugg, *Eur. Phys. J. A* **16**, 537 (2003).
- [32] F. James and M. Roos, *Comput. Phys. Commun.* **10**, 343 (1975).
- [33] C. Langenbruch, *Eur. Phys. J. C* **82**, 393 (2022).
- [34] P. A. Zyla *et al.* (Particle Data Group), *Prog. Theor. Exp. Phys.* **8**, 083C01 (2020).
- [35] D. V. Bugg, *Phys. Rep.* **397**, 257 (2004).
- [36] M. Ablikim *et al.* (BESIII Collaboration), *Phys. Rev. D* **87**, 032008 (2013).
- [37] M. Jacob and G. C. Wick, *Ann. Phys. (N.Y.)* **7**, 404 (1959).
- [38] G. Costa *et al.* (BARI-BONN-CERN-GLASGOW-LIVERPOOL-MILAN-VIENNA Collaboration), *Nucl. Phys.* **B175**, 402 (1980).
- [39] M. Ablikim *et al.* (BES Collaboration), *Phys. Lett. B* **642**, 441 (2006).