

# Inside the tumor: p53 modulates calcium homeostasis

Carlotta Giorgi<sup>†</sup>, Massimo Bonora<sup>†</sup>, and Paolo Pinton\*

Department of Morphology; Surgery and Experimental Medicine; Section of Pathology; Oncology and Experimental Biology and LTTA center; University of Ferrara; Ferrara, Italy

<sup>†</sup>These authors equally contributed to this work.

Calcium ( $\text{Ca}^{2+}$ ) is an essential element of signal transduction.  $\text{Ca}^{2+}$  signaling is involved in various cellular processes, such as proliferation and apoptosis and thus is also crucial in cancer. In particular, modulation of  $\text{Ca}^{2+}$  signaling can change cells' sensitivity to apoptotic signals, such as chemotherapeutic agents.<sup>1</sup>

Previous studies indicate that both tumor-suppressors (PML, PTEN, Bax) and oncogenes (Bcl2, Ras, Akt) are able to regulate  $\text{Ca}^{2+}$  dependent apoptosis by modulating  $\text{Ca}^{2+}$  release from the endoplasmic reticulum (ER) store.<sup>2</sup> In turn, these factors can mediate the opening of the mitochondrial permeability transition pore.<sup>3</sup>

In contrast, little information was available on the transformation-related protein 53 (TRP53), better known as p53 in the regulation of  $\text{Ca}^{2+}$ -dependent apoptosis. This oncosuppressor becomes activated in response to a myriad of stressors, and its activity is crucial for the regulation of cell death pathways. Since the p53s prominent pro-apoptotic role was demonstrated, the mechanism through which p53 mediates apoptosis has been a matter of intense study. Numerous publications have described the importance of p53 in both extrinsic and intrinsic apoptotic pathways.<sup>4</sup>

In a recent study we demonstrated that p53 is present at the ER and mitochondria-associated membranes (MAMs), where it becomes enriched upon activation. At these sites, p53 interacts with the C-terminal portion of the sarco/ER  $\text{Ca}^{2+}$ -ATPase (SERCA) pump, changing

its oxidative state and, in turn, increasing ER  $\text{Ca}^{2+}$  loading. Furthermore, p53 activation enhances ER  $\text{Ca}^{2+}$  transfer to mitochondria, thus triggering pro-apoptotic mitochondrial  $\text{Ca}^{2+}$  overload and mitochondrial morphological alterations, leading to release of pro-apoptotic factors. Consistently, pharmacological p53 inactivation, as well as naturally occurring loss-of-function mutations of p53, inhibit p53s ability to increase  $\text{Ca}^{2+}$  signaling to the mitochondria, and allow for its oncogenic function.<sup>5</sup>

Whereas most of the mechanisms concerning intracellular  $\text{Ca}^{2+}$  handling have been successfully elucidated *in vitro*, we still know very little about the actual physiological role of these processes in the context of the tumor environment. Recent advancements in intravital imaging and genetically-encoded sensor technologies have allowed us to visualize  $\text{Ca}^{2+}$  transient changes in live mice, overcoming previous technical limitations.

In order to elucidate chemoresistance signaling and, thus, to study the effect of novel drugs on the induction of apoptosis, we investigated the role of intra-tumor  $\text{Ca}^{2+}$  signaling and apoptosis in a 3D tumor mass, by intravital fluorescence microscopy of mouse tumor models (Fig. 1).<sup>6</sup>

To that end, we generated a clone deriving from  $p53^{-/-}$  mouse embryo fibroblasts (MEFs) transduced with H-RAS<sup>V12</sup> and a clone from  $p53^{-/-}$  MEFs transduced with H-RAS<sup>V12</sup> with re-introduced wild-type p53. These clones were then transplanted to a dorsal skinfold chamber, in athymic mice to allow tumor

growth. We treated the mice with aluminum phthalocyanine chloride a photosensitizer commonly used in the photodynamic therapy (PDT) of cancer. Aluminum phthalocyanine chloride accumulates in intracellular organelles, including mitochondria and ER, where it initiates the apoptotic pathway by increasing cytosolic and mitochondrial  $\text{Ca}^{2+}$  upon photoactivation.<sup>7</sup>

We found that p53 is necessary and sufficient for the control of  $\text{Ca}^{2+}$ -dependent apoptosis. In particular, we were able to demonstrate that  $\text{Ca}^{2+}$  response induced by p53 is correlated with the ability of PDT to initiate apoptosis.

To investigate whether the observed  $\text{Ca}^{2+}$  modulation was a side effect of our genetic manipulation of p53, or whether it was a result of p53s mechanism of action, we used various pharmacological and molecular approaches to mimic p53s effects on intracellular  $\text{Ca}^{2+}$  homeostasis. Independently of the mechanism, all conditions that increased  $\text{Ca}^{2+}$  responses rescue the sensitivity to apoptosis in  $p53^{-/-}$  cells, while treatments that blunted the  $\text{Ca}^{2+}$  responses in p53 wild-type background were associated with the inhibition of apoptosis as in  $p53^{-/-}$  cells.<sup>6</sup>

Finally, the methodology used in this study is compatible with all the fluorescent probes currently available to follow intracellular parameters.


It is of the utmost importance to understand all the signaling pathways deregulated in cancer cells. Understanding the molecular mechanisms underlying the

\*Correspondence to: Paolo Pinton; Email: pnp@unife.it

Submitted: 01/12/2015; Accepted: 01/14/2015

<http://dx.doi.org/10.1080/15384101.2015.1010973>

Comment on: Giorgi C, et al. Intravital imaging reveals p53-dependent cancer cell death induced by phototherapy via calcium signaling. *Oncotarget* 2015; 6(3):1435-45; PMID:25544762.



**Figure 1.** Imaging of calcium signaling and apoptosis into tumor mass. Tumors grown within the skinfold chamber are loaded with aluminum phtalocyanine and the  $Ca^{2+}$  indicator Fura-2. Phtalocyanine activation and  $Ca^{2+}$  signal recording are performed in the same situ using the microscope optics. After  $Ca^{2+}$  live imaging, apoptosis is measured by intravenous administration of a fluorescent marker measuring caspase activity.

regulation of tumor cell fate is a great challenge in cancer research and will provide further insights for the development of new approaches for the treatment and cure of malignancy.

The development of novel *in vivo* imaging techniques will facilitate an even greater understanding of the  $Ca^{2+}$  signal-function relationship in live organism and in tumors, and will be an invaluable tool in the identification and classification of new pharmacological targets, as well as in the optimization of current treatments.

#### References

1. Roderick HL, Cook SJ. Nat Rev Cancer 2008; 8:361-75; PMID:18432251; <http://dx.doi.org/10.1038/nrc2374>
2. Giorgi C, et al. Antioxid Redox Signal 2015 PMID:25557408; <http://dx.doi.org/10.1089/ars.2014.6223>
3. Bonora M, et al. Cell Cycle 2013; 12:674-83; PMID:23343770; <http://dx.doi.org/10.4161/cc.23599>
4. Vousden KH, Prives C. Cell 2009; 137:413-31; PMID:19410540; <http://dx.doi.org/10.1016/j.cell.2009.04.037>
5. Giorgi C, et al. Proc Natl Acad Sci USA 2015; 112 (6):1779-84; <http://dx.doi.org/10.1073/pnas.1410723112>
6. Giorgi C, et al. Oncotarget 2015; 6(3):1435-45; PMID:25544762.
7. Shahzidi S, et al. Photochem Photobiol Sci 2011; 10:1773-82; PMID:21881674; <http://dx.doi.org/10.1039/c1pp05169e>