

Spectroscopy of geo-neutrinos from 2056 days of Borexino data

M. Agostini,¹ S. Appel,¹ G. Bellini,² J. Benziger,³ D. Bick,⁴ G. Bonfini,⁵ D. Bravo,⁶ B. Caccianiga,² F. Calaprice,⁷ A. Caminata,⁸ P. Cavalcante,⁵ A. Chepurinov,⁹ K. Choi,¹⁰ D. D'Angelo,² S. Davini,¹¹ A. Derbin,¹² L. Di Noto,⁸ I. Drachnev,¹¹ A. Empl,¹³ A. Etenko,¹⁴ G. Fiorentini,¹⁵ K. Fomenko,¹⁶ D. Franco,¹⁷ F. Gabriele,⁵ C. Galbiati,⁷ C. Ghiano,⁸ M. Giammarchi,² M. Goeger-Neff,¹ A. Goretti,⁷ M. Gromov,⁹ C. Hagner,⁴ E. Hungerford,¹³ Aldo Ianni,⁵ Andrea Ianni,⁷ K. Jedrzejczak,¹⁸ M. Kaiser,⁴ V. Kobychyev,¹⁹ D. Korablev,¹⁶ G. Korga,⁵ D. Kryn,¹⁷ M. Laubenstein,⁵ B. Lehnert,²⁰ E. Litvinovich,^{14,21} F. Lombardi,⁵ P. Lombardi,² L. Ludhova,² G. Lukyanchenko,^{14,21} I. Machulin,^{14,21} S. Manecki,⁶ W. Maneschg,²² F. Mantovani,¹⁵ S. Marocci,¹¹ E. Meroni,² M. Meyer,⁴ L. Miramonti,² M. Misiaszek,^{18,5} M. Montuschi,¹⁵ P. Mosteiro,⁷ V. Muratova,¹² L. Oberauer,¹ M. Obolensky,¹⁷ F. Ortica,²³ K. Otis,²⁴ L. Pagani,⁸ M. Pallavicini,⁸ L. Papp,¹ L. Perasso,² A. Pocar,²⁴ G. Ranucci,² A. Razeto,⁵ A. Re,² B. Ricci,¹⁵ A. Romani,²³ R. Roncin,⁵ N. Rossi,⁵ S. Schönert,¹ D. Semenov,¹² H. Simgen,²² M. Skorokhvatov,^{14,21} O. Smirnov,¹⁶ A. Sotnikov,¹⁶ S. Sukhotin,¹⁴ Y. Suvorov,^{25,14} R. Tartaglia,⁵ G. Testera,⁸ J. Thurn,²⁰ M. Toropova,¹⁴ E. Unzhakov,¹² R.B. Vogelaar,⁶ F. von Feilitzsch,¹ H. Wang,²⁵ S. Weinz,²⁶ J. Winter,²⁶ M. Wojcik,¹⁸ M. Wurm,²⁶ Z. Yokley,⁶ O. Zaimidoroga,¹⁶ S. Zavatarelli,⁸ K. Zuber,²⁰ and G. Zuzel¹⁸

(Borexino Collaboration)

¹*Physik-Department and Excellence Cluster Universe, Technische Universität München, 85748 Garching, Germany.*

²*Dipartimento di Fisica, Università degli Studi e INFN, 20133 Milano, Italy*

³*Chemical Engineering Department, Princeton University, Princeton, NJ 08544, USA*

⁴*Institut für Experimentalphysik, Universität, 22761 Hamburg, Germany*

⁵*INFN Laboratori Nazionali del Gran Sasso, 67010 Assergi (AQ), Italy*

⁶*Physics Department, Virginia Polytechnic Institute and State University, Blacksburg, VA 24061, USA*

⁷*Physics Department, Princeton University, Princeton, NJ 08544, USA*

⁸*Dipartimento di Fisica, Università degli Studi e INFN, Genova 16146, Italy*

⁹*Lomonosov Moscow State University Skobeltsyn Institute of Nuclear Physics, 119234 Moscow, Russia.*

¹⁰*Department of Physics and Astronomy, University of Hawaii, Honolulu, HI 96822, USA.*

¹¹*Gran Sasso Science Institute (INFN), 67100 L'Aquila, Italy.*

¹²*St. Petersburg Nuclear Physics Institute NRC Kurchatov Institute, 188350 Gatchina, Russia*

¹³*Department of Physics, University of Houston, Houston, TX 77204, USA.*

¹⁴*NRC Kurchatov Institute, 123182 Moscow, Russia*

¹⁵*Dipartimento di Fisica e Scienze della Terra Università degli Studi di Ferrara e INFN, Via Saragat 1-44122, Ferrara, Ital*

¹⁶*Joint Institute for Nuclear Research, 141980 Dubna, Russia*

¹⁷*Laboratoire AstroParticule et Cosmologie, 75231 Paris cedex 13, France*

¹⁸*M. Smoluchowski Institute of Physics, Jagiellonian University, 30059 Krakow, Poland*

¹⁹*Kiev Institute for Nuclear Research, 06380 Kiev, Ukraine*

²⁰*Department of Physics, Technische Universität Dresden, 01062 Dresden, Germany.*

²¹*National Research Nuclear University MEPhI (Moscow Engineering Physics Institute), 115409 Moscow, Russia.*

²²*Max-Planck-Institut für Kernphysik, 69117 Heidelberg, Germany*

²³*Dipartimento di Chimica, Università e INFN, 06123 Perugia, Italy*

²⁴*Amherst Center for Fundamental Interactions and Physics*

Department, University of Massachusetts, Amherst, MA 01003, USA

²⁵*Physics and Astronomy Department, University of California Los Angeles (UCLA), Los Angeles, California 90095, USA.*

²⁶*Institute of Physics and Excellence Cluster PRISMA, Johannes Gutenberg-Universität Mainz, 55099 Mainz, Germany.*

(Dated: June 17, 2015)

We report an improved geo-neutrino measurement with Borexino from 2056 days of data taking. The present exposure is $(5.5 \pm 0.3) \times 10^{31}$ proton \times yr. Assuming a chondritic Th/U mass ratio of 3.9, we obtain $23.7_{-5.7}^{+6.5}(\text{stat})_{-0.6}^{+0.9}(\text{sys})$ geo-neutrino events. The null observation of geo-neutrinos with Borexino alone has a probability of 3.6×10^{-9} (5.9σ). A geo-neutrino signal from the mantle is obtained at 98% C.L. The radiogenic heat production for U and Th from the present best-fit result is restricted to the range 23-36 TW, taking into account the uncertainty on the distribution of heat producing elements inside the Earth.

PACS numbers: 13.35.Hb, 14.60.St, 26.65.+t, 95.55.Vj, 29.40.Mc

Geo-neutrinos are electron anti-neutrinos ($\bar{\nu}_e$) produced by β decays of long-lived isotopes, which are naturally present in the interior of the Earth, such as decays in the ^{238}U and ^{232}Th chains, and ^{40}K [1, 2]. Geo- $\bar{\nu}_e$

measurements have been reported by Borexino [3, 4] and KamLAND [5, 6]. Here we present improved reactor neutrino and geo- $\bar{\nu}_e$ measurements performed by Borexino. Borexino is an unsegmented liquid scintillator detector in

operation at the underground Gran Sasso National Laboratory, Italy [8]. In liquid scintillator detectors $\bar{\nu}_e$ are detected via the inverse β decay process, $\bar{\nu}_e + p \rightarrow e^+ + n$, with a threshold of 1.806 MeV. The deposited $\bar{\nu}_e$ energy results in a prompt signal induced by the positron, and includes annihilation photons. The visible energy is related to the $\bar{\nu}_e$ energy as $E_{vis} = E_{\bar{\nu}_e} - 0.784$ MeV. A delayed signal induced by the neutron capture on protons produces the 2.22 MeV gamma-ray, providing a delayed coincidence signal with a mean capture time of $259.7 \pm 1.3(\text{stat}) \pm 2.0(\text{syst}) \mu\text{s}$ [7]. In Borexino 278 tons of ultra-pure organic liquid scintillator (pseudocumene doped with 1.5 g/l of diphenyloxazole) is confined within a thin, spherical nylon inner vessel (IV) with a nominal radius of 4.25 m and concentrically placed inside a stainless steel sphere (SSS) with radius of 13.7 m and equipped with 2212 8" photomultipliers (PMTs). The volume between the SSS and the IV, divided in two regions by a second outer nylon vessel (OV) is filled with 890 tons of pseudocumene with 3 g/l of light quencher dimethylphthalate to shield the core of the detector against γ radiation and radon from the PMTs and the SSS. The SSS is immersed in an water tank (WT) instrumented with 208 PMTs as a Čerenkov active muon veto, also acting as a shield against γ -rays and neutrons from the surrounding rock.

In Borexino, known sources of $\bar{\nu}_e$ events are nuclear reactors and geo-neutrinos. For geo-neutrinos, ^{238}U and ^{232}Th are the only isotopes abundant enough to significantly contribute events in Borexino above the inverse β decay threshold. The light yield in Borexino is measured to be about 500 photoelectrons (p.e.) / MeV and the energy resolution scales as $\sim 5\%/\sqrt{E}$ [8]. The Borexino liquid scintillator shows a high pulse shape discrimination efficiency in separating high ionizing particles (protons, α 's) from electrons and gamma-rays [9].

The data reported here were collected between December 15, 2007 and March 8, 2015 for a total of 2055.9 days before any selection cut. The number of PMTs for the present data set has been declining with time, from 1931 to 1525 (average 1730), with small run by run fluctuations. As reported in [3] the event energy is a calibrated non-linear function of the number of detected p.e. We perform the analysis directly in number of p.e. We discard events occurring within 2 ms of every muon crossing the outer detector and within 2 s of muons crossing the inner detector to reject neutrons and long-lived cosmogenic radioactivity, respectively. This cut reduces the live-time to 1841.9 days. We apply the following additional selection cuts: (1) prompt scintillation light: $Q_p > 408$ p.e. (i.e. 1.022 MeV corrected for the energy resolution); (2) delayed signal scintillation light: $860 < Q_d < 1300$ p.e. (neutron capture peak); (3) correlation distance between prompt and delayed signals: $\Delta R < 1\text{m}$; (4) correlated time between prompt and delayed signals: $20 < \Delta t < 1280 \mu\text{s}$; (5) pulse shape discrimination with

Gatti filter [10]: $g_{\alpha\beta} < 0.015$ for delayed signals; (6) multiplicity cut: selected event are neither preceded or followed by neutron-like events within a 2 ms window; (7) dynamical fiducial volume [9]: every prompt signal has a reconstructed vertex >30 cm away from the time-varying IV surface; (8) FADC cut: independent check of candidate events features by a 400 MHz digitizer acquisition system. The combined efficiency of the cuts is determined by Monte Carlo to be $(84.2 \pm 1.5)\%$. The total efficiency-corrected exposure for the present data set is 907 ± 44 ton \times yr. We have identified 77 $\bar{\nu}_e$ candidates passing all the selection cuts.

TABLE I. Estimated backgrounds for $\bar{\nu}_e$ given in number of events. Upper limits are given for 90% C.L.

$^9\text{Li}-^8\text{He}$	$0.194^{+0.125}_{-0.089}$
Accidental coincidences	0.221 ± 0.004
Time correlated	$0.035^{+0.029}_{-0.028}$
(α, n) in scintillator	0.165 ± 0.010
(α, n) in buffer	< 0.51
Fast n's (μ in WT)	< 0.01
Fast n's (μ in rock)	< 0.43
untagged muons	0.12 ± 0.01
Fission in PMTs	0.032 ± 0.003
$^{214}\text{Bi}-^{214}\text{Po}$	0.009 ± 0.013
Total	$0.78^{+0.13}_{-0.10}$
	$< 0.65(\text{combined})$

The probability of $\bar{\nu}_e$ -mimicking background events leaking into the dataset was evaluated as follows: (1) The rate of accidental coincidences has been searched for by shifting the delayed time window to 2-20 seconds and keeping all other cuts unchanged. The energy spectrum of these events is limited to <3 MeV. (2) Time correlated events have been searched for in the (2 ms, 2 s) time window. A negligible amount of correlated events with a ~ 1 s time constant were identified and their contribution in the $\bar{\nu}_e$ time window determined. (3) (α, n) background for $\bar{\nu}_e$ search has been extensively discussed elsewhere [3–6]. The average rate of ^{210}Po in the dataset is determined to be (14.1 ± 0.2) counts/(day \cdot ton). (4) Cosmogenic radioactive isotopes which decay via $\beta+n$, namely $^9\text{Li}-^8\text{He}$ [7], have been studied in the (2 ms, 2 s) time window after a muon crossing the inner detector. These events have a wide energy spectrum with maximum at ~ 5 MeV. (5) ^{222}Rn in the liquid scintillator can produce background events through time-correlated $\beta+(\alpha+\gamma)$ decays of ^{214}Bi and ^{214}Po . This decay sequence has a time constant close to the neutron capture time following a $\bar{\nu}_e$ interaction. This background has been estimated by individually tagging $^{214}\text{Bi}-^{214}\text{Po}$ decays. (6) Backgrounds from fast neutrons, untagged muons and spontaneous fission decays in the PMTs are the same as in previous papers [3, 4]. Table I summarizes the estimated backgrounds for $\bar{\nu}_e$ candidates, expressed in number of events. The combined upper limit is obtained by Monte Carlo. The

signal-to-background ratio is ~ 100 .

We have performed an un-binned likelihood fit of the energy spectrum of selected prompt $\bar{\nu}_e$ candidate events [3], shown in Fig. 1. The reactor and geo-neutrinos spectra are obtained by Monte Carlo and the backgrounds considered in this analysis are reported in Table I. The Monte Carlo spectra have been determined as reported in [4]. The reactor neutrinos signal has been calculated adopting the data from IAEA [11] updated to 2014 and the method described in [12]. For the first quarter of 2015 we have used the values from 2014. For the present exposure we predict (87 ± 4) TNU events from nuclear reactors, where 1 Terrestrial Neutrino Unit (TNU) = 1 event / year / 10^{32} protons. The log-likelihood function has two signal components, S_{geo} and S_{react} , left free, and three background components, S_{LiHe} , $S_{\alpha n}$, S_{acc} , constrained to the values and errors reported in Tab. I. These components account for 75% of the total background. The other components were left out due to the uncertainty in their energy spectrum. Combined, they contribute $\sim 1\%$ to the best fit and their contribution to the systematic uncertainty is absorbed in the uncertainty on the energy scale.

Using the value ratio for the masses of Th and U, $m(Th)/m(U) = 3.9$, suggested by the chondritic model, our best fit yields $S_{geo} = 23.7_{-5.7}^{+6.5}(stat)_{-0.6}^{+0.9}(sys)$ events $(43.5_{-10.4}^{+11.8}(stat)_{-2.4}^{+2.7}(sys)$ TNU) and $S_{react} = 52.7_{-7.7}^{+8.5}(stat)_{-0.9}^{+0.7}(sys)$ events $(96.6_{-14.2}^{+15.6}(stat)_{-5.0}^{+4.9}(sys)$ TNU). When expressing the results in TNU, systematic uncertainties from both the exposure (4.8%) and the Monte Carlo energy calibration (1%) are included. Only the Monte Carlo calibration uncertainty is relevant when using the number of decays.

In Fig. 2 we show the 1, 3 and 5σ contours from the log-likelihood fit. Borexino alone observes geo-neutrinos with 5.9σ significance (Fig. 2). The null hypothesis for geo-neutrino observation has a probability equal to 3.6×10^{-9} . The measured geo-neutrino signal corresponds to $\bar{\nu}_e$ fluxes at the detector from decays in the U and Th chains of $\phi(U) = (2.7 \pm 0.7) \times 10^6 \text{ cm}^{-2}\text{s}^{-1}$ and $\phi(Th) = (2.3 \pm 0.6) \times 10^6 \text{ cm}^{-2}\text{s}^{-1}$, respectively. Statistical and systematic uncertainties are added in quadrature.

Fig. 3 shows the probability contours obtained by performing the fit leaving the U and Th spectral contributions as free parameters. The U and Th best-fit contributions are shown in Fig. 1. This measurement shows how Borexino, with larger exposure, could separate the contributions from U and Th, and demonstrates the ability of this detection technique to perform real-time spectroscopy of geo-neutrinos.

The radiogenic heat production for U and Th, $H(U + Th)$, from the present best-fit result is restricted in the range 23-36 TW (see Fig. 4). The range of values includes the uncertainty on the distribution of heat producing elements inside the Earth. The model-independent




FIG. 1. Prompt light yield spectrum, in units of photoelectrons (p.e.), of $\bar{\nu}_e$ candidates and the best-fit. The best-fit shows the geo-neutrino and reactor neutrino spectra (dotted lines) assuming the chondritic ratio. Colored areas show the result of a separate fit with U (blue) and Th (light blue) set as free and independent parameters.

analysis yields a radiogenic heat interval 11-52 TW (69% C.L.) for $H(U + Th)$. Adopting the chondritic mass ratio above and a potassium-to-uranium mass ratio $m(K)/m(U) = 10^4$, the total measured terrestrial radiogenic power is $P(U + Th + K) = 33_{-20}^{+28}$ TW, to be compared with the global terrestrial power output $P_{tot} = 47 \pm 2$ TW [14].

The contribution to the total geo-neutrino signal from the local crust (LOC) is estimated to be $S_{geo}(LOC) = (9.7 \pm 1.3)$ TNU [15]. Considering the contribution from the rest of the crust (ROC) [16], the signal from the crust in Borexino is calculated as $S_{geo}(LOC+ROC) = (23.4 \pm 2.8)$ TNU. In order to estimate the significance of a positive signal from the mantle we have determined the likelihood of $S_{geo}(Mantle) = S_{geo} - S_{geo}(LOC+ROC)$ using the experimental likelihood profile of S_{geo} and a gaussian approximation for the crust contribution. The non-physical region, $S_{geo}(Mantle) < 0$, is excluded. This approach gives a signal from the mantle equal to $S_{geo}(Mantle) = 20.9_{-10.3}^{+15.1}$ TNU, with the null hypothesis rejected at 98% C.L..

An updated measurement of $\bar{\nu}_e$'s with Borexino is presented. We show that Borexino-only data measure geo-neutrinos with 5.9σ significance. We also shows that the background level in Borexino allows to perform a real time spectroscopy of geo-neutrinos, currently limited only by the size of the detector.

The Borexino program is made possible by funding from INFN (Italy), NSF (USA), BMBF, DFG, and MPG (Germany), RFBR: Grants 14-22-03031 and 13-02-12140, RFBR-ASPERA-13-02-92440 (Russia), and NCN Poland (UMO-2012/06/M/ST2/00426). We acknowledge the financial support from the UnivEarthS Labex program of Sorbonne Paris Cit (ANR-10-LABX-0023 and ANR-11-IDEX-0005-02). We acknowledge the generous support and hospitality of the Laboratori Nazionali del Gran




FIG. 2. Best-fit contours for 1, 3 and 5σ for the statistics reported in this paper.




FIG. 3. Best-fit contours for 1, 2 and 3σ for the statistics reported in this paper and for an unbinned likelihood fit with U and Th kept as distinct and free parameters. All other parameters in the fit were kept unchanged. Dashed line corresponds to the chondritic assumption.

Sasso (LNGS). We acknowledge the collaboration of J. Mandula for providing detailed data on reactor load factors.

[1] G. Fiorentini, M. Lissia and F. Mantovani, Phys.Rept.453:117-172,2007.

- [2] G. Bellini et al., Progress in Particle and Nuclear Physics 73 (2013) 1-34.
 [3] G. Bellini et al. (Borexino Collaboration), Phys.Lett. B687 (2010) 299-304.
 [4] G. Bellini et al. (Borexino Collaboration), Phys.Lett. B722 (2013) 295-300.




FIG. 4. The expected geo-neutrino signal in Borexino from U and Th as a function of radiogenic heat released in radioactive decays of U and Th [2]. The three filled regions delimit, from the left to the right, the cosmochemical, geochemical and geodynamical BSE models [13]. Best values from Borexino together with $\pm 1\sigma$ errors are reported: the experimental statistical and systematic uncertainties have been added in quadrature.

- [5] T. Araki et al. (KamLAND Collaboration), Nature 436 (2005) 499.
 [6] S. Abe et al. (KamLAND Collaboration), Phys. Rev. Lett. 100 (2008) 221803
 [7] G. Alimonti et al. (Borexino Collaboration), JCAP08 (2013) 049.
 [8] G. Alimonti et al. (Borexino Collaboration), Nucl. Instr. and Meth. A **600**, 58 (2009).
 [9] G. Bellini et al. (Borexino Collaboration), Phys.Rev. D89 (2014) 11, 112007.
 [10] E. Gatti, F. De Martini, A New Linear Method of Discrimination between Elementary Particles in Scintillation Counters, Nuclear Electronics, 265-276, Vol. 2, 1962
 [11] Nuclear Power Engineering Section, IAEA-PRIS database, <http://www.iaea.org/pris/>, Power Reactor Information System (IAEA-PRIS database).
 [12] M. Baldoncini et al., Physical Review D 91, 065002 (2015).
 [13] O. Sramek et al., Earth and Planetary Science Letters 361 (2013) 356-366.
 [14] J.H. Davies and D.R. Davies, Solid Earth 1 (2010) 5-24.
 [15] M. Coltorti et al., Earth Planet. Sci. Lett. 293 (2010) 259.
 [16] Y. Huang et al. Geochemistry, Geophysics, Geosystems 14, 2003 (2013).