



## Evidence for Exotic Hadron Contributions to $\Lambda_b^0 \rightarrow J/\psi p \pi^-$ Decays

R. Aaij *et al.*\*

(LHCb Collaboration)

(Received 22 June 2016; published 18 August 2016; corrected 24 August 2016)

A full amplitude analysis of  $\Lambda_b^0 \rightarrow J/\psi p \pi^-$  decays is performed with a data sample acquired with the LHCb detector from 7 and 8 TeV  $pp$  collisions, corresponding to an integrated luminosity of  $3 \text{ fb}^{-1}$ . A significantly better description of the data is achieved when, in addition to the previously observed nucleon excitations  $N \rightarrow p \pi^-$ , either the  $P_c(4380)^+$  and  $P_c(4450)^+ \rightarrow J/\psi p$  states, previously observed in  $\Lambda_b^0 \rightarrow J/\psi p K^-$  decays, or the  $Z_c(4200)^- \rightarrow J/\psi \pi^-$  state, previously reported in  $B^0 \rightarrow J/\psi K^+ \pi^-$  decays, or all three, are included in the amplitude models. The data support a model containing all three exotic states, with a significance of more than three standard deviations. Within uncertainties, the data are consistent with the  $P_c(4380)^+$  and  $P_c(4450)^+$  production rates expected from their previous observation taking account of Cabibbo suppression.

DOI: 10.1103/PhysRevLett.117.082003

From the birth of the quark model, it has been anticipated that baryons could be constructed not only from three quarks, but also four quarks and an antiquark [1,2], hereafter referred to as pentaquarks [3,4]. The distribution of the  $J/\psi p$  mass ( $m_{J/\psi p}$ ) in  $\Lambda_b^0 \rightarrow J/\psi p K^-$ ,  $J/\psi \rightarrow \mu^+ \mu^-$  decays (charge conjugation is implied throughout the text) observed with the LHCb detector at the LHC shows a narrow peak suggestive of  $uudc\bar{c}$  pentaquark formation, amidst the dominant formation of various excitations of the  $\Lambda$  [ $uds$ ] baryon ( $\Lambda^*$ ) decaying to  $K^- p$  [5,6]. It was demonstrated that these data cannot be described with  $K^- p$  contributions alone without a specific model of them [7]. Amplitude model fits were also performed on all relevant masses and decay angles of the six-dimensional data [5], using the helicity formalism and Breit-Wigner amplitudes to describe all resonances. In addition to the previously well-established  $\Lambda^*$  resonances, two pentaquark resonances, named the  $P_c(4380)^+$  ( $9\sigma$  significance) and  $P_c(4450)^+$  ( $12\sigma$ ), are required in the model for a good description of the data [5]. The mass, width, and fractional yields (fit fractions) were determined to be  $4380 \pm 8 \pm 29 \text{ MeV}$ ,  $205 \pm 18 \pm 86 \text{ MeV}$ ,  $(8.4 \pm 0.7 \pm 4.3)\%$ , and  $4450 \pm 2 \pm 3 \text{ MeV}$ ,  $39 \pm 5 \pm 19 \text{ MeV}$ ,  $(4.1 \pm 0.5 \pm 1.1)\%$ , respectively. Observations of the same two  $P_c^+$  states in another decay would strengthen their interpretation as genuine exotic baryonic states, rather than kinematical effects related to the so-called triangle singularity [8], as pointed out in Ref. [9].

In this Letter,  $\Lambda_b^0 \rightarrow J/\psi p \pi^-$  decays are analyzed, which are related to  $\Lambda_b^0 \rightarrow J/\psi p K^-$  decays via Cabibbo suppression. LHCb has measured the relative branching fraction  $\mathcal{B}(\Lambda_b^0 \rightarrow J/\psi p \pi^-)/\mathcal{B}(\Lambda_b^0 \rightarrow J/\psi p K^-) = 0.0824 \pm 0.0024 \pm 0.0042$  [10] with the same data sample as used here, corresponding to  $3 \text{ fb}^{-1}$  of integrated luminosity acquired by the LHCb experiment in  $pp$  collisions at 7 and 8 TeV center-of-mass energy. The LHCb detector is a single-arm forward spectrometer covering the pseudorapidity range  $2 < \eta < 5$ , described in detail in Refs. [11,12]. The data selection is similar to that described in Ref. [5], with the  $K^-$  replaced by a  $\pi^-$  candidate. In the preselection a larger significance for the  $\Lambda_b^0$  flight distance and a tighter alignment between the  $\Lambda_b^0$  momentum and the vector from the primary to the secondary vertex are required. To remove specific  $\bar{B}^0$  and  $\bar{B}_s^0$  backgrounds, candidates are vetoed within a  $3\sigma$  invariant mass window around the corresponding nominal  $B$  mass [13] when interpreted as  $\bar{B}^0 \rightarrow J/\psi \pi^+ K^-$  or as  $\bar{B}_s^0 \rightarrow J/\psi K^+ K^-$ . In addition, residual long-lived  $\Lambda \rightarrow p \pi^-$  background is excluded if the  $p \pi^-$  invariant mass ( $m_{p\pi}$ ) lies within  $\pm 5 \text{ MeV}$  of the known  $\Lambda$  mass [13]. The resulting invariant mass spectrum of  $\Lambda_b^0$  candidates is shown in Fig. 1. The signal yield is  $1885 \pm 50$ , determined by an unbinned extended maximum likelihood fit to the mass spectrum. The signal is described by a double-sided crystal ball function [14]. The combinatorial background is modeled by an exponential function. The background of  $\Lambda_b^0 \rightarrow J/\psi p K^-$  events is described by a histogram obtained from simulation, with yield free to vary. This fit is used to assign weights to the candidates using the  $s$ Plot technique [15], which allows the signal component to be projected out by weighting each event depending on the  $J/\psi p \pi^-$  mass. Amplitude fits are performed by minimizing a six-dimensional unbinned negative log likelihood,  $-2 \ln \mathcal{L}$ , with the background subtracted using these

\*Full author list given at the end of the article.

Published by the American Physical Society under the terms of the Creative Commons Attribution 3.0 License. Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI.




FIG. 1. Invariant mass spectrum for the selected  $\Lambda_b^0 \rightarrow J/\psi p \pi^-$  candidates.

weights and the efficiency folded into the signal probability density function, as discussed in detail in Ref. [5].

Amplitude models for the  $\Lambda_b^0 \rightarrow J/\psi p \pi^-$  decays are constructed to examine the possibility of exotic hadron contributions from the  $P_c(4380)^+$  and  $P_c(4450)^+ \rightarrow J/\psi p$  states and from the  $Z_c(4200)^- \rightarrow J/\psi \pi^-$  state, previously reported by the Belle Collaboration in  $B^0 \rightarrow J/\psi K^+ \pi^-$  decays [16] (spin parity  $J^P = 1^+$ , mass and width of  $4196^{+31}_{-29} +^{17}_{-13}$  MeV and  $370 \pm 70^{+70}_{-132}$  MeV, respectively). By analogy with kaon decays [17],  $p \pi^-$  contributions from conventional nucleon excitations (denoted as  $N^*$ ) produced with  $\Delta I = 1/2$  in  $\Lambda_b^0$  decays are expected to dominate over  $\Delta$  excitations with  $\Delta I = 3/2$ , where  $I$  is isospin. The decay matrix elements for the two interfering decay chains,  $\Lambda_b^0 \rightarrow J/\psi N^*$ ,  $N^* \rightarrow p \pi^-$  and  $\Lambda_b^0 \rightarrow P_c^+ \pi^-$ ,  $P_c^+ \rightarrow J/\psi p$  with  $J/\psi \rightarrow \mu^+ \mu^-$  in both cases, are identical to those used in the  $\Lambda_b^0 \rightarrow J/\psi p K^-$  analysis [5], with  $K^-$  and  $\Lambda^*$  replaced by  $\pi^-$  and  $N^*$ . The additional decay chain,  $\Lambda_b^0 \rightarrow Z_c^- p$ ,  $Z_c^- \rightarrow J/\psi \pi^-$ , is also included. Helicity couplings, describing the dynamics of the decays, are expressed in terms of  $LS$  couplings [5], where  $L$  is the decay orbital angular momentum, and  $S$  is the sum of spins of the decay products. This is a convenient way to incorporate parity conservation in strong decays and to allow for reduction of the number of free parameters by excluding high  $L$  values for phase-space suppressed decays.

Table I lists the  $N^*$  resonances considered in the amplitude model of  $p \pi^-$  contributions. There are 15 well-established  $N^*$  resonances [13]. The high-mass and high-spin states ( $9/2$  and  $11/2$ ) are not included, since they require  $L \geq 3$  in the  $\Lambda_b^0$  decay and therefore are unlikely to be produced near the upper kinematic limit of  $m_{p \pi^-}$ . Theoretical models of baryon resonances predict many more high-mass states [18], which have not yet been observed. Their absence could arise from decreased couplings of the higher  $N^*$  excitations to the simple production and decay channels [19] and possibly also from experimental difficulties in identifying broad resonances

TABLE I. The  $N^*$  resonances used in the different fits. Parameters are taken from the PDG [13]. The number of  $LS$  couplings is listed in the columns to the right for the two versions (RM and EM) of the  $N^*$  model discussed in the text. To fix overall phase and magnitude conventions, the  $N(1535)$  complex coupling of lowest  $LS$  is set to  $(1, 0)$ .

State	$J^P$	Mass (MeV)	Width (MeV)	RM	EM
$NR$ $p \pi$	$1/2^-$	...	...	4	4
$N(1440)$	$1/2^+$	1430	350	3	4
$N(1520)$	$3/2^-$	1515	115	3	3
$N(1535)$	$1/2^-$	1535	150	4	4
$N(1650)$	$1/2^-$	1655	140	1	4
$N(1675)$	$5/2^-$	1675	150	3	5
$N(1680)$	$5/2^+$	1685	130	...	3
$N(1700)$	$3/2^-$	1700	150	...	3
$N(1710)$	$1/2^+$	1710	100	...	4
$N(1720)$	$3/2^+$	1720	250	3	5
$N(1875)$	$3/2^-$	1875	250	...	3
$N(1900)$	$3/2^+$	1900	200	...	3
$N(2190)$	$7/2^-$	2190	500	...	3
$N(2300)$	$1/2^+$	2300	340	...	3
$N(2570)$	$5/2^-$	2570	250	...	3
Free parameters				40	106

and insufficient statistics at high masses in scattering experiments. The possibility of high-mass, low-spin  $N^*$  states is explored by including two very significant, but unconfirmed, resonances claimed by the BESIII Collaboration in  $\psi(2S) \rightarrow p \bar{p} \pi^0$  decays [20]:  $1/2^+$   $N(2300)$  and  $5/2^-$   $N(2570)$ . A nonresonant  $J^P = 1/2^-$   $p \pi^-$   $S$ -wave component is also included. Two models, labeled “reduced” (RM) and “extended” (EM), are considered and differ in the number of resonances and of  $LS$  couplings included in the fit as listed in Table I. The reduced model, used for the central values of fit fractions, includes only the resonances and  $L$  couplings that give individually significant contributions. The systematic uncertainties and the significances for the exotic states are evaluated with the extended model by including all well-motivated resonances and the maximal number of  $LS$  couplings for which the fit is able to converge.

All  $N^*$  resonances are described by Breit-Wigner functions [5] to model their line shape and phase variation as a function of  $m_{p \pi^-}$ , except for the  $N(1535)$ , which is described by a Flatté function [21] to account for the threshold of the  $n \eta$  channel. The mass and width are fixed to the values determined from previous experiments [13]. The couplings to the  $n \eta$  and  $p \pi^-$  channels for the  $N(1535)$  state are determined by the branching fractions of the two channels [22]. The nonresonant  $S$ -wave component is described with a function that depends inversely on  $m_{p \pi^-}^2$ , as this is found to be preferred by the data. An alternative description of the  $1/2^-$   $p \pi^-$  contributions, including the  $N(1535)$  and nonresonant components, is provided by a  $K$ -matrix model obtained from multichannel partial wave




FIG. 2. Background-subtracted data and fit projections onto  $m_{p\pi}$ . Fits are shown with models containing  $N^*$  states only (EM) and with  $N^*$  states (RM) plus exotic contributions.

analysis by the Bonn-Gatchina group [22,23] and is used to estimate systematic uncertainties.

The limited number of signal events and the large number of free parameters in the amplitude fits prevent an open-ended analysis of  $J/\psi p$  and  $J/\psi \pi^-$  contributions. Therefore, the data are examined only for the presence of the previously observed  $P_c(4380)^+$ ,  $P_c(4450)^+$  states [5] and the claimed  $Z_c(4200)^-$  resonance [16]. In the fits, the mass and width of each exotic state are fixed to the reported central values. The  $LS$  couplings describing  $P_c^+ \rightarrow J/\psi p$  decays are also fixed to the values obtained from the Cabibbo-favored channel. This leaves four free parameters per  $P_c^+$  state for the  $\Lambda_b^0 \rightarrow P_c^+ \pi^-$  couplings. The nominal fits are performed for the most likely  $(3/2^-, 5/2^+)$   $J^P$  assignment to the  $P_c(4380)^+$ ,  $P_c(4450)^+$  states [5]. All couplings for the  $1^+ Z_c(4200)^-$  contribution are allowed to vary (ten free parameters).

The fits show a significant improvement when exotic contributions are included. When all three exotic

contributions are added to the EM  $N^*$ -only model, the  $\Delta(-2 \ln \mathcal{L})$  value is 49.0, which corresponds to their combined statistical significance of  $3.9\sigma$ . Including the systematic uncertainties discussed later lowers their significance to  $3.1\sigma$ . The systematic uncertainties are included in subsequent significance figures. Because of the ambiguity between the  $P_c(4380)^+$ ,  $P_c(4450)^+$  and  $Z_c(4200)^-$  contributions, no single one of them makes a significant difference to the model. Adding either state to a model already containing the other two, or the two  $P_c^+$  states to a model already containing the  $Z_c(4200)^-$  contribution, yields significances below  $1.7\sigma$  [ $0.4\sigma$  for adding the  $Z_c(4200)^-$  after the two  $P_c^+$  states]. If the  $Z_c(4200)^-$  contribution is assumed to be negligible, adding the two  $P_c^+$  states to a model without exotics yields a significance of  $3.3\sigma$ . On the other hand, under the assumption that no  $P_c^+$  states are produced, adding the  $Z_c(4200)^-$  to a model without exotics yields a significance of  $3.2\sigma$ . The significances are determined using Wilks' theorem [24], the applicability of which has been verified by simulation.

A satisfactory description of the data is already reached with the RM  $N^*$  model if either the two  $P_c^+$ , or the  $Z_c^-$ , or all three states, are included in the fit. The projections of the full amplitude fit onto the invariant masses and the decay angles reasonably well reproduce the data, as shown in Figs. 2–5. The EM  $N^*$ -only model does not give good descriptions of the peaking structure in  $m_{J/\psi p}$  observed for  $m_{p\pi} > 1.8$  GeV [Fig. 3(b)]. In fact, all contributions to  $\Delta(-2 \ln \mathcal{L})$  favoring the exotic components belong to this  $m_{p\pi}$  region. The models with the  $P_c^+$  states describe the  $m_{J/\psi p}$  peaking structure better than with the  $Z_c(4200)^-$  alone (see Supplemental Material [25]).

The model with all three exotic resonances is used when determining the fit fractions. The sources of systematic uncertainty are listed in Table II. They include varying the masses and widths of  $N^*$  resonances, varying the masses and widths of the exotic states, considering  $N^*$  model




FIG. 3. Background-subtracted data and fit projections onto  $m_{J/\psi p}$  for (a) all events and (b) the  $m_{p\pi} > 1.8$  GeV region. See the legend and caption of Fig. 2 for a description of the components.




FIG. 4. Background-subtracted data and fit projections onto  $m_{J/\psi\pi}$  for (a) all events and (b) the  $m_{p\pi} > 1.8$  GeV region. See the legend and caption of Fig. 2 for a description of the components.

dependence and other possible spin parities  $J^P$  for the two  $P_c^+$  states, varying the Blatt-Weisskopf radius [5] between 1.5 and 4.5  $\text{GeV}^{-1}$ , changing the angular momenta  $L$  in  $\Lambda_b^0$  decays that are used in the resonant mass description by one or two units, using the  $K$ -matrix model for the  $S$ -wave  $p\pi$  resonances, varying the fixed couplings of the  $P_c^+$  decay by their uncertainties, and splitting  $\Lambda_b^0$  and  $J/\psi$  helicity angles into bins when determining the weights for the background subtraction to account for correlations between the invariant mass of  $J/\psi p\pi^-$  and these angles. A putative

$Z_c(4430)^-$  contribution [16,26,27] hardly improves the value of  $-2 \ln \mathcal{L}$  relative to the EM  $N^*$ -only model, and thus is considered among systematic uncertainties. Exclusion of the  $Z_c(4200)^-$  state from the fit model is also considered to determine the systematic uncertainties for the two  $P_c^+$  states.

The EM model is used to assess the uncertainty due to the  $N^*$  modeling when computing significances. The RM model gives larger significances. All sources of systematic uncertainties, including the ambiguities in the quantum number assignments to the two  $P_c^+$  states, are accounted for in the calculation of the significance of various contributions, by using the smallest  $\Delta(-2 \ln \mathcal{L})$  among the fits representing different systematic variations.

The fit fractions for the  $P_c(4380)^+$ ,  $P_c(4450)^+$  and  $Z_c(4200)^-$  states are measured to be  $(5.1 \pm 1.5^{+2.6}_{-1.6})\%$ ,




FIG. 5. Background-subtracted data and fit projections of decay angles describing the  $N^*$  decay chain, which are included in the amplitude fit. The helicity angle of particle  $P$ ,  $\theta_P$ , is the polar angle in the rest frame of  $P$  between a decay product of  $P$  and the boost direction from the particle decaying to  $P$ . The azimuthal angle between decay planes of  $\Lambda_b^0$  and  $N^*$  (of  $J/\psi$ ) is denoted as  $\phi_\pi$  ( $\phi_\mu$ ). See Ref. [5] for more details.

TABLE II. Summary of absolute systematic uncertainties of the fit fractions in units of percent.

Source	$P_c(4450)^+$	$P_c(4380)^+$	$Z_c(4200)^-$
$N^*$ masses and widths	$\pm 0.05$	$\pm 0.23$	$\pm 0.31$
$P_c^+$ , $Z_c^-$ masses and widths	$\pm 0.32$	$\pm 1.27$	$\pm 1.56$
Additional $N^*$	+0.08 -0.23	+0.59 -0.55	+0.71 -2.92
Inclusion of $Z_c(4430)^-$	+0.01	+0.97	+2.87
Exclusion of $Z_c(4200)^-$	-0.15	+1.61	...
Other $J^P$	+0.38 -0.00	+0.92 -0.28	+0.00 -2.16
Blatt-Weisskopf radius	$\pm 0.11$	$\pm 0.17$	$\pm 0.21$
$L_{\Lambda_b^0}^{N^*}$ in $\Lambda_b^0 \rightarrow J/\psi N^*$	$\pm 0.07$	$\pm 0.46$	$\pm 0.04$
$L_{\Lambda_b^0}^{P_c}$ in $\Lambda_b^0 \rightarrow P_c^+ \pi^-$	-0.05	-0.17	+0.09
$L_{\Lambda_b^0}^{Z_c}$ in $\Lambda_b^0 \rightarrow Z_c^- p$	$\pm 0.07$	$\pm 0.22$	$\pm 0.53$
$K$ -matrix model	-0.03	+0.11	-0.02
$P_c^+$ couplings	$\pm 0.14$	$\pm 0.31$	$\pm 0.36$
Background subtraction	-0.07	-0.13	-0.39
Total	+0.55 -0.48	+2.61 -1.58	+3.43 -4.04

$(1.6_{-0.6}^{+0.8+0.6})\%$ , and  $(7.7 \pm 2.8_{-4.0}^{+3.4})\%$  respectively, and to be less than 8.9%, 2.9%, and 13.3% at 90% confidence level, respectively. When the two  $P_c^+$  states are not considered, the fraction for the  $Z_c(4200)^-$  state is surprisingly large,  $(17.2 \pm 3.5)\%$ , where the uncertainty is statistical only, given that its fit fraction was measured to be only  $(1.9_{-0.5}^{+0.7+0.9})\%$  in  $B^0 \rightarrow J/\psi K^+ \pi^-$  decays [16]. Conversely, the fit fractions of the two  $P_c^+$  states remain stable regardless of the inclusion of the  $Z_c(4200)^-$  state. We measure the relative branching fraction  $R_{\pi/K} \equiv \mathcal{B}(\Lambda_b^0 \rightarrow \pi^- P_c^+)/\mathcal{B}(\Lambda_b^0 \rightarrow K^- P_c^+)$  to be  $0.050 \pm 0.016_{-0.016}^{+0.026} \pm 0.025$  for  $P_c(4380)^+$  and  $0.033_{-0.014}^{+0.016+0.011} \pm 0.009$  for  $P_c(4450)^+$ , respectively, where the first error is statistical, the second is systematic, and the third is due to the systematic uncertainty on the fit fractions of the  $P_c^+$  states in  $J/\psi p K^-$  decays. The results are consistent with a prediction of  $(0.07-0.08)$  [28], where the assumption is made that an additional diagram with internal  $W$  emission, which can only contribute to the Cabibbo-suppressed mode, is negligible. Our measurement rules out the proposal that the  $P_c^+$  state in the  $\Lambda_b^0 \rightarrow J/\psi p K^-$  decay is produced mainly by the charmless  $\Lambda_b^0$  decay via the  $b \rightarrow u\bar{u}s$  transition, since this predicts a very large value for  $R_{\pi/K} = 0.58 \pm 0.05$  [29].

In conclusion, we have performed a full amplitude fit to  $\Lambda_b^0 \rightarrow J/\psi p \pi^-$  decays allowing for previously observed conventional ( $p\pi^-$ ) and exotic ( $J/\psi p$  and  $J/\psi \pi^-$ ) resonances. A significantly better description of the data is achieved by either including the two  $P_c^+$  states observed in  $\Lambda_b^0 \rightarrow J/\psi p K^-$  decays [5], or the  $Z_c(4200)^-$  state reported by the Belle Collaboration in  $B^0 \rightarrow J/\psi \pi^- K^+$  decays [16]. If both types of exotic resonances are included, the total significance for them is  $3.1\sigma$ . Individual exotic hadron components, or the two  $P_c^+$  states taken together, are not significant as long as the other(s) is (are) present. Within the statistical and systematic errors, the data are consistent with the  $P_c(4380)^+$  and  $P_c(4450)^+$  production rates expected from their previous observation and Cabibbo suppression. Assuming that the  $Z_c(4200)^-$  contribution is negligible, there is a  $3.3\sigma$  significance for the two  $P_c^+$  states taken together.

We thank the Bonn-Gatchina group who provided us with the  $K$ -matrix  $p\pi^-$  model. We express our gratitude to our colleagues in the CERN accelerator departments for the excellent performance of the LHC. We thank the technical and administrative staff at the LHCb institutes. We acknowledge support from CERN and from the following national agencies: CAPES, CNPq, FAPERJ, and FINEP (Brazil); NSFC (China); CNRS/IN2P3 (France); BMBF, DFG, and MPG (Germany); INFN (Italy); FOM and NWO (Netherlands); MNiSW and NCN (Poland); MEN/IFA (Romania); MinES and FANO (Russia); MinECo (Spain); SNSF and SER (Switzerland); NASU (Ukraine);

STFC (United Kingdom); and NSF (USA). We acknowledge the computing resources that are provided by CERN, IN2P3 (France), KIT and DESY (Germany), INFN (Italy), SURF (Netherlands), PIC (Spain), GridPP (United Kingdom), RRCKI and Yandex LLC (Russia), CSCS (Switzerland), IFIN-HH (Romania), CBPF (Brazil), PL-GRID (Poland), and OSC (USA). We are indebted to the communities behind the multiple open source software packages on which we depend. Individual groups or members have received support from AvH Foundation (Germany), EPLANET, Marie Skłodowska-Curie Actions and ERC (European Union), Conseil Général de Haute-Savoie, Labex ENIGMASS, and OCEVU, Région Auvergne (France), RFBR and Yandex LLC (Russia), GVA, XuntaGal, and GENCAT (Spain), Herchel Smith Fund, The Royal Society, Royal Commission for the Exhibition of 1851, and the Leverhulme Trust (United Kingdom).

Deutsches Elektronen-Synchrotron

- 
- [1] M. Gell-Mann, A schematic model of baryons and mesons, *Phys. Lett.* **8**, 214 (1964).
  - [2] G. Zweig, Report No. CERN-TH-401, 1964.
  - [3] L. Montanet, G. C. Rossi, and G. Veneziano, Baryonium physics, *Phys. Rep.* **63**, 151 (1980).
  - [4] H. J. Lipkin, New possibilities for exotic hadrons: anti-charmed strange baryons, *Phys. Lett. B* **195**, 484 (1987).
  - [5] R. Aaij *et al.* (LHCb Collaboration), Observation of  $J/\psi p$  Resonances Consistent with Pentaquark States in  $\Lambda_b^0 \rightarrow J/\psi p K^-$  Decays, *Phys. Rev. Lett.* **115**, 072001 (2015).
  - [6] R. Aaij *et al.* (LHCb Collaboration), Study of the productions of  $\Lambda_b^0$  and  $\bar{B}^0$  hadrons in  $pp$  collisions and first measurement of the  $\Lambda_b^0 \rightarrow J/\psi p K^-$  branching fraction, *Chin. Phys. C* **40**, 011001 (2016).
  - [7] R. Aaij *et al.* (LHCb Collaboration), preceding Letter, Model-Independent Evidence for  $J/\psi p$  Contributions to  $\Lambda_b^0 \rightarrow J/\psi p K^-$  Decays, *Phys. Rev. Lett.* **117**, 082002 (2016).
  - [8] F.-K. Guo, U.-G. Meiner, W. Wang, and Z. Yang, How to reveal the exotic nature of the  $P_c(4450)$ , *Phys. Rev. D* **92**, 071502 (2015); X.-H. Liu, Q. Wang, and Q. Zhao, Understanding the newly observed heavy pentaquark candidates, *Phys. Lett. B* **757**, 231 (2016); M. Mikhasenko, A triangle singularity and the LHCb pentaquarks, [arXiv:1507.06552](https://arxiv.org/abs/1507.06552).
  - [9] T. J. Burns, Phenomenology of  $P_c(4380)^+$ ,  $P_c(4450)^+$ , and related states, *Eur. Phys. J. A* **51**, 152 (2015); E. Wang, H.-X. Chen, L.-S. Geng, D.-M. Li, and E. Oset, Hidden-charm pentaquark state in  $\Lambda_b^0 \rightarrow J/\psi p \pi^-$  decay, *Phys. Rev. D* **93**, 094001 (2016).
  - [10] R. Aaij *et al.* (LHCb Collaboration), Observation of the  $\Lambda_b^0 \rightarrow J/\psi p \pi^-$  decay, *J. High Energy Phys.* **07** (2014) 103.
  - [11] A. A. Alves Jr. *et al.* (LHCb Collaboration), The LHCb detector at the LHC, *J. Instrum.* **3**, S08005 (2008).
  - [12] R. Aaij *et al.* (LHCb Collaboration), LHCb detector performance, *Int. J. Mod. Phys. A* **30**, 1530022 (2015).
  - [13] K. A. Olive *et al.* (Particle Data Group), Review of particle physics, *Chin. Phys. C* **38**, 090001 (2014).

- [14] T. Skwarnicki, Ph.D. thesis, Institute of Nuclear Physics [Report No. DESY-F31-86-02, 1986].
- [15] M. Pivk and F.R. Le Diberder, *sPlot*: a statistical tool to unfold data distributions, *Nucl. Instrum. Methods Phys. Res., Sect. A* **555**, 356 (2005).
- [16] K. Chilikin *et al.* (Belle Collaboration), Observation of a new charged charmoniumlike state in  $\bar{B}^0 \rightarrow J/\psi K^- \pi^+$  decays, *Phys. Rev. D* **90**, 112009 (2014).
- [17] J.F. Donoghue, E. Golowich, W.A. Ponce, and B.R. Holstein, Analysis of  $\Delta S = 1$  nonleptonic weak decays and the  $\Delta I = 1/2$  rule, *Phys. Rev. D* **21**, 186 (1980).
- [18] U. Loring, B. C. Metsch, and H. R. Petry, The light baryon spectrum in a relativistic quark model with instanton induced quark forces: the nonstrange baryon spectrum and ground states, *Eur. Phys. J. A* **10**, 395 (2001).
- [19] R. Koniuk and N. Isgur, Where Have all the Resonances Gone? An Analysis of Baryon Couplings in a Quark Model with Chromodynamics, *Phys. Rev. Lett.* **44**, 845 (1980).
- [20] M. Ablikim *et al.* (BESIII Collaboration), Observation of Two New  $N^*$  Resonances in the Decay  $\psi(3686) \rightarrow p \bar{p} \pi^0$ , *Phys. Rev. Lett.* **110**, 022001 (2013).
- [21] S. M. Flatté, Coupled-channel analysis of the  $\pi\eta$  and  $K\bar{K}$  systems near  $K\bar{K}$  threshold, *Phys. Lett. B* **63B**, 224 (1976).
- [22] A. V. Anisovich, R. Beck, E. Klempt, V. A. Nikonov, A. V. Sarantsev, and U. Thoma, Properties of baryon resonances from a multichannel partial wave analysis, *Eur. Phys. J. A* **48**, 15 (2012).
- [23] A. V. Anisovich, E. Klempt, V. A. Nikonov, M. A. Matveev, A. V. Sarantsev, and U. Thoma, Photoproduction of pions and properties of baryon resonances from a Bonn-Gatchina partial wave analysis, *Eur. Phys. J. A* **44**, 203 (2010).
- [24] S. S. Wilks, The large-sample distribution of the likelihood ratio for testing composite hypotheses, *Ann. Math. Stat.* **9**, 60 (1938).
- [25] See Supplemental Material at <http://link.aps.org/supplemental/10.1103/PhysRevLett.117.082003> for the Dalitz plot distribution and additional fit results.
- [26] K. Chilikin *et al.* (Belle Collaboration), Experimental constraints on the spin and parity of the  $Z(4430)^+$ , *Phys. Rev. D* **88**, 074026 (2013).
- [27] R. Aaij *et al.* (LHCb Collaboration), Observation of the Resonant Character of the  $Z(4430)^-$  State, *Phys. Rev. Lett.* **112**, 222002 (2014); R. Aaij *et al.* (LHCb Collaboration), A model-independent confirmation of the  $Z(4430)^-$  state, *Phys. Rev. D* **92**, 112009 (2015).
- [28] H.-Y. Cheng and C.-K. Chua, Bottom baryon decays to pseudoscalar meson and pentaquark, *Phys. Rev. D* **92**, 096009 (2015).
- [29] Y. K. Hsiao and C. Q. Geng, Pentaquarks from intrinsic charms in  $\Lambda_b^0$  decays, *Phys. Lett. B* **751**, 572 (2015).

R. Aaij,<sup>39</sup> C. Abellán Beteta,<sup>41</sup> B. Adeva,<sup>38</sup> M. Adinolfi,<sup>47</sup> Z. Ajaltouni,<sup>5</sup> S. Akar,<sup>6</sup> J. Albrecht,<sup>10</sup> F. Alessio,<sup>39</sup> M. Alexander,<sup>52</sup> S. Ali,<sup>42</sup> G. Alkhazov,<sup>31</sup> P. Alvarez Cartelle,<sup>54</sup> A. A. Alves Jr.,<sup>58</sup> S. Amato,<sup>2</sup> S. Amerio,<sup>23</sup> Y. Amhis,<sup>7</sup> L. An,<sup>40</sup> L. Anderlini,<sup>18</sup> G. Andreassi,<sup>40</sup> M. Andreotti,<sup>17,a</sup> J. E. Andrews,<sup>59</sup> R. B. Appleby,<sup>55</sup> O. Aquines Gutierrez,<sup>11</sup> F. Archilli,<sup>1</sup> P. d'Argent,<sup>12</sup> J. Arnau Romeu,<sup>6</sup> A. Artamonov,<sup>36</sup> M. Artuso,<sup>60</sup> E. Aslanides,<sup>6</sup> G. Auriemma,<sup>26,b</sup> M. Baalouch,<sup>5</sup> S. Bachmann,<sup>12</sup> J. J. Back,<sup>49</sup> A. Badalov,<sup>37</sup> C. Baesso,<sup>61</sup> W. Baldini,<sup>17</sup> R. J. Barlow,<sup>55</sup> C. Barschel,<sup>39</sup> S. Barsuk,<sup>7</sup> W. Barter,<sup>39</sup> V. Batozskaya,<sup>29</sup> V. Battista,<sup>40</sup> A. Bay,<sup>40</sup> L. Beaucourt,<sup>4</sup> J. Beddow,<sup>52</sup> F. Bedeschi,<sup>24</sup> I. Bediaga,<sup>1</sup> L. J. Bel,<sup>42</sup> V. Bellee,<sup>40</sup> N. Belloli,<sup>21,c</sup> K. Belous,<sup>36</sup> I. Belyaev,<sup>32</sup> E. Ben-Haim,<sup>8</sup> G. Bencivenni,<sup>19</sup> S. Benson,<sup>39</sup> J. Benton,<sup>47</sup> A. Berezhnov,<sup>33</sup> R. Bernet,<sup>41</sup> A. Bertolin,<sup>23</sup> M.-O. Bettler,<sup>39</sup> M. van Beuzekom,<sup>42</sup> S. Bifani,<sup>46</sup> P. Billoir,<sup>8</sup> T. Bird,<sup>55</sup> A. Birnkraut,<sup>10</sup> A. Bitadze,<sup>55</sup> A. Bizzeti,<sup>18,d</sup> T. Blake,<sup>49</sup> F. Blanc,<sup>40</sup> J. Blouw,<sup>11</sup> S. Blusk,<sup>60</sup> V. Bocci,<sup>26</sup> T. Boettcher,<sup>57</sup> A. Bondar,<sup>35</sup> N. Bondar,<sup>31,39</sup> W. Bonivento,<sup>16</sup> S. Borghi,<sup>55</sup> M. Borisyak,<sup>67</sup> M. Borsato,<sup>38</sup> F. Bossu,<sup>7</sup> M. Boubdir,<sup>9</sup> T. J. V. Bowcock,<sup>53</sup> E. Bowen,<sup>41</sup> C. Bozzi,<sup>17,39</sup> S. Braun,<sup>12</sup> M. Britsch,<sup>12</sup> T. Britton,<sup>60</sup> J. Brodzicka,<sup>55</sup> E. Buchanan,<sup>47</sup> C. Burr,<sup>55</sup> A. Bursche,<sup>2</sup> J. Buytaert,<sup>39</sup> S. Cadeddu,<sup>16</sup> R. Calabrese,<sup>17,a</sup> M. Calvi,<sup>21,c</sup> M. Calvo Gomez,<sup>37,e</sup> P. Campana,<sup>19</sup> D. Campora Perez,<sup>39</sup> L. Capriotti,<sup>55</sup> A. Carbone,<sup>15,f</sup> G. Carboni,<sup>25,g</sup> R. Cardinale,<sup>20,h</sup> A. Cardini,<sup>16</sup> P. Carniti,<sup>21,c</sup> L. Carson,<sup>51</sup> K. Carvalho Akiba,<sup>2</sup> G. Casse,<sup>53</sup> L. Cassina,<sup>21,c</sup> L. Castillo Garcia,<sup>40</sup> M. Cattaneo,<sup>39</sup> Ch. Cauet,<sup>10</sup> G. Cavallero,<sup>20</sup> R. Cenci,<sup>24,i</sup> M. Charles,<sup>8</sup> Ph. Charpentier,<sup>39</sup> G. Chatzikonstantinidis,<sup>46</sup> M. Chefdeville,<sup>4</sup> S. Chen,<sup>55</sup> S.-F. Cheung,<sup>56</sup> V. Chobanova,<sup>38</sup> M. Chrzaszcz,<sup>41,27</sup> X. Cid Vidal,<sup>38</sup> G. Ciezarek,<sup>42</sup> P. E. L. Clarke,<sup>51</sup> M. Clemencic,<sup>39</sup> H. V. Cliff,<sup>48</sup> J. Closier,<sup>39</sup> V. Coco,<sup>58</sup> J. Cogan,<sup>6</sup> E. Cogneras,<sup>5</sup> V. Cogoni,<sup>16,j</sup> L. Cojocariu,<sup>30</sup> G. Collazuol,<sup>23,k</sup> P. Collins,<sup>39</sup> A. Comerma-Montells,<sup>12</sup> A. Contu,<sup>39</sup> A. Cook,<sup>47</sup> S. Coquereau,<sup>8</sup> G. Corti,<sup>39</sup> M. Corvo,<sup>17,a</sup> C. M. Costa Sobral,<sup>49</sup> B. Couturier,<sup>39</sup> G. A. Cowan,<sup>51</sup> D. C. Craik,<sup>51</sup> A. Crocombe,<sup>49</sup> M. Cruz Torres,<sup>61</sup> S. Cunliffe,<sup>54</sup> R. Currie,<sup>54</sup> C. D'Ambrosio,<sup>39</sup> E. Dall'Occo,<sup>42</sup> J. Dalseno,<sup>47</sup> P. N. Y. David,<sup>42</sup> A. Davis,<sup>58</sup> O. De Aguiar Francisco,<sup>2</sup> K. De Bruyn,<sup>6</sup> S. De Capua,<sup>55</sup> M. De Cian,<sup>12</sup> J. M. De Miranda,<sup>1</sup> L. De Paula,<sup>2</sup> P. De Simone,<sup>19</sup> C.-T. Dean,<sup>52</sup> D. Decamp,<sup>4</sup> M. Deckenhoff,<sup>10</sup> L. Del Buono,<sup>8</sup> M. Demmer,<sup>10</sup> D. Derkach,<sup>67</sup> O. Deschamps,<sup>5</sup> F. Dettori,<sup>39</sup> B. Dey,<sup>22</sup> A. Di Canto,<sup>39</sup> H. Dijkstra,<sup>39</sup> F. Dordei,<sup>39</sup> M. Dorigo,<sup>40</sup> A. Dosil Suárez,<sup>38</sup> A. Dovbnya,<sup>44</sup> K. Dreimanis,<sup>53</sup> L. Dufour,<sup>42</sup> G. Dujany,<sup>55</sup> K. Dungs,<sup>39</sup> P. Durante,<sup>39</sup> R. Dzhelyadin,<sup>36</sup> A. Dziurda,<sup>39</sup> A. Dzyuba,<sup>31</sup> N. Déleage,<sup>4</sup> S. Easo,<sup>50</sup> U. Egede,<sup>54</sup> V. Egorychev,<sup>32</sup> S. Eidelman,<sup>35</sup> S. Eisenhardt,<sup>51</sup> U. Eitschberger,<sup>10</sup> R. Ekelhof,<sup>10</sup> L. Eklund,<sup>52</sup> Ch. Elsasser,<sup>41</sup> S. Ely,<sup>60</sup> S. Esen,<sup>12</sup> H. M. Evans,<sup>48</sup> T. Evans,<sup>56</sup> A. Falabella,<sup>15</sup> N. Farley,<sup>46</sup>

S. Farry,<sup>53</sup> R. Fay,<sup>53</sup> D. Ferguson,<sup>51</sup> V. Fernandez Albor,<sup>38</sup> F. Ferrari,<sup>15,39</sup> F. Ferreira Rodrigues,<sup>1</sup> M. Ferro-Luzzi,<sup>39</sup> S. Filippov,<sup>34</sup> M. Fiore,<sup>17,a</sup> M. Fiorini,<sup>17,a</sup> M. Firlej,<sup>28</sup> C. Fitzpatrick,<sup>40</sup> T. Fiutowski,<sup>28</sup> F. Fleuret,<sup>7,1</sup> K. Fohl,<sup>39</sup> M. Fontana,<sup>16</sup> F. Fontanelli,<sup>20,h</sup> D. C. Forshaw,<sup>60</sup> R. Forty,<sup>39</sup> M. Frank,<sup>39</sup> C. Frei,<sup>39</sup> M. Frosini,<sup>18</sup> J. Fu,<sup>22,m</sup> E. Furfaro,<sup>25,g</sup> C. Färber,<sup>39</sup> A. Gallas Torreira,<sup>38</sup> D. Galli,<sup>15,f</sup> S. Gallorini,<sup>23</sup> S. Gambetta,<sup>51</sup> M. Gandelman,<sup>2</sup> P. Gandini,<sup>56</sup> Y. Gao,<sup>3</sup> J. García Pardiñas,<sup>38</sup> J. Garra Tico,<sup>48</sup> L. Garrido,<sup>37</sup> P. J. Garsed,<sup>48</sup> D. Gascon,<sup>37</sup> C. Gaspar,<sup>39</sup> L. Gavardi,<sup>10</sup> G. Gazzoni,<sup>5</sup> D. Gerick,<sup>12</sup> E. Gersabeck,<sup>12</sup> M. Gersabeck,<sup>55</sup> T. Gershon,<sup>49</sup> Ph. Ghez,<sup>4</sup> S. Gianì,<sup>40</sup> V. Gibson,<sup>48</sup> O. G. Girard,<sup>40</sup> L. Giubega,<sup>30</sup> K. Gizdov,<sup>51</sup> V. V. Gligorov,<sup>8</sup> D. Golubkov,<sup>32</sup> A. Golutvin,<sup>54,39</sup> A. Gomes,<sup>1,n</sup> I. V. Gorelov,<sup>33</sup> C. Gotti,<sup>21,c</sup> M. Grabalosa Gándara,<sup>5</sup> R. Graciani Diaz,<sup>37</sup> L. A. Granado Cardoso,<sup>39</sup> E. Graugés,<sup>37</sup> E. Graverini,<sup>41</sup> G. Graziani,<sup>18</sup> A. Grecu,<sup>30</sup> P. Griffith,<sup>46</sup> L. Grillo,<sup>12</sup> B. R. Gruber Cazon,<sup>56</sup> O. Grünberg,<sup>65</sup> E. Gushchin,<sup>34</sup> Yu. Guz,<sup>36</sup> T. Gys,<sup>39</sup> C. Göbel,<sup>61</sup> T. Hadavizadeh,<sup>56</sup> C. Hadjivasiliou,<sup>60</sup> G. Haefeli,<sup>40</sup> C. Haen,<sup>39</sup> S. C. Haines,<sup>48</sup> S. Hall,<sup>54</sup> B. Hamilton,<sup>59</sup> X. Han,<sup>12</sup> S. Hansmann-Menzemer,<sup>12</sup> N. Harnew,<sup>56</sup> S. T. Harnew,<sup>47</sup> J. Harrison,<sup>55</sup> J. He,<sup>39</sup> T. Head,<sup>40</sup> A. Heister,<sup>9</sup> K. Hennessy,<sup>53</sup> P. Henrard,<sup>5</sup> L. Henry,<sup>8</sup> J. A. Hernando Morata,<sup>38</sup> E. van Herwijnen,<sup>39</sup> M. Heß,<sup>65</sup> A. Hicheur,<sup>2</sup> D. Hill,<sup>56</sup> C. Hombach,<sup>55</sup> W. Hulsbergen,<sup>42</sup> T. Humair,<sup>54</sup> M. Hushchyn,<sup>67</sup> N. Hussain,<sup>56</sup> D. Hutchcroft,<sup>53</sup> M. Idzik,<sup>28</sup> P. Ilten,<sup>57</sup> R. Jacobsson,<sup>39</sup> A. Jaeger,<sup>12</sup> J. Jalocha,<sup>56</sup> E. Jans,<sup>42</sup> A. Jawahery,<sup>59</sup> M. John,<sup>56</sup> D. Johnson,<sup>39</sup> C. R. Jones,<sup>48</sup> C. Joram,<sup>39</sup> B. Jost,<sup>39</sup> N. Jurik,<sup>60</sup> S. Kandybei,<sup>44</sup> W. Kanso,<sup>6</sup> M. Karacson,<sup>39</sup> J. M. Kariuki,<sup>47</sup> S. Karodia,<sup>52</sup> M. Kecke,<sup>12</sup> M. Kelsey,<sup>60</sup> I. R. Kenyon,<sup>46</sup> M. Kenzie,<sup>39</sup> T. Ketel,<sup>43</sup> E. Khairullin,<sup>67</sup> B. Khanji,<sup>21,39,c</sup> C. Khurewathanakul,<sup>40</sup> T. Kirm,<sup>9</sup> S. Klaver,<sup>55</sup> K. Klimaszewski,<sup>29</sup> S. Koliiev,<sup>45</sup> M. Kolpin,<sup>12</sup> I. Komarov,<sup>40</sup> R. F. Koopman,<sup>43</sup> P. Koppenburg,<sup>42</sup> A. Kozachuk,<sup>33</sup> M. Kozeiha,<sup>5</sup> L. Kravchuk,<sup>34</sup> K. Kreplin,<sup>12</sup> M. Kreps,<sup>49</sup> P. Krokovny,<sup>35</sup> F. Kruse,<sup>10</sup> W. Krzemien,<sup>29</sup> W. Kucewicz,<sup>27,o</sup> M. Kucharczyk,<sup>27</sup> V. Kudryavtsev,<sup>35</sup> A. K. Kuonen,<sup>40</sup> K. Kurek,<sup>29</sup> T. Kvaratskheliya,<sup>32,39</sup> D. Lacarrere,<sup>39</sup> G. Lafferty,<sup>55,39</sup> A. Lai,<sup>16</sup> D. Lambert,<sup>51</sup> G. Lanfranchi,<sup>19</sup> C. Langenbruch,<sup>49</sup> B. Langhans,<sup>39</sup> T. Latham,<sup>49</sup> C. Lazzeroni,<sup>46</sup> R. Le Gac,<sup>6</sup> J. van Leerdam,<sup>42</sup> J.-P. Lees,<sup>4</sup> A. Leflat,<sup>33,39</sup> J. Lefrançois,<sup>7</sup> R. Lefèvre,<sup>5</sup> F. Lemaître,<sup>39</sup> E. Lemos Cid,<sup>38</sup> O. Leroy,<sup>6</sup> T. Lesiak,<sup>27</sup> B. Leverington,<sup>12</sup> Y. Li,<sup>7</sup> T. Likhomanenko,<sup>67,66</sup> R. Lindner,<sup>39</sup> C. Linn,<sup>39</sup> F. Lionetto,<sup>41</sup> B. Liu,<sup>16</sup> X. Liu,<sup>3</sup> D. Loh,<sup>49</sup> I. Longstaff,<sup>52</sup> J. H. Lopes,<sup>2</sup> D. Lucchesi,<sup>23,k</sup> M. Lucio Martinez,<sup>38</sup> H. Luo,<sup>51</sup> A. Lupato,<sup>23</sup> E. Luppi,<sup>17,a</sup> O. Lupton,<sup>56</sup> A. Lusiani,<sup>24</sup> X. Lyu,<sup>62</sup> F. Machefert,<sup>7</sup> F. Maciuc,<sup>30</sup> O. Maev,<sup>31</sup> K. Maguire,<sup>55</sup> S. Malde,<sup>56</sup> A. Malinin,<sup>66</sup> T. Maltsev,<sup>35</sup> G. Manca,<sup>7</sup> G. Mancinelli,<sup>6</sup> P. Manning,<sup>60</sup> J. Maratas,<sup>5</sup> J. F. Marchand,<sup>4</sup> U. Marconi,<sup>15</sup> C. Marin Benito,<sup>37</sup> P. Marino,<sup>24,i</sup> J. Marks,<sup>12</sup> G. Martellotti,<sup>26</sup> M. Martin,<sup>6</sup> M. Martinelli,<sup>40</sup> D. Martinez Santos,<sup>38</sup> F. Martinez Vidal,<sup>68</sup> D. Martins Tostes,<sup>2</sup> L. M. Massacrier,<sup>7</sup> A. Massafferri,<sup>1</sup> R. Matev,<sup>39</sup> A. Mathad,<sup>49</sup> Z. Mathe,<sup>39</sup> C. Matteuzzi,<sup>21</sup> A. Mauri,<sup>41</sup> B. Maurin,<sup>40</sup> A. Mazurov,<sup>46</sup> M. McCann,<sup>54</sup> J. McCarthy,<sup>46</sup> A. McNab,<sup>55</sup> R. McNulty,<sup>13</sup> B. Meadows,<sup>58</sup> F. Meier,<sup>10</sup> M. Meissner,<sup>12</sup> D. Melnychuk,<sup>29</sup> M. Merk,<sup>42</sup> E. Michielin,<sup>23</sup> D. A. Milanes,<sup>64</sup> M.-N. Minard,<sup>4</sup> D. S. Mitzel,<sup>12</sup> J. Molina Rodriguez,<sup>61</sup> I. A. Monroy,<sup>64</sup> S. Monteil,<sup>5</sup> M. Morandin,<sup>23</sup> P. Morawski,<sup>28</sup> A. Mordà,<sup>6</sup> M. J. Morello,<sup>24,i</sup> J. Moron,<sup>28</sup> A. B. Morris,<sup>51</sup> R. Mountain,<sup>60</sup> F. Muheim,<sup>51</sup> M. Mulder,<sup>42</sup> M. Mussini,<sup>15</sup> D. Müller,<sup>55</sup> J. Müller,<sup>10</sup> K. Müller,<sup>41</sup> V. Müller,<sup>10</sup> P. Naik,<sup>47</sup> T. Nakada,<sup>40</sup> R. Nandakumar,<sup>50</sup> A. Nandi,<sup>56</sup> I. Nasteva,<sup>2</sup> M. Needham,<sup>51</sup> N. Neri,<sup>22</sup> S. Neubert,<sup>12</sup> N. Neufeld,<sup>39</sup> M. Neuner,<sup>12</sup> A. D. Nguyen,<sup>40</sup> C. Nguyen-Mau,<sup>40,p</sup> V. Niess,<sup>5</sup> S. Nieswand,<sup>9</sup> R. Niet,<sup>10</sup> N. Nikitin,<sup>33</sup> T. Nikodem,<sup>12</sup> A. Novoselov,<sup>36</sup> D. P. O'Hanlon,<sup>49</sup> A. Oblakowska-Mucha,<sup>28</sup> V. Obraztsov,<sup>36</sup> S. Ogilvy,<sup>19</sup> O. Okhrimenko,<sup>45</sup> R. Oldeman,<sup>48</sup> C. J. G. Onderwater,<sup>69</sup> J. M. Otalora Goicochea,<sup>2</sup> A. Otto,<sup>39</sup> P. Owen,<sup>54</sup> A. Oyangueren,<sup>68</sup> P. R. Pais,<sup>40</sup> A. Palano,<sup>14,q</sup> F. Palombo,<sup>22,m</sup> M. Palutan,<sup>19</sup> J. Panman,<sup>39</sup> A. Papanestis,<sup>50</sup> M. Pappagallo,<sup>52</sup> L. L. Pappalardo,<sup>17,a</sup> C. Pappenheimer,<sup>58</sup> W. Parker,<sup>59</sup> C. Parkes,<sup>55</sup> G. Passaleva,<sup>18</sup> G. D. Patel,<sup>53</sup> M. Patel,<sup>54</sup> C. Patrignani,<sup>15,f</sup> A. Pearce,<sup>55,50</sup> A. Pellegrino,<sup>42</sup> G. Penso,<sup>26,r</sup> M. Pepe Altarelli,<sup>39</sup> S. Perazzini,<sup>39</sup> P. Perret,<sup>5</sup> L. Pescatore,<sup>46</sup> K. Petridis,<sup>47</sup> A. Petrolini,<sup>20,h</sup> A. Petrov,<sup>66</sup> M. Petruzzo,<sup>22,m</sup> E. Picatoste Olloqui,<sup>37</sup> B. Pietrzyk,<sup>4</sup> M. Pikiés,<sup>27</sup> D. Pinci,<sup>26</sup> A. Pistone,<sup>20</sup> A. Piucci,<sup>12</sup> S. Playfer,<sup>51</sup> M. Plo Casasus,<sup>38</sup> T. Poikela,<sup>39</sup> F. Polci,<sup>8</sup> A. Poluektov,<sup>49,35</sup> I. Polyakov,<sup>32</sup> E. Polycarpo,<sup>2</sup> G. J. Pomery,<sup>47</sup> A. Popov,<sup>36</sup> D. Popov,<sup>11,39</sup> B. Popovici,<sup>30</sup> C. Potterat,<sup>2</sup> E. Price,<sup>47</sup> J. D. Price,<sup>53</sup> J. Prisciandaro,<sup>38</sup> A. Pritchard,<sup>53</sup> C. Prouve,<sup>47</sup> V. Pugatch,<sup>45</sup> A. Puig Navarro,<sup>40</sup> G. Punzi,<sup>24,s</sup> W. Qian,<sup>56</sup> R. Quagliani,<sup>7,47</sup> B. Rachwal,<sup>27</sup> J. H. Rademacker,<sup>47</sup> M. Rama,<sup>24</sup> M. Ramos Pernas,<sup>38</sup> M. S. Rangel,<sup>2</sup> I. Raniuk,<sup>44</sup> G. Raven,<sup>43</sup> F. Redi,<sup>54</sup> S. Reichert,<sup>10</sup> A. C. dos Reis,<sup>1</sup> C. Remon Alepuz,<sup>68</sup> V. Renaudin,<sup>7</sup> S. Ricciardi,<sup>50</sup> S. Richards,<sup>47</sup> M. Rihl,<sup>39</sup> K. Rinnert,<sup>53,39</sup> V. Rives Molina,<sup>37</sup> P. Robbe,<sup>7</sup> A. B. Rodrigues,<sup>1</sup> E. Rodrigues,<sup>58</sup> J. A. Rodriguez Lopez,<sup>64</sup> P. Rodriguez Perez,<sup>55</sup> A. Rogozhnikov,<sup>67</sup> S. Roiser,<sup>39</sup> V. Romanovskiy,<sup>36</sup> A. Romero Vidal,<sup>38</sup> J. W. Ronayne,<sup>13</sup> M. Rotondo,<sup>23</sup> T. Ruf,<sup>39</sup> P. Ruiz Valls,<sup>68</sup> J. J. Saborido Silva,<sup>38</sup> E. Sadykhov,<sup>32</sup> N. Sagidova,<sup>31</sup> B. Saitta,<sup>16,j</sup> V. Salustino Guimaraes,<sup>2</sup> C. Sanchez Mayordomo,<sup>68</sup> B. Sanmartin Sedes,<sup>38</sup> R. Santacesaria,<sup>26</sup> C. Santamarina Rios,<sup>38</sup> M. Santimaria,<sup>19</sup> E. Santovetti,<sup>25,g</sup> A. Sarti,<sup>19,r</sup> C. Satriano,<sup>26,b</sup> A. Satta,<sup>25</sup> D. M. Saunders,<sup>47</sup> D. Savrina,<sup>32,33</sup> S. Schael,<sup>9</sup> M. Schiller,<sup>39</sup> H. Schindler,<sup>39</sup> M. Schlupp,<sup>10</sup> M. Schmelling,<sup>11</sup> T. Schmelzer,<sup>10</sup> B. Schmidt,<sup>39</sup>

O. Schneider,<sup>40</sup> A. Schopper,<sup>39</sup> M. Schubiger,<sup>40</sup> M.-H. Schune,<sup>7</sup> R. Schwemmer,<sup>39</sup> B. Sciascia,<sup>19</sup> A. Sciubba,<sup>26,r</sup>  
 A. Semennikov,<sup>32</sup> A. Sergi,<sup>46</sup> N. Serra,<sup>41</sup> J. Serrano,<sup>6</sup> L. Sestini,<sup>23</sup> P. Seyfert,<sup>21</sup> M. Shapkin,<sup>36</sup> I. Shapoval,<sup>17,44,a</sup>  
 Y. Shcheglov,<sup>31</sup> T. Shears,<sup>53</sup> L. Shekhtman,<sup>35</sup> V. Shevchenko,<sup>66</sup> A. Shires,<sup>10</sup> B. G. Siddi,<sup>17</sup> R. Silva Coutinho,<sup>41</sup>  
 L. Silva de Oliveira,<sup>2</sup> G. Simi,<sup>23,k</sup> M. Sirendi,<sup>48</sup> N. Skidmore,<sup>47</sup> T. Skwarnicki,<sup>60</sup> E. Smith,<sup>54</sup> I. T. Smith,<sup>51</sup> J. Smith,<sup>48</sup>  
 M. Smith,<sup>55</sup> H. Snoek,<sup>42</sup> M. D. Sokoloff,<sup>58</sup> F. J. P. Soler,<sup>52</sup> D. Souza,<sup>47</sup> B. Souza De Paula,<sup>2</sup> B. Spaan,<sup>10</sup> P. Spradlin,<sup>52</sup>  
 S. Sridharan,<sup>39</sup> F. Stagni,<sup>39</sup> M. Stahl,<sup>12</sup> S. Stahl,<sup>39</sup> P. Stefko,<sup>40</sup> S. Stefkova,<sup>54</sup> O. Steinkamp,<sup>41</sup> O. Stenyakin,<sup>36</sup> S. Stevenson,<sup>56</sup>  
 S. Stoica,<sup>30</sup> S. Stone,<sup>60</sup> B. Storaci,<sup>41</sup> S. Stracka,<sup>24,i</sup> M. Straticiu,<sup>30</sup> U. Straumann,<sup>41</sup> L. Sun,<sup>58</sup> W. Sutcliffe,<sup>54</sup> K. Swientek,<sup>28</sup>  
 V. Syropoulos,<sup>43</sup> M. Szczekowski,<sup>29</sup> T. Szumlak,<sup>28</sup> S. T'Jampens,<sup>4</sup> A. Tayduganov,<sup>6</sup> T. Tekampe,<sup>10</sup> G. Tellarini,<sup>17,a</sup>  
 F. Teubert,<sup>39</sup> C. Thomas,<sup>56</sup> E. Thomas,<sup>39</sup> J. van Tilburg,<sup>42</sup> V. Tisserand,<sup>4</sup> M. Tobin,<sup>40</sup> S. Tolc,<sup>48</sup> L. Tomassetti,<sup>17,a</sup>  
 D. Tonelli,<sup>39</sup> S. Topp-Joergensen,<sup>56</sup> F. Toriello,<sup>60</sup> E. Tournefier,<sup>4</sup> S. Tourneur,<sup>40</sup> K. Trabelsi,<sup>40</sup> M. Traill,<sup>52</sup> M. T. Tran,<sup>40</sup>  
 M. Tresch,<sup>41</sup> A. Trisovic,<sup>39</sup> A. Tsaregorodtsev,<sup>6</sup> P. Tsopelas,<sup>42</sup> A. Tully,<sup>48</sup> N. Tuning,<sup>42</sup> A. Ukleja,<sup>29</sup> A. Ustyuzhanin,<sup>67,66</sup>  
 U. Uwer,<sup>12</sup> C. Vacca,<sup>16,39,j</sup> V. Vagnoni,<sup>15,39</sup> S. Valat,<sup>39</sup> G. Valenti,<sup>15</sup> A. Vallier,<sup>7</sup> R. Vazquez Gomez,<sup>19</sup> P. Vazquez Regueiro,<sup>38</sup>  
 S. Vecchi,<sup>17</sup> M. van Veghel,<sup>42</sup> J. J. Velthuis,<sup>47</sup> M. Veltri,<sup>18,t</sup> G. Veneziano,<sup>40</sup> A. Venkateswaran,<sup>60</sup> M. Vesterinen,<sup>12</sup> B. Viaud,<sup>7</sup>  
 D. Vieira,<sup>1</sup> M. Vieites Diaz,<sup>38</sup> X. Vilasis-Cardona,<sup>37,e</sup> V. Volkov,<sup>33</sup> A. Vollhardt,<sup>41</sup> B. Voneki,<sup>39</sup> D. Voong,<sup>47</sup> A. Vorobyev,<sup>31</sup>  
 V. Vorobyev,<sup>35</sup> C. Voß,<sup>65</sup> J. A. de Vries,<sup>42</sup> C. Vázquez Sierra,<sup>38</sup> R. Waldi,<sup>65</sup> C. Wallace,<sup>49</sup> R. Wallace,<sup>13</sup> J. Walsh,<sup>24</sup> J. Wang,<sup>60</sup>  
 D. R. Ward,<sup>48</sup> H. M. Wark,<sup>53</sup> N. K. Watson,<sup>46</sup> D. Websdale,<sup>54</sup> A. Weiden,<sup>41</sup> M. Whitehead,<sup>39</sup> J. Wicht,<sup>49</sup> G. Wilkinson,<sup>56,39</sup>  
 M. Wilkinson,<sup>60</sup> M. Williams,<sup>39</sup> M. P. Williams,<sup>46</sup> M. Williams,<sup>57</sup> T. Williams,<sup>46</sup> F. F. Wilson,<sup>50</sup> J. Wimberley,<sup>59</sup> J. Wishahi,<sup>10</sup>  
 W. Wislicki,<sup>29</sup> M. Witek,<sup>27</sup> G. Wormser,<sup>7</sup> S. A. Wotton,<sup>48</sup> K. Wraight,<sup>52</sup> S. Wright,<sup>48</sup> K. Wyllie,<sup>39</sup> Y. Xie,<sup>63</sup> Z. Xu,<sup>40</sup> Z. Yang,<sup>3</sup>  
 H. Yin,<sup>63</sup> J. Yu,<sup>63</sup> X. Yuan,<sup>35</sup> O. Yushchenko,<sup>36</sup> M. Zangoli,<sup>15</sup> K. A. Zarebski,<sup>46</sup> M. Zavertyaev,<sup>11,u</sup> L. Zhang,<sup>3</sup> Y. Zhang,<sup>7</sup>  
 Y. Zhang,<sup>62</sup> A. Zhelezov,<sup>12</sup> Y. Zheng,<sup>62</sup> A. Zhokhov,<sup>32</sup> V. Zhukov,<sup>9</sup> and S. Zucchelli<sup>15</sup>

(LHCb Collaboration)

<sup>1</sup>Centro Brasileiro de Pesquisas Físicas (CBPF), Rio de Janeiro, Brazil<sup>2</sup>Universidade Federal do Rio de Janeiro (UFRJ), Rio de Janeiro, Brazil<sup>3</sup>Center for High Energy Physics, Tsinghua University, Beijing, China<sup>4</sup>LAPP, Université Savoie Mont-Blanc, CNRS/IN2P3, Annecy-Le-Vieux, France<sup>5</sup>Clermont Université, Université Blaise Pascal, CNRS/IN2P3, LPC, Clermont-Ferrand, France<sup>6</sup>CPPM, Aix-Marseille Université, CNRS/IN2P3, Marseille, France<sup>7</sup>LAL, Université Paris-Sud, CNRS/IN2P3, Orsay, France<sup>8</sup>LPNHE, Université Pierre et Marie Curie, Université Paris Diderot, CNRS/IN2P3, Paris, France<sup>9</sup>I. Physikalisches Institut, RWTH Aachen University, Aachen, Germany<sup>10</sup>Fakultät Physik, Technische Universität Dortmund, Dortmund, Germany<sup>11</sup>Max-Planck-Institut für Kernphysik (MPIK), Heidelberg, Germany<sup>12</sup>Physikalisches Institut, Ruprecht-Karls-Universität Heidelberg, Heidelberg, Germany<sup>13</sup>School of Physics, University College Dublin, Dublin, Ireland<sup>14</sup>Sezione INFN di Bari, Bari, Italy<sup>15</sup>Sezione INFN di Bologna, Bologna, Italy<sup>16</sup>Sezione INFN di Cagliari, Cagliari, Italy<sup>17</sup>Sezione INFN di Ferrara, Ferrara, Italy<sup>18</sup>Sezione INFN di Firenze, Firenze, Italy<sup>19</sup>Laboratori Nazionali dell'INFN di Frascati, Frascati, Italy<sup>20</sup>Sezione INFN di Genova, Genova, Italy<sup>21</sup>Sezione INFN di Milano Bicocca, Milano, Italy<sup>22</sup>Sezione INFN di Milano, Milano, Italy<sup>23</sup>Sezione INFN di Padova, Padova, Italy<sup>24</sup>Sezione INFN di Pisa, Pisa, Italy<sup>25</sup>Sezione INFN di Roma Tor Vergata, Roma, Italy<sup>26</sup>Sezione INFN di Roma La Sapienza, Roma, Italy<sup>27</sup>Henryk Niewodniczanski Institute of Nuclear Physics Polish Academy of Sciences, Kraków, Poland<sup>28</sup>AGH - University of Science and Technology, Faculty of Physics and Applied Computer Science, Kraków, Poland<sup>29</sup>National Center for Nuclear Research (NCBJ), Warsaw, Poland<sup>30</sup>Horia Hulubei National Institute of Physics and Nuclear Engineering, Bucharest-Magurele, Romania<sup>31</sup>Petersburg Nuclear Physics Institute (PNPI), Gatchina, Russia



- <sup>32</sup>*Institute of Theoretical and Experimental Physics (ITEP), Moscow, Russia*
- <sup>33</sup>*Institute of Nuclear Physics, Moscow State University (SINP MSU), Moscow, Russia*
- <sup>34</sup>*Institute for Nuclear Research of the Russian Academy of Sciences (INR RAN), Moscow, Russia*
- <sup>35</sup>*Budker Institute of Nuclear Physics (SB RAS) and Novosibirsk State University, Novosibirsk, Russia*
- <sup>36</sup>*Institute for High Energy Physics (IHEP), Protvino, Russia*
- <sup>37</sup>*Universitat de Barcelona, Barcelona, Spain*
- <sup>38</sup>*Universidad de Santiago de Compostela, Santiago de Compostela, Spain*
- <sup>39</sup>*European Organization for Nuclear Research (CERN), Geneva, Switzerland*
- <sup>40</sup>*Ecole Polytechnique Fédérale de Lausanne (EPFL), Lausanne, Switzerland*
- <sup>41</sup>*Physik-Institut, Universität Zürich, Zürich, Switzerland*
- <sup>42</sup>*Nikhef National Institute for Subatomic Physics, Amsterdam, The Netherlands*
- <sup>43</sup>*Nikhef National Institute for Subatomic Physics and VU University Amsterdam, Amsterdam, The Netherlands*
- <sup>44</sup>*NSC Kharkiv Institute of Physics and Technology (NSC KIPT), Kharkiv, Ukraine*
- <sup>45</sup>*Institute for Nuclear Research of the National Academy of Sciences (KINR), Kyiv, Ukraine*
- <sup>46</sup>*University of Birmingham, Birmingham, United Kingdom*
- <sup>47</sup>*H.H. Wills Physics Laboratory, University of Bristol, Bristol, United Kingdom*
- <sup>48</sup>*Cavendish Laboratory, University of Cambridge, Cambridge, United Kingdom*
- <sup>49</sup>*Department of Physics, University of Warwick, Coventry, United Kingdom*
- <sup>50</sup>*STFC Rutherford Appleton Laboratory, Didcot, United Kingdom*
- <sup>51</sup>*School of Physics and Astronomy, University of Edinburgh, Edinburgh, United Kingdom*
- <sup>52</sup>*School of Physics and Astronomy, University of Glasgow, Glasgow, United Kingdom*
- <sup>53</sup>*Oliver Lodge Laboratory, University of Liverpool, Liverpool, United Kingdom*
- <sup>54</sup>*Imperial College London, London, United Kingdom*
- <sup>55</sup>*School of Physics and Astronomy, University of Manchester, Manchester, United Kingdom*
- <sup>56</sup>*Department of Physics, University of Oxford, Oxford, United Kingdom*
- <sup>57</sup>*Massachusetts Institute of Technology, Cambridge, Massachusetts, USA*
- <sup>58</sup>*University of Cincinnati, Cincinnati, Ohio, USA*
- <sup>59</sup>*University of Maryland, College Park, Maryland, USA*
- <sup>60</sup>*Syracuse University, Syracuse, New York, USA*
- <sup>61</sup>*Pontifícia Universidade Católica do Rio de Janeiro (PUC-Rio), Rio de Janeiro, Brazil (associated with Institution Universidade Federal do Rio de Janeiro (UFRJ), Rio de Janeiro, Brazil)*
- <sup>62</sup>*University of Chinese Academy of Sciences, Beijing, China (associated with Institution Center for High Energy Physics, Tsinghua University, Beijing, China)*
- <sup>63</sup>*Institute of Particle Physics, Central China Normal University, Wuhan, Hubei, China (associated with Institution Center for High Energy Physics, Tsinghua University, Beijing, China)*
- <sup>64</sup>*Departamento de Física, Universidad Nacional de Colombia, Bogota, Colombia (associated with Institution LPNHE, Université Pierre et Marie Curie, Université Paris Diderot, CNRS/IN2P3, Paris, France)*
- <sup>65</sup>*Institut für Physik, Universität Rostock, Rostock, Germany (associated with Institution Physikalisches Institut, Ruprecht-Karls-Universität Heidelberg, Heidelberg, Germany)*
- <sup>66</sup>*National Research Centre Kurchatov Institute, Moscow, Russia (associated with Institution Institute of Theoretical and Experimental Physics (ITEP), Moscow, Russia)*
- <sup>67</sup>*Yandex School of Data Analysis, Moscow, Russia (associated with Institution Institute of Theoretical and Experimental Physics (ITEP), Moscow, Russia)*
- <sup>68</sup>*Instituto de Física Corpuscular (IFIC), Universitat de Valencia-CSIC, Valencia, Spain (associated with Institution Universitat de Barcelona, Barcelona, Spain)*
- <sup>69</sup>*Van Swinderen Institute, University of Groningen, Groningen, The Netherlands (associated with Institution Nikhef National Institute for Subatomic Physics, Amsterdam, The Netherlands)*

<sup>a</sup>Also at Universidade Federal do Triângulo Mineiro (UFTM), Uberaba-MG, Brazil.

<sup>b</sup>Also at Università di Roma La Sapienza, Roma, Italy.

<sup>c</sup>Also at Università della Basilicata, Potenza, Italy.

<sup>d</sup>Also at Università di Urbino, Urbino, Italy.

<sup>e</sup>Also at Università di Ferrara, Ferrara, Italy.

<sup>f</sup>Also at P.N. Lebedev Physical Institute, Russian Academy of Science (LPI RAS), Moscow, Russia.

<sup>g</sup>Also at Università di Bari, Bari, Italy.

<sup>h</sup>Also at Università degli Studi di Milano, Milano, Italy.

<sup>i</sup>Also at Università di Roma Tor Vergata, Roma, Italy.

<sup>j</sup>Also at Scuola Normale Superiore, Pisa, Italy.

<sup>k</sup>Also at Università di Milano Bicocca, Milano, Italy.

<sup>l</sup>Also at Hanoi University of Science, Hanoi, Viet Nam.

<sup>m</sup>Also at Università di Padova, Padova, Italy.

<sup>n</sup>Also at AGH - University of Science and Technology, Faculty of Computer Science, Electronics and Telecommunications, Kraków, Poland.

<sup>o</sup>Also at Università di Cagliari, Cagliari, Italy.

<sup>p</sup>Also at Università di Genova, Genova, Italy.

<sup>q</sup>Also at Laboratoire Leprince-Ringuet, Palaiseau, France.

<sup>r</sup>Also at Università di Bologna, Bologna, Italy.

<sup>s</sup>Also at Università di Modena e Reggio Emilia, Modena, Italy.

<sup>t</sup>Also at Università di Pisa, Pisa, Italy.

<sup>u</sup>Also at LIFAELS, La Salle, Universitat Ramon Llull, Barcelona, Spain.