

Title: Association of Television viewing with mental health and mild cognitive impairment in the elderly in three European countries, data from the MentDis_ICF65+ project

Authors: Da Ronch C. (1), Canuto A. (2), Volkert J. (3), Massarenti S. (1), Weber K. (2), Dehoust M.C. (3), Nanni M.G. (1), Andreas S. (3,4), Sehner S. (5), Schulz H. (3), Härter M. (3), Grassi L. (1)

Affiliations:

1. Section of Psychiatry, Department of Biomedical and Specialty Surgical Sciences, Via Fossato di Mortara, 64/b, Ferrara 44121 Italy
2. Division of Liaison Psychiatry and Crisis Intervention, Department of Psychiatry and Mental Health, University Hospitals of Geneva (HUG), Rue Gabrielle Perret-Gentil 4, 1211 Geneva 14, Switzerland
3. Department of Medical Psychology, University Medical Centre Hamburg-Eppendorf, Martinistrasse 52, Building W 26, 20246 Hamburg, Germany
4. Institute of Psychology, Alpen-Adria Universität Klagenfurt, A-9020 Klagenfurt, Austria
5. Department of Biometry and Epidemiology, University Medical Centre Hamburg-Eppendorf, Martinistrasse 52, Building W 34, D-20246 Hamburg, Germany

E-Mail addresses:

SA: sandreas@uke.de

SS: s.sehner@uke.de

CDR: chiara.daronch@unife.it

HS: schulz@uke.de

AC: alessandra.canuto@hcuge.ch

MH: m.haerter@uke.de

KW: kerstin.weber@hcuge.ch

MGN: mariagiulia.nanni@unife.it

JV: jvolkert@uke.de

SM: sara.massarenti@unife.it

MCD: m.dehoust@uke.de

LG: luigi.grassi@unife.it

Word count:

abstract 236, article 3774 words, 2 tables, 1 graph, total 28 pages

Corresponding address:

Chiara Da Ronch

Section of Psychiatry

Department of Biomedical and Specialty Surgical Sciences

Via Fossato di Mortara, 64/b

Ferrara 44121 - Italy

Tel. +39 (0) 532 23 6409

Fax +39 (0) 532 21 2240

Association of Television viewing with mental health and mild cognitive impairment in the elderly in three European countries, data from the MentDis_ICF65+ project

Highlights

- There are no multi-country studies on the effects of TV on the elderly
- Data from the multi-centric MentDis_ICF65+ project were used
- 85.5% (n= 1175) of the sample watched TV for 5-7 days a week
- We found an association between TV viewing and Mild Cognitive Impairment
- No association was found between TV viewing and Mental Health

1
2
3 **Title: Association of Television viewing with mental health and mild cognitive**
4 **impairment in the elderly in three European countries, data from the MentDis_ICF65+**
5 **project**
6

7
8
9 Word count:

10 abstract 236, article 3774 words, 2 tables, 1 graph, total 28 pages
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Abstract

Background: Time spent watching TV by Europeans has been calculated to be 22.1 hours per week on average and it has shown to be correlated with a series of physical and mental problems in adults. Very little research is available in population over 65. This study aimed at evaluating the association between TV viewing and mental disorders and cognitive performance, taking into account the general physical activity level and socio-demographic characteristics in Europe.

Methods: Within the MentDis_ICF65+ study, a subsample of 1383 subjects aged 65-84 years were assessed by the Composite International Diagnostic Interview (CIDI65+) and the International Physical Activity Questionnaire (IPAQ) for physical activity evaluation. Time spent in watching TV was assessed through a self report instrument.

Results: Forty-three per cent of the total sample watched TV for 5-7 days a week for 2 or more hours every day. Females, people who lived alone, older subjects and those with lower education significantly watched TV for a longer time. Stepwise multiple regression showed statistically significant inverse correlation between Mini-Mental State Examination scores and TV viewing time ($p < 0.001$). Apart from a negative association with Major Depressive Disorder, no particular associations were found between TV viewing and psychopathological diagnoses.

Conclusions: Given the relationship of time spending watching TV with cognitive impairment, awareness should be raised about the possible negative effects of TV viewing on the elderly and programs to reduce TV viewing time should be set up.

Key words: elderly, sedentary behaviour, television, mental health, cognition, Europe

1. Introduction

Television (TV) viewing is a common, widespread way of spending leisure time at all ages. Time spent watching TV by Europeans has lately been calculated to be 22.1 hours per week on average (Nationmaster, 2009). In younger adults, TV viewing, frequently used as a measure of sedentary behaviour, has been correlated with a number of serious health risks, such as higher risk of obesity and type 2 diabetes mellitus in women (Hu, Li, Colditz, Willett, & Manson, 2003), and in men (Grontved & Hu, 2011; Hu et al., 2001), higher risk of obesity and cardiovascular diseases (Bowman, 2006; Fung et al., 2000; Jakes et al., 2003; Marshall, Biddle, Gorely, Cameron, & Murdey, 2004), neck and low back pain (Kuster, 2004), metabolic syndrome (Dunstan et al., 2005; Gao, Nelson, & Tucker, 2007; Wijndaele et al., 2009), atherosclerosis (Kronenberg et al., 2000), sleep problems (Basner & Dinges, 2009) and others (Dunstan, Howard, Healy, & Owen, 2012; Grontved & Hu, 2011).

In the young/adult sector of the population, TV viewing has also been associated with risk for mental health disorders. In a Spanish study, Sanchez and colleagues found an association between sedentary activities (watching TV and using the computer) and the incidence of mental disorders (Sanchez-Villegas et al., 2008). In a Scottish survey, Hamer et al. concluded that TV and screen-based entertainment were associated with poorer mental health scores in a representative population sample of adults (Hamer, Stamatakis, & Mishra, 2010). De Wit and colleagues report that, independently of physical activity amount, sedentary behaviors occur more frequently among persons with a mental disorder. In particular in this Dutch study, TV viewing was higher in people with dysthymia, panic disorder and agoraphobia compared to controls (de Wit, van Straten, Lamers, Cuijpers, & Penninx, 2011). Atkin, Adams et al. found a negative association between sedentary activities (including TV viewing) and mental health scores in women in the UK (Atkin, Adams, Bull, & Biddle, 2012). Finally, in an important longitudinal study, the English Longitudinal Study

1 for Ageing, Hamer and associates found a cross-sectional and longitudinal association
2 between TV viewing and depressive symptoms (Hamer, Poole, & Messerli-Burgy, 2013;
3 Hamer & Stamatakis, 2013).
4
5

6
7 Furthermore, TV viewing has been linked to a possible increased risk of cognitive
8 impairment. In the previously cited Longitudinal study, Hamer et al. found an inverse
9 association between TV viewing and cognitive function levels (Hamer & Stamatakis, 2013).
10
11 A 5-year longitudinal study in China found that watching television was associated with a
12 higher risk of developing a cognitive impairment (Wang et al., 2006). Finally, another 6-
13 years longitudinal study done in France found a negative association between time spent
14 watching TV and executive functioning (Kesse-Guyot et al., 2012).
15
16
17
18
19
20
21
22
23

24 A recent review on the prevalence of sitting behaviour in people of 60+ years indicated
25 that over 55% referred watching more than 2 hours of TV daily . However, while the
26 previously cited studies have mostly focussed on adults, little has been published on the
27 effects of TV viewing on people over 65 years old (Kikuchi et al., 2013; Lucas et al., 2011).
28
29
30
31
32
33

34 A recent review on the prevalence of sitting behaviour in of people of 60+ years indicated
35 that over 55% referred watching more than 2 hours of TV daily (Harvey, Chastin, & Skelton,
36 2013). Furthermore, all currently available data have been derived from single-country
37 studies and no international or cross-national studies exist.
38
39
40
41
42

43 The present study aims at providing some further elements in order to understand how TV
44 viewing relates with elderly mental health and cognitive performance across three European
45 countries.
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

2. Methods

2.1 Sample

Data have been gathered as part of the MentDis_ICF65+ study, on “Prevalence, 1-year incidence and symptom severity of mental disorders in the elderly: Relationship to impairment, functioning (ICF) and service utilization” (MentDis_ICF65+) funded by the European Commission within the 7th framework program. Aims, design, and methods have been described in greater detail elsewhere (Andreas et al., 2013) and are only shortly reported here.

The MentDis_ICF65+ study is a multicentre research project designed to adapt existing diagnostic tools and questionnaires to the elderly, and then to gather data on the prevalence (cross-sectional phase) and incidence (longitudinal phase) of mental disorders in the general 65-84 years population across different countries of the European Union (Spain, Italy, UK, Germany) and two other related states (Switzerland, Israel). The sample was age- and gender-stratified and randomly selected.

Inclusion criteria were living in the defined catchment area, being aged between 65 and 84 years, having a sufficient level of corresponding language and a Mini Mental State Examination score indicating sufficient cognitive function (MMSE>18, age and education adjusted). In this study we examined results from a subsample of 1383 subjects from three centres, Italy, Switzerland and Germany, where specific additional data on TV viewing and physical activity were gathered.

The sample consisted of 1383 subjects. There were 52.4% (n= 725) male, 47.6% (n= 658) female, mean age 72.5 (S.D. 5.69), 35.1% (n= 486) from Ferrara (Italy), 27.5% (n= 380) from Hamburg (Germany) and 37.4% (n= 517) from Geneva (Switzerland).

1 The majority of people interviewed were living with someone (overall 70.8%; 70.5% in
2 Hamburg, 77.6% in Ferrara, 64.6% in Geneva, $p < .001$) while only 29.2% were living alone.
3
4 The participants attended school for 10.3 years on average (10.3 in Hamburg, 8.8 in Ferrara
5 and 11.8 in Geneva, $p < .001$) and most graduated from the last school they attended (88.7%
6 overall 96.8%, in Hamburg, 88.4% in Ferrara, 83.2% in Geneva, $p < .001$). At the time of the
7 interview the vast majority of participants were not working (94.7% overall, 97.9% in
8 Hamburg, 97.0% in Ferrara, 90.3% in Geneva, $p < .001$).
9
10
11
12
13
14
15
16
17

18 **2.2 Measurements**

19 For the present analyses we used the data of the first wave (cross-sectional) of the
20
21 MentDis_ICF65+ study. All participants underwent a global assessment during individual
22 meetings either at their homes or at the local research structures between January and October
23 2011 and gave written informed consent for participation according to the local institutional
24 Ethics Committees (Andreas, et al., 2013). Such evaluation covered a wide range of physical
25 as well as mental health aspects, socio-demographic data, physical activity, and TV viewing
26 and was carried out both through semi-structured interview and self reported information.
27
28
29
30
31
32
33
34
35
36
37
38

39 **2.2.1 Depressive and anxiety disorder diagnoses**

40 Data on mental health were collected using a Computer Assisted Personal Interviewing
41 (CAPI) version of the *Composite International Diagnostic Interview for the Elderly*
42 (CIDI65+), the adjusted version of the CIDI (Haro et al., 2006; Kessler et al., 2004; Wittchen
43 et al., 1991) and adapted by the study group to the particular social, cognitive and
44 psychological abilities and needs of the elderly. The syndrome domains and sections, the
45 translation and adaptation procedure and the psychometric assessment of the CIDI65+ are
46 described elsewhere in more detail (Wittchen et al., 2014). Interviewers received extensive
47 training before the start of the study and were monitored and supervised constantly during the
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 data collection phase. The same procedure was followed in each centre and the same –
2 translated - version of the CIDI 65+ was used. The interview covers several mental health
3
4 problems including anxiety disorders and affective disorders according to the Diagnostic and
5
6 Statistical Manual for Mental Disorders, DSM-IV criteria (APA, 2000; Wittchen, 1994) and
7
8 leads to specific diagnoses according to pre-defined diagnostic algorithms taking also into
9
10 account diagnostic hierarchy rules and possible organic exclusions. The present study
11
12 focused on depressive disorders such as Dysthymic disorder (with hierarchy), Major
13
14 depressive disorder (MDD), as well as anxiety disorders, including Agoraphobia, Generalized
15
16 anxiety disorder (GAD6 - DSM-IV), Social phobia (lifetime), any Simple phobia, Panic
17
18 disorder. In a previous similar work by de Wit et al (de Wit, et al., 2011) the timeframe chose
19
20 for presence of diagnosis was the 6 months preceding the study. For comparability purposes,
21
22 as our study only had current (one month), one year and lifetime available, we chose to select
23
24 the one-year timeframe for inclusion as closer to the 6-months criterion.
25
26
27
28
29
30

31 32 **2.2.2 Cognitive impairment**

33
34
35 Cognitive functioning was gathered through the Mini Mental State Evaluation (MMSE).
36
37 This is a short tool (30 items with an average delivery time of 10 minutes) normally used for
38
39 cognitive impairment screening and severity evaluation (Folstein, Folstein, & McHugh,
40
41 1975). It needs to be adjusted for age and education (O'Connor, Pollitt, Treasure, Brook, &
42
43 Reiss, 1989) and even though it does not provide a definitive diagnosis for dementia, it has
44
45 sufficient psychometric properties to be considered useful to generally investigate cognitive
46
47 functioning (Crum, Anthony, Bassett, & Folstein, 1993; Tombaugh & McIntyre, 1992).
48
49
50

51
52 In this study people with severe cognitive impairment were excluded. People were
53
54 considered as having a Mild Cognitive impairment with a 18-26 MMSE score, while with 27-
55
56 30 score they were considered with no cognitive impairment.
57
58
59
60
61
62
63
64
65

2.2.3 Television viewing

1
2 Time spent watching TV in the past week was collected by a self report instrument.
3
4 Similarly to the questions regarding physical activity, the subjects were asked to indicate on
5
6 how many days they watched TV during the previous week (0-7) and to indicate for how long
7
8 on average each day by choosing between six options (<10 minutes, 10-30 minutes, 30 min-1
9
10 hour, 1-2 hours, 2-3 hours, >3 hours). The weekly hours TV viewed number was calculated
11
12 by multiplying the average time spent daily (e.g. for the “1-2 hours” category we considered
13
14 150 minutes) by the number of days (range 0-1260 minutes per week).
15
16
17
18
19

2.2.4 Covariates

20
21
22
23 Analyses were corrected for possible confounding variables, specifically gender, age,
24
25 level of education (high: >13 years, intermediate: 9-12 years, low: <8 years), employment
26
27 status (unemployed/retired, employed), financial situation (very good, good, just enough or
28
29 very low), living status (alone, not alone), study centre (Ferrara, Hamburg, Geneva). Physical
30
31 activity levels were gathered through the self-report International Physical Activity
32
33 Questionnaire (IPAQ), using the short last 7 days telephone version for time saving purposes,
34
35 even though it was used during the face to face interview (Craig et al., 2003; Hallal &
36
37 Victora, 2004). This simple questionnaire gathers information on the time that people spent
38
39 doing three different types of physical activities (vigorous, moderate and walking) during the
40
41 previous week. The original version included a fourth area related to the sedentary behaviour
42
43 that included time spent watching TV. For the purposes of our study the question was slightly
44
45 modified to separate TV viewing time from the time spent in other sedentary behaviours
46
47 (meal, reading, travelling by car and so on).
48
49
50
51
52
53

54
55 In order to facilitate completion by the elderly, instead of asking to recollect for how
56
57 many hours and minutes they performed each type of activity on average per day, they were
58
59 asked to choose between six options (<10 minutes, 10-30 minutes, 30 min-1 hour, 1-2 hours,
60
61
62
63
64
65

1 2-3 hours, >3 hours). In fact less than 10 minutes is considered not sufficient physical activity
2 (Sun, Norman, & While, 2013), more than 3 hours can be considered as a very high level of
3 physical activity for the 65-84 age group.
4
5

6
7 Based on such data it was possible to compute an overall energy expenditure estimate in
8 Metabolic Equivalent Total (MET)-units. This is a standard and comparable way to measure
9 the energy “cost” of physical activities: it allows to reduce to a single number different types
10 of activities (Sedentary-Behaviour-Research-Network, 2013). The Total weekly MET
11 minutes were derived from multiplying duration of activity times frequency of activity times
12 MET intensity, as provided within the IPAQ guidelines (IPAQ-Group, 2005). Low physical
13 activity has been considered when total weekly MET were lower than 600, moderate between
14 600 and 3000, high levels when higher than 3000.
15
16

17
18 Physical activity and sedentary behaviour can be influenced also by the level of
19 functioning of the person (the inverse concept of Disability (WHO, 2002)) and by their
20 physical health. Therefore we included the WHODAS II simple sum score derived from the
21 12-item self-administered version of the WHO Disability Assessment Schedule II
22 (WHODAS-II) (Sousa et al., 2010), used to assess level of functioning in relation to
23 cognition, mobility, self-care, getting along, life activities and participation. Furthermore we
24 included the self-reported number of diagnosed medical illnesses in the previous year.
25
26

27 **2.3 Statistical analyses**

28
29 IBM SPSS 20.0 was used to perform all statistical analyses. We computed TV viewing
30 outcomes and covariates in relation to the depression and anxiety diagnosis (control subjects
31 with no depression nor anxiety, depression alone, anxiety alone, comorbid anxiety and
32 depression) and to cognitive functioning (mild cognitive impairment, no cognitive
33 impairment) and tested differences with χ^2 analyses or T-test where appropriate. Stepwise
34 multiple regression analyses, adjusted for covariates (as described above: gender, age, level
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 of education, employment status, financial situation, living status, study centre, physical
2 activity level (weekly MET total), Level of Functioning (WHODAS II score) were performed
3
4 to test the association between TV viewing as dependant variable (between 0 and 1260
5 minutes per week)) and cognitive impairment rated as continuous MMSE score, as well as
6
7 diagnosis of depression/anxiety as independent variables, rated as binary variable (0= control,
8
9 1= case). We analysed diagnoses both as general diagnoses and including the specific
10
11 diagnoses (Dysthymic disorder (with hierarchy), Major depressive disorder (MDD),
12
13 Agoraphobia, Generalized anxiety disorder (GAD6 - DSM-IV), Social phobia (lifetime), Any
14
15 simple phobia, Panic disorder).
16
17
18
19
20
21
22

23 3. Results

24
25 The sample included a total of 1383 subjects, 166 had to be excluded since the
26
27 questionnaires were not complete (IPAQ-Group, 2005). Regarding psychiatry diagnoses,
28
29 13.4% (n= 216) presented a depressive disorder only (13.2% MDD and 3.0% Dysthymia),
30
31 15.6% (n=186) an anxiety disorder (3.3% agoraphobia, 3.0% GAD, 1.4% social phobia, 7.5%
32
33 simple phobia, 2.9% PD) and 4.3% (n= 59) comorbid anxiety and depression disorder. The
34
35 subjects with depressive and/or anxiety disorders were more often female (p< 0.001),
36
37 younger (p< 0.01), living more often alone than controls (p< 0.05), with lower levels of
38
39 functioning (p< 0.001) and a higher rate of medical diagnoses (p< 0.001). Regarding the
40
41 variable “financial situation” , those with depressive disorder more frequently reported to
42
43 have a “good” situation than the others (p< 0.05).
44
45
46
47
48
49

50 Concerning cognitive functioning 251 subjects (18.1%) presented a mild cognitive
51
52 impairment (MCI) and 1132 (81.9%) no cognitive impairment. MCI was more common in
53
54 female (p< 0.05), older people (p< 0.001), with lower levels of education (p< 0.001), living
55
56 alone (p< 0.05), from Geneva (p< 0.001), with lower levels of functioning (p< 0.001), and
57
58 with lower levels of physical activity (p< 0.001). People with MCI watched an average of
59
60
61
62
63
64
65

1 1.91 (S.D. 0.95) hours TV per day while people with no cognitive impairment watched only
2
3 1.61 (S.D. 0.97) ($p < 0.001$).

4 The average daily hours of TV viewing was 1.81 (S.D. 0.94, range 0-3), sitting only was
5
6 2.01 (S.D. 0.96 range 0-3) and total daily sedentary time was 3.7 (S.D. 1.44, range 0-6).
7
8 Additionally, 85.5% ($n = 1175$) of the sample watched 5-7 days/week (43.75% ($n = 605$) for
9
10 >2 hours per day, 23.9% ($n = 330$) for 1-2 hour per day, 17.4% ($n = 240$) for < 1 hour per day),
11
12 7.8% ($n = 108$) watched for 3-4 days, 7.2% watched between 0 and 2 days ($n = 100$), only
13
14 3.3% ($n = 45$) of the total sample declared to have never watched TV in the previous week
15
16 ($p < 0.001$) (see graph 1).
17
18
19
20
21
22
23

24 --- *Insert graph 1 around here* ---
25
26
27
28

29 There was no association between having a diagnosis of depression, anxiety or both and
30
31 time spent viewing TV ($p = 0.61$) or any sedentary activity ($p = 0.64$) nor levels of physical
32
33 activity ($p = 0.09$). However, there was a general difference between controls and people with
34
35 a diagnosis in terms of physical activity levels (Weekly MET Total) ($p < 0.05$, $F = 10.462$, df
36
37 1381).
38
39
40

41 Subjects with MCI presented a higher time spent watching TV ($p < 0.001$) and total
42
43 sedentary time (sitting plus TV viewing) ($p < 0.001$) but not with higher time in other sitting
44
45 time ($p = 0.31$).
46
47
48
49
50

51 --- *Insert table 1 around here* ---
52
53
54
55
56

57 ***3.1 Psychiatric diagnoses, Cognitive impairment and TV use*** 58 59 60 61 62 63 64 65

1 Stepwise multiple regression analyses, adjusted for covariates (gender, age, study centre,
2 level of education, living status, work status (employed vs. retired or unemployed), level of
3 functioning, number of physical illnesses, physical activity), were employed to examine
4 further the hypothesized association with either a general depressive or anxiety disorder, or
5 more specific ones with time spent watching TV. According to the model produced no
6 significant association was found between a general diagnosis of either anxiety or depression
7 or both and time spent watching TV. There was only a significant negative association
8 between the specific diagnosis of Major Depressive Disorder (MDD) and TV viewing (see
9 table 2). On the other side cognitive functioning (MMSE score) and TV viewing were
10 inversely related, after adjusting for covariates. The results can be summarized in a model
11 with 6 significant predictors of TV viewing. Lower average time of TV watching per day is
12 significantly predicted by the following factors: higher level of education, active work status,
13 higher cognitive functioning (as assessed by MMSE scores), higher level of physical activity,
14 and presence of Major Depressive Disorder (MDD). In contrast, severity of comorbid
15 physical illnesses (number of medical diagnoses) increases the risk for high average time of
16 TV watching (see table 2).
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40

41 --- Insert table 2 around here---
42
43
44

45 **4. Discussion**

46
47 The hypothesis of a relationship between mental health and TV viewing in the elderly
48 general population was only partially confirmed in this study. While a general association
49 between TV viewing and general diagnoses of depression, anxiety or both was not present, it
50 emerged that persons with a Major Depressive Disorder (MDD) tended to watch less
51 Television than those without such a diagnosis. These results were significant after
52 controlling for possible covariates.
53
54
55
56
57
58
59
60
61
62
63
64
65

1 This result is in general not in line with other studies that found a relationship between
2 sedentary behaviour and mental health (Arredondo et al., 2013; Hamer & Stamatakis, 2013;
3 Hamer, et al., 2010; Lucas, et al., 2011; Teychenne, Ball, & Salmon, 2010). A possible
4 hypothesis of such result may refer to the clinical features of such condition, that include the
5 characteristic anhedonia, i.e. inability to enjoy previously appreciated activities such as TV
6 viewing. Furthermore the task of recalling how many hours of such activity per day might be
7 influenced by the possible memory diminished efficiency with an underestimation of the
8 actual amount.
9

10
11
12
13
14
15
16
17
18
19 On the other side it was demonstrated that, even after controlling for possible covariates,
20 subjects with lower cognitive functioning have a tendency to spend more time viewing
21 television than those with no cognitive impairment. This is in agreement with previous
22 research (Fogel & Carlson, 2006; Hamer & Stamatakis, 2013; Kesse-Guyot, et al., 2012;
23 Lindstrom et al., 2005; Rundek & Bennett, 2006; Wang, et al., 2006).
24
25
26
27
28
29
30

31 Furthermore, the need to separately evaluate different types of sedentary activity, in
32 particular to assess TV viewing time, as there are different associations and effects (Hamer &
33 Stamatakis, 2013; Kesse-Guyot, et al., 2012), was confirmed in our study.
34
35
36
37
38

39 The main strength of these findings, in comparison with the previously cited studies,
40 regards the target sample, since, to our knowledge, this is the first study investigating the
41 association of TV viewing on the general elderly population mental health and cognitive
42 functioning across different European Countries.
43
44
45
46
47

48 The main limitation of this study refers to the measure of TV viewing. More specifically,
49 the time categories proposed (maximum value “3 hours or more”) limited the power of the
50 analyses and reflected only a normal use of the TV and sitting time, not the extremes or
51 abuses. A scale with up to 12 hours per day would have been more accurate in measuring the
52 real behaviour. Even better, in order to avoid possible biases due to memory-based self-
53
54
55
56
57
58
59
60
61
62
63
64
65

1 reporting instruments, would be to gather direct information and measures of actual
2 behaviour through, for example, an accelerometer and a measure of electronic TV monitor
3
4 (Otten, Littenberg, & Harvey-Berino, 2010). Another limitation is the cross-sectional nature
5
6 of the study as it prevents us from inferring any causal relationship between variables, and we
7
8 can only describe associations. We also are aware of the fact that that self-report measures
9
10 generally tend to underestimate the real behaviour (Harvey, et al., 2013) and both TV
11
12 viewing and sitting time could have been even higher than what was reported. Moreover,
13
14 since the Mini Mental State Examination is mainly a screening tool, not specifically designed
15
16 to perform a detailed evaluation of cognitive functioning (Folstein, et al., 1975), more
17
18 specific measures of cognitive efficiency should be used in further studies. Finally, regarding
19
20 the selection process it is possible that the final sample did not include the most fragile or
21
22 sick at the time of the recruitment therefore our subjects could be slightly more “sane”, and
23
24 probably more active, than the real population.
25
26
27
28
29
30

31
32
33
34 In conclusion, even with such limits, this research has provided further evidence of the
35
36 association between time spent watching Television and some specific characteristics, in
37
38 particular Mild Cognitive Impairment (MCI), in the 65+ population. The interesting result
39
40 regarding the inverse association found between TV viewing and Major Depressive Disorder
41
42 would deserve further investigations, taking into account further variables, such as, for
43
44 example, sleeping hours or other leisure activities such as reading, listening to the radio or
45
46 having the possibility of daily social interaction or not (i.e. level of isolation).
47
48
49

50
51 Television viewing might displace more physically and/or mentally challenging activities
52
53 (Sugiyama, Healy, Dunstan, Salmon, & Owen, 2008; Teychenne, Ball, & Salmon, 2008) thus
54
55 preventing people from engaging in an healthier lifestyle. The inverse association between
56
57 Television time and Physical activity levels found in our data would also support such
58
59
60
61
62
63
64
65

1 statement. However recent studies demonstrated that sitting for long time (thus typically TV
2 viewing) per se could be harmful for health, independently of time spent on physical
3 activities (Ekelund et al., 2006; M. T. Hamilton, Hamilton, & Zderic, 2004, 2007; M. T.
4 Hamilton, Healy, Dunstan, Zderic, & Owen, 2008; Katzmarzyk, Church, Craig, & Bouchard,
5 2009). It is well known how challenging it could be to modify people’s habits and increase
6 their physical activity (Sun, et al., 2013). An important step to promoting a better ageing
7 could be to start from increasing awareness of the negative impact of sitting time and design
8 programs aimed at drastically reduce the time spent viewing TV while increasing time spent
9 on other more challenging mental activities such as reading, playing cards, interacting with
10 other people (Chen et al., 2012; Fratiglioni, Paillard-Borg, & Winblad, 2004; Paillard-Borg,
11 Fratiglioni, Xu, Winblad, & Wang, 2012; Wang, et al., 2006). This should be done bearing in
12 mind that the most at risk people could be those with a Mild Cognitive Impairment. As
13 Hamer points out, “even small adverse health effects of this behaviour (TV viewing) could
14 have profound impact at the population level” (Hamer & Stamatakis, 2013).
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34

35 **Acknowledgments**

36 All participants to the study are gratefully acknowledged. We also express our gratitude
37 to all interviewers, dedicated collaborators and local institutions that made this study
38 possible.
39
40
41
42
43
44

45 **Funding Sources:** This study is funded by a grant from the European Commission (Grant
46 No: 223105) within the 7th Framework Research Program of the EU.
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

References

- Andreas, S., Harter, M., Volkert, J., Hausberg, M., Sehner, S., Wegscheider, K., . . . Schulz, H. (2013). The MentDis_ICF65+ study protocol: prevalence, 1-year incidence and symptom severity of mental disorders in the elderly and their relationship to impairment, functioning (ICF) and service utilisation. *BMC Psychiatry*, 13, 62. doi: 10.1186/1471-244X-13-62
- 1471-244X-13-62 [pii]
- APA. (2000). *Diagnostic and statistical manual of mental disorders : DSM-IV-TR (4th ed., text revision. ed.)*. Washington, DC: American Psychiatric Association.
- Arredondo, M. A., Lemus, H., Elder, J. P., Molina, M., Martinez, S., Sumek, C., & Ayala, G. X. (2013). The relationship between sedentary behavior and depression among Latinos. *Ment Health Phys Act*, 6, 3-9.
- Atkin, A. J., Adams, E., Bull, F. C., & Biddle, S. J. (2012). Non-occupational sitting and mental well-being in employed adults. *Ann Behav Med*, 43(2), 181-188. doi: 10.1007/s12160-011-9320-y
- Basner, M., & Dinges, D. F. (2009). Dubious bargain: trading sleep for Leno and Letterman. *Sleep*, 32(6), 747-752.
- Bowman, S. A. (2006). Television-viewing characteristics of adults: correlations to eating practices and overweight and health status. *Prev Chronic Dis*, 3(2), A38. doi: A38 [pii]
- Chen, X. R., Jiang, Y., Wang, L. M., Li, Y. C., Zhang, M., Hu, N., & Zhao, W. H. (2012). [Leisure-time physical activity and sedentary behaviors among Chinese adults in 2010]. *Zhonghua Yu Fang Yi Xue Za Zhi*, 46(5), 399-403.

1 Craig, C. L., Marshall, A. L., Sjostrom, M., Bauman, A. E., Booth, M. L., Ainsworth, B. E., .

2 . . Oja, P. (2003). International physical activity questionnaire: 12-country reliability
3 and validity. *Med Sci Sports Exerc*, 35(8), 1381-1395. doi:
4

5 10.1249/01.MSS.0000078924.61453.FB
6

7
8
9 Crum, R. M., Anthony, J. C., Bassett, S. S., & Folstein, M. F. (1993). Population-based
10 norms for the Mini-Mental State Examination by age and educational level. *JAMA*,
11
12 269(18), 2386-2391.
13
14

15
16 de Wit, L., van Straten, A., Lamers, F., Cuijpers, P., & Penninx, B. (2011). Are sedentary
17 television watching and computer use behaviors associated with anxiety and
18
19 depressive disorders? *Psychiatry Res*, 186(2-3), 239-243. doi:
20
21

22 10.1016/j.psychres.2010.07.003
23
24

25
26 S0165-1781(10)00377-X [pii]
27

28
29 Dunstan, D. W., Howard, B., Healy, G. N., & Owen, N. (2012). Too much sitting--a health
30 hazard. *Diabetes Res Clin Pract*, 97(3), 368-376. doi: 10.1016/j.diabres.2012.05.020
31
32

33
34 S0168-8227(12)00208-2 [pii]
35

36
37 Dunstan, D. W., Salmon, J., Owen, N., Armstrong, T., Zimmet, P. Z., Welborn, T. A., . . .

38
39 Shaw, J. E. (2005). Associations of TV viewing and physical activity with the
40 metabolic syndrome in Australian adults. *Diabetologia*, 48(11), 2254-2261. doi:
41
42

43 10.1007/s00125-005-1963-4
44
45

46
47 Ekelund, U., Brage, S., Froberg, K., Harro, M., Anderssen, S. A., Sardinha, L. B., . . .

48
49 Andersen, L. B. (2006). TV viewing and physical activity are independently
50 associated with metabolic risk in children: the European Youth Heart Study. *PLoS*
51
52

53
54 *Med*, 3(12), e488. doi: 06-PLME-RA-0438R2 [pii]
55

56 10.1371/journal.pmed.0030488
57
58
59
60
61
62
63
64
65

1 Fogel, J., & Carlson, M. C. (2006). Soap operas and talk shows on television are associated
2 with poorer cognition in older women. *South Med J*, 99(3), 226-233.
3

4 Folstein, M. F., Folstein, S. E., & McHugh, P. R. (1975). "Mini-mental state." A practical
5 method for grading the cognitive state of patients for the clinician. *Journal of*
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Psychiatric Research, 12(3), 189-198.

Fratiglioni, L., Paillard-Borg, S., & Winblad, B. (2004). An active and socially integrated
lifestyle in late life might protect against dementia. *Lancet Neurol*, 3(6), 343-353. doi:
10.1016/S1474-4422(04)00767-7
S1474442204007677 [pii]

Fung, T. T., Hu, F. B., Yu, J., Chu, N. F., Spiegelman, D., Tofler, G. H., . . . Rimm, E. B.
(2000). Leisure-time physical activity, television watching, and plasma biomarkers of
obesity and cardiovascular disease risk. *Am J Epidemiol*, 152(12), 1171-1178.

Gao, X., Nelson, M. E., & Tucker, K. L. (2007). Television viewing is associated with
prevalence of metabolic syndrome in Hispanic elders. *Diabetes Care*, 30(3), 694-700.
doi: 30/3/694 [pii]
10.2337/dc06-1835

Grontved, A., & Hu, F. B. (2011). Television viewing and risk of type 2 diabetes,
cardiovascular disease, and all-cause mortality: a meta-analysis. *JAMA*, 305(23),
2448-2455. doi: 10.1001/jama.2011.812
305/23/2448 [pii]

Hallal, P. C., & Victora, C. G. (2004). Reliability and validity of the International Physical
Activity Questionnaire (IPAQ). *Med Sci Sports Exerc*, 36(3), 556. doi: 00005768-
200403000-00027 [pii]

1 Hamer, M., Poole, L., & Messerli-Burgy, N. (2013). Television viewing, C-reactive protein,
2 and depressive symptoms in older adults. *Brain Behav Immun*, 33, 29-32. doi:
3
4 10.1016/j.bbi.2013.05.001
5
6

7 S0889-1591(13)00181-5 [pii]
8

9 Hamer, M., & Stamatakis, E. (2013). Prospective Study of Sedentary Behavior, Risk of
10 Depression, and Cognitive Impairment. *Med Sci Sports Exerc*. doi:
11
12 10.1249/MSS.0000000000000156
13
14
15

16 Hamer, M., Stamatakis, E., & Mishra, G. D. (2010). Television- and screen-based activity
17 and mental well-being in adults. *Am J Prev Med*, 38(4), 375-380. doi:
18
19 10.1016/j.amepre.2009.12.030
20
21
22

23 S0749-3797(10)00010-3 [pii]
24

25 Hamilton, M. T., Hamilton, D. G., & Zderic, T. W. (2004). Exercise physiology versus
26 inactivity physiology: an essential concept for understanding lipoprotein lipase
27 regulation. *Exerc Sport Sci Rev*, 32(4), 161-166. doi: 00003677-200410000-00007
28
29
30
31
32
33 [pii]
34
35

36 Hamilton, M. T., Hamilton, D. G., & Zderic, T. W. (2007). Role of low energy expenditure
37 and sitting in obesity, metabolic syndrome, type 2 diabetes, and cardiovascular
38 disease. *Diabetes*, 56(11), 2655-2667. doi: db07-0882 [pii]
39
40
41

42 10.2337/db07-0882
43

44 Hamilton, M. T., Healy, G. N., Dunstan, D. W., Zderic, T. W., & Owen, N. (2008). Too little
45 exercise and too much sitting: Inactivity physiology and the need for new
46 recommendations on sedentary behaviour. *Current Cardiovascular Risk Reports*, 2(4),
47
48 292-298.
49
50
51

52 Haro, J. M., Arbabzadeh-Bouchez, S., Brugha, T. S., de Girolamo, G., Guyer, M. E., Jin, R., .
53
54 . . Kessler, R. C. (2006). Concordance of the Composite International Diagnostic
55
56
57
58
59

1 Interview Version 3.0 (CIDI 3.0) with standardized clinical assessments in the WHO
2 World Mental Health surveys. *Int J Methods Psychiatr Res*, 15(4), 167-180.
3

4 Harvey, J. A., Chastin, S. F., & Skelton, D. A. (2013). Prevalence of sedentary behavior in
5 older adults: a systematic review. *Int J Environ Res Public Health*, 10(12), 6645-6661.
6
7 doi: 10.3390/ijerph10126645
8
9

10
11
12 ijerph10126645 [pii]
13

14 Hu, F. B., Leitzmann, M. F., Stampfer, M. J., Colditz, G. A., Willett, W. C., & Rimm, E. B.
15
16 (2001). Physical activity and television watching in relation to risk for type 2 diabetes
17 mellitus in men. *Arch Intern Med*, 161(12), 1542-1548. doi: ioi00636 [pii]
18
19

20
21 Hu, F. B., Li, T. Y., Colditz, G. A., Willett, W. C., & Manson, J. E. (2003). Television
22 watching and other sedentary behaviors in relation to risk of obesity and type 2
23
24 diabetes mellitus in women. *JAMA*, 289(14), 1785-1791. doi:
25
26
27 10.1001/jama.289.14.1785
28
29

30
31 289/14/1785 [pii]
32
33

34 IPAQ-Group, T. (2005). Guidelines for Data Processing and Analysis of the International
35
36 Physical Activity Questionnaire (IPAQ).
37

38 Jakes, R. W., Day, N. E., Khaw, K. T., Luben, R., Oakes, S., Welch, A., . . . Wareham, N. J.
39
40 (2003). Television viewing and low participation in vigorous recreation are
41
42 independently associated with obesity and markers of cardiovascular disease risk:
43
44 EPIC-Norfolk population-based study. *Eur J Clin Nutr*, 57(9), 1089-1096. doi:
45
46
47 10.1038/sj.ejcn.1601648
48
49

50
51 1601648 [pii]
52
53

54 Katzmarzyk, P. T., Church, T. S., Craig, C. L., & Bouchard, C. (2009). Sitting time and
55
56 mortality from all causes, cardiovascular disease, and cancer. *Med Sci Sports Exerc*,
57
58 41(5), 998-1005. doi: 10.1249/MSS.0b013e3181930355
59
60

00005768-200905000-00005 [pii]

1
2 Kesse-Guyot, E., Charreire, H., Andreeva, V. A., Touvier, M., Hercberg, S., Galan, P., &
3
4 Oppert, J. M. (2012). Cross-sectional and longitudinal associations of different
5
6 sedentary behaviors with cognitive performance in older adults. *PLoS One*, 7(10),
7
8 e47831. doi: 10.1371/journal.pone.0047831
9
10

11 PONE-D-12-17909 [pii]

12
13 Kessler, R. C., Abelson, J., Demler, O., Escobar, J. I., Gibbon, M., Guyer, M. E., . . . Zheng,
14
15 H. (2004). Clinical calibration of DSM-IV diagnoses in the World Mental Health
16
17 (WMH) version of the World Health Organization (WHO) Composite International
18
19 Diagnostic Interview (WMHCIDI). *Int J Methods Psychiatr Res*, 13(2), 122-139.
20
21

22 Kikuchi, H., Inoue, S., Sugiyama, T., Owen, N., Oka, K., & Shimomitsu, T. (2013).
23
24 Correlates of prolonged television viewing time in older Japanese men and women.
25
26 *BMC Public Health*, 13, 213. doi: 10.1186/1471-2458-13-213
27
28

29 1471-2458-13-213 [pii]

30
31
32 Kronenberg, F., Pereira, M. A., Schmitz, M. K., Arnett, D. K., Evenson, K. R., Crapo, R. O., .
33
34 . . Hunt, S. C. (2000). Influence of leisure time physical activity and television
35
36 watching on atherosclerosis risk factors in the NHLBI Family Heart Study.
37
38 *Atherosclerosis*, 153(2), 433-443. doi: S0021915000004263 [pii]
39
40

41
42 Kuster, M. (2004). [Effects of sports and media consumption on the trunk muscle strength,
43
44 posture and flexibility of the spine in 12- to 14- year-old adolescents]. *Sportverletz*
45
46 *Sportschaden*, 18(2), 90-96. doi: 10.1055/s-2004-812910
47
48

49
50
51 Lindstrom, H. A., Fritsch, T., Petot, G., Smyth, K. A., Chen, C. H., Debanne, S. M., . . .
52
53 Friedland, R. P. (2005). The relationships between television viewing in midlife and
54
55 the development of Alzheimer's disease in a case-control study. *Brain Cogn*, 58(2),
56
57 157-165. doi: S0278-2626(04)00291-X [pii]
58
59
60
61
62
63
64
65

10.1016/j.bandc.2004.09.020

1
2 Lucas, M., Mekary, R., Pan, A., Mirzaei, F., O'Reilly, E. J., Willett, W. C., . . . Ascherio, A.
3
4 (2011). Relation between clinical depression risk and physical activity and time spent
5 watching television in older women: a 10-year prospective follow-up study. *Am J*
6
7 *Epidemiol*, 174(9), 1017-1027. doi: 10.1093/aje/kwr218
8
9
10

11
12 kwr218 [pii]

13
14 Marshall, S. J., Biddle, S. J., Gorely, T., Cameron, N., & Murdey, I. (2004). Relationships
15 between media use, body fatness and physical activity in children and youth: a meta-
16
17 analysis. *Int J Obes Relat Metab Disord*, 28(10), 1238-1246. doi:
18
19 10.1038/sj.ijo.0802706
20
21
22

23
24 0802706 [pii]

25
26 Nationmaster. (2009). Media statistics: Television viewing (most recent) bu country, from
27
28 http://www.nationmaster.com/graph/med_tel_vie-media-television-viewing
29
30

31
32 O'Connor, D. W., Pollitt, P. A., Treasure, F. P., Brook, C. P., & Reiss, B. B. (1989). The
33 influence of education, social class and sex on Mini-Mental State scores. *Psychol*
34
35 *Med*, 19(3), 771-776.
36
37

38
39 Otten, J. J., Littenberg, B., & Harvey-Berino, J. R. (2010). Relationship between self-report
40 and an objective measure of television-viewing time in adults. *Obesity (Silver*
41
42 *Spring)*, 18(6), 1273-1275. doi: 10.1038/oby.2009.371
43
44
45

46 oby2009371 [pii]

47
48 Paillard-Borg, S., Fratiglioni, L., Xu, W., Winblad, B., & Wang, H. X. (2012). An active
49 lifestyle postpones dementia onset by more than one year in very old adults. *J*
50
51 *Alzheimers Dis*, 31(4), 835-842. doi: 10.3233/JAD-2012-120724
52
53
54

55
56 L08604U20280N88X [pii]

- 1
2 Rundek, T., & Bennett, D. A. (2006). Cognitive leisure activities, but not watching TV, for
3 future brain benefits. *Neurology*, 66(6), 794-795. doi: 66/6/794 [pii]
4
5 10.1212/01.wnl.0000209497.38834.d7
6
7 Sanchez-Villegas, A., Ara, I., Guillen-Grima, F., Bes-Rastrollo, M., Varo-Cenarruzabeitia, J.
8
9 J., & Martinez-Gonzalez, M. A. (2008). Physical activity, sedentary index, and mental
10 disorders in the SUN cohort study. *Med Sci Sports Exerc*, 40(5), 827-834. doi:
11
12 10.1249/MSS.0b013e31816348b9
13
14
15
16 Sedentary-Behaviour-Research-Network. (2013). Letter to the Editor: Standardized use of the
17 terms “sedentary” and “sedentary behaviours”. *Mental Health and Physical Activity*,
18
19 6(1), 55-56. doi: <http://dx.doi.org/10.1016/j.mhpa.2012.06.001>
20
21
22
23
24 Sousa, R. M., Dewey, M. E., Acosta, D., Jotheeswaran, A. T., Castro-Costa, E., Ferri, C. P., .
25
26 . . Prince, M. J. (2010). Measuring disability across cultures--the psychometric
27 properties of the WHODAS II in older people from seven low- and middle-income
28 countries. The 10/66 Dementia Research Group population-based survey. *Int J*
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
- Sun, F., Norman, I. J., & While, A. E. (2013). Physical activity in older people: a systematic
review. *BMC Public Health*, 13, 449. doi: 10.1186/1471-2458-13-449
1471-2458-13-449 [pii]
- Teychenne, M., Ball, K., & Salmon, J. (2008). Physical activity and likelihood of depression
in adults: a review. *Prev Med*, 46(5), 397-411. doi: 10.1016/j.ypmed.2008.01.009
S0091-7435(08)00047-9 [pii]

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
- Teychenne, M., Ball, K., & Salmon, J. (2010). Sedentary behavior and depression among adults: a review. *Int J Behav Med*, 17(4), 246-254. doi: 10.1007/s12529-010-9075-z
- Tombaugh, T. N., & McIntyre, N. J. (1992). The mini-mental state examination: a comprehensive review. *J Am Geriatr Soc*, 40(9), 922-935.
- Wang, J. Y., Zhou, D. H., Li, J., Zhang, M., Deng, J., Tang, M., . . . Chen, M. (2006). Leisure activity and risk of cognitive impairment: the Chongqing aging study. *Neurology*, 66(6), 911-913. doi: 01.wnl.0000192165.99963.2a [pii]
10.1212/01.wnl.0000192165.99963.2a
- WHO. (2002). Towards a Common Language for Functioning, Disability and Health ICF. In <http://www.who.int/classifications/icf/en/> (Ed.).
- Wijndaele, K., Duvigneaud, N., Matton, L., Duquet, W., Delecluse, C., Thomis, M., . . . Philippaerts, R. M. (2009). Sedentary behaviour, physical activity and a continuous metabolic syndrome risk score in adults. *Eur J Clin Nutr*, 63(3), 421-429. doi: 1602944 [pii]
10.1038/sj.ejcn.1602944
- Wittchen, H. U. (1994). Reliability and validity studies of the WHO--Composite International Diagnostic Interview (CIDI): a critical review. *J Psychiatr Res*, 28(1), 57-84. doi: 0022-3956(94)90036-1 [pii]
- Wittchen, H. U., Robins, L. N., Cottler, L. B., Sartorius, N., Burke, J. D., & Regier, D. (1991). Cross-cultural feasibility, reliability and sources of variance of the Composite International Diagnostic Interview (CIDI). The Multicentre WHO/ADAMHA Field Trials. *Br J Psychiatry*, 159, 645-653, 658.
- Wittchen, H. U., Strehle, J., Gerschler, A., Volkert, J., Dehoust, M. C., Sehner, S., . . . Andreas, S. (2014). Measuring symptoms and diagnosing mental disorders in the

elderly community: the test-retest reliability of the CIDI65+. Int J Methods Psychiatr

Res. doi: 10.1002/mpr.1455

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Table 1
Description of socio-demographic variables and TV viewing across depression, anxiety and cognitive functioning

	Value	N	%	Controls (n=1040; 75.2%)	Depressive disorder only (n= 216; 13.4%)	Anxiety disorder only (n= 186; 15.6%)	Co morbid depressive and anxiety disorders (n= 59; 4.3%)	P-value	Mini- Mental score (S.D.)	No cognitive impairment (n= 1132, 81.9%)	Mild Cognitive Impairment (n=251, 18.1%)	P-value
	TOTAL	1383							27.66 (1.65)			
Gender	Female	658	47.6	41.6	61.8	66.1	74.6	< 0.001	27.57 (1.74)	46.0	54.6	< 0.05
	Male	725	52.4	58.4	38.2	33.9	25.4		27.74 (1.56)	54.0	45.4	
Age	Mean age (S.D.)	73.1 4	(5.69)	73.37 (.16)	72.46 (.43)	72.5 (.45)	71.49 (.63)	< 0.01		72.67 (5.61)	75.23 (5.57)	< 0.001
Level of education	High (> 13 yrs)	544	39.4	39.6	45.2	33.9	33.9	0.12	27.87 (1.36)	42.2	26.8	< 0.001
	Intermediate (9-12 yrs)	460	33.4	31.9	35.0	40.9	37.3		27.54 (1.72)	32.1	39.2	
	Low (< 8 yrs)	375	27.2	22.5	19.7	25.2	28.8		27.52 (1.90)	25.7	34.0	
Employment status	Unemployed /retired	1297	94.7	94.8	94.9	92.9	96.6	0.73	27.65 (1.65)	95.4	96.0	0.2
	Employed	72	5.3	5.2	5.1	7.1	3.4		27.72 (1.75)	5.5	4.0	
Financial situation	Very good	179	13.0	13.0	12.8	14.2	10.2	< 0.05	27.75 (1.38)	13.7	9.6	0.19
	Good	652	47.3	45.8	60.9	45.7	40.7		27.61 (1.59)	46.7	49.8	
	Just enough, low or very low	548	39.7	41.2	26.3	40.2	49.2		27.69 (1.80)	39.5	40.6	
Living status	Alone	403	29.2	26.9	35.7	34.9	39.7	< 0.05	27.50 (1.64)	28.0	34.8	< 0.05
	Not alone	977	70.8	73.1	64.3	65.1	60.3		27.73 (1.66)	72.0	65.2	
Center	Ferrara	486	35.1	37.6	17.8	36.2	35.6	< 0.001	28.10 (1.48)	37.9	22.7	< 0.001
	Hamburg	380	27.5	29.4	15.3	26.0	28.8		27.71 (1.60)	27.6	27.1	
	Geneva	517	37.4	33.0	66.9	37.8	35.6		27.22 (1.73)	34.5	50.2	
Physical activity	Low	208	15.0	14.2	13.4	21.3	20.3	0.09	27.20 (2.17)	13.4	22.3	< 0.001
	Moderate	589	42.6	41.8	51.0	39.4	40.7		27.71 (1.55)	43.3	39.4	
	High	586	42.4	43.9	35.7	39.4	39.0		27.78 (1.51)	43.3	38.2	
WHODAS II simple sum score	Mean (S.D.)	16.3 8	(5.67)	15.82 (5.22)	17.90 (7.03)	17.83 (6.13)	19.02 (6.14)	< 0.001		15.99 (5.21)	18.13 (7.14)	< 0.001
Medical problems past year	No	220	15.9	18.9	5.9	4.1	0.9	< 0.001		82.7	17.3	0.85
	Yes	1162	84.1	89.1	12.4	10.2	4.9			81.7	18.2	
Daily hours of TV watching	Mean (S.D.)	1.81	(0.94)	1.81 (0.9)	1.7 (1.0)	1.9 (0.9)	1.86 (0.9)	0.61		1.62 (.97)	1.91 (.95)	< 0.001
Daily hours sitting	Mean (S.D.)	2.01	(0.96)	2.0 (0.9)	2.1 (0.9)	1.99 (0.9)	2.07 (0.9)	0.64		2.07 (.94)	2.00 (0.96)	0.31
Total daily sedentary hours	Mean (S.D.)	3.7	(1.44)	3.67 (1.4)	3.6 (1.4)	3.7 (1.3)	3.7 (1.4)	0.89		3.62 (1.4)	3.98 (1.42)	< 0.001

Table 2

Regression analyses between TV viewing, Mini Mental Score Evaluation, Depressive and Anxiety diagnoses and covariates

Model	Standardised coefficients		Sig.	Confidence interval 95.0% for B	
	Beta	t		Lower limit	Upper limit
(Constant)		9.911	0.000	1439.281	2149.625
Years of education	-0.176	-6.652	0.000	-32.722	-17.817
Number of medical diagnoses	0.0147	5.586	0.000	18.732	39.011
Work status	-0.137	-5.246	0.000	-173.024	-78.834
Mini Mental Status - score	-0.105	-4.056	0.000	-38.755	-13.489
Physical activity levels	-0.064	-2.441	0.015	-.013	-0.001
Major depressive disorder (MDD) (year)	-0.061	-2.333	0.020	-135.983	-11.758

Regression statistics based on linear regression analyses (adjusted for gender, age, study centre, years of education, living status, level of functioning, number of medical diagnosis and physical activity). Corrected R²= 0.105

Graph 1

Distribution of days and hours of TV viewing in the sample

