

**TITLE PAGE:**

**Discovery and Structure-activity Relationships of Nociceptin Receptor Partial Agonists that Afford Symptom Ablation in Parkinson's Disease Models**

*Uma Gayathri Kamakolanu,<sup>1#</sup> Michael E. Meyer,<sup>1#</sup> Dennis Yasuda,<sup>1</sup> Willma E. Polgar,<sup>1</sup> Matteo Marti,<sup>2</sup> Daniela Mercatelli,<sup>3</sup> Clarissa Anna Pisanò,<sup>3</sup> Alberto Brugnoli,<sup>3</sup> Michele Morari,<sup>3</sup> Nurulain T. Zaveri<sup>1\*</sup>*

<sup>1</sup> Astraea Therapeutics, 320 Logue Avenue, Suite 142, Mountain View, CA 94043, USA

<sup>2</sup> Department of Morphology, Surgery and Experimental Medicine, Section of Legal Medicine, University of Ferrara, Ferrara (Italy)

<sup>3</sup> Department of Medical Sciences, Section of Pharmacology, University of Ferrara, Ferrara (Italy)

,

<sup>#</sup> Both authors contributed equally to this work.

**\* Address correspondence to:**

Dr. Nurulain T. Zaveri

Astraea Therapeutics,

320 Logue Avenue, Suite 142,

Mountain View, CA 94043, USA.

Tel: 650-254-0786

Email: [nurulain@astraeatherapeutics.com](mailto:nurulain@astraeatherapeutics.com)

## ABSTRACT

We report a novel series of C(3)-substituted piperdinylindoles as partial agonists at the nociceptin opioid peptide receptor (NOP), developed to explore a pharmacological hypothesis that NOP partial agonists would afford a dual pharmacological action of attenuating Parkinson's Disease (PD) motor symptoms and development of levodopa-induced dyskinesias (LID). Small-molecule and peptidic NOP antagonists have been shown to attenuate Parkinson's disease (PD)-like motor symptoms in animal models of PD; however, they may worsen dyskinesia development. On the other hand, NOP agonists were found to suppress the expression of levodopa-induced dyskinesias (LID), but had a narrow dose separation window vis-a-vis sedative effects characteristic of NOP agonists. Selective small-molecule NOP partial agonists or their SAR have not been systematically studied previously. We identified a novel class of piperidinyllindoles as selective NOP partial agonists and report here a detailed structure-activity relationship (SAR) study around the C(3) substituents of the indole moiety, designed to investigate the SAR of this scaffold with respect to NOP binding affinity, intrinsic activity and selectivity for the NOP receptor versus the other three opioid receptors, mu, delta and kappa. While the C-3 substituted indoles gave selective and high affinity NOP ligands, positively charged aminergic C-3 substituents significantly affected the intrinsic activity of the ligands, to afford partial agonists with a range of intrinsic efficacies. Selected NOP partial agonists were further characterized for their DMPK properties and brain penetration, and evaluated in the 6-OHDA hemilesioned rat model of PD. NOP partial agonists **1-3** with intrinsic efficacy between 25-35% were found to significantly attenuate parkinsonian motor deficits in hemilesioned rats. Further, unlike NOP antagonists, which appear to worsen dyskinesia expression, NOP partial agonists did not attenuate or worsen dyskinesia expression. These results suggest that NOP partial agonists

may be a viable alternative for a nondopaminergic approach for PD treatment. The SAR of the novel class of piperidinyl indole-based NOP partial agonists reported here will be useful for the future discovery of drug-like NOP partial agonists of varying efficacies.

## INTRODUCTION

Parkinson's Disease (PD) is the second most common neurodegenerative disorder after Alzheimer's disease affecting over 6 million individuals worldwide.(Global burden Lancet neurol ref 2018) PD is clinically characterized as having classic motor symptoms (hypo/akinesia, rigidity, gait disturbance and resting tremor), as well as non-motor complications such as depression and cognitive decline.(Schneider et al. Neurodegn Dis Mgt 2017) The dopamine (DA) precursor levodopa (L,3,4-dihydroxyphenylalanine; L-DOPA) is the cornerstone of PD therapy, often given in combination with COMT and MAO inhibitors to extend its bioavailability and therapeutic action (ref needed). However, chronic L-DOPA therapy is associated with the eventual appearance (within 10 years in ~80% of patients) of motor fluctuations and dyskinesias (involuntary movements) that limit its clinical effectiveness and greatly reduce the quality of life of patients (Obeso, *Trends Neurosci*, 2000).<sup>9</sup> The pathogenic hallmark of PD is the neurodegeneration of dopamine producing neurons in the substantia nigra pars compacta leading to striatal dopamine depletion, which leads to the classic motor features. On the other hand, long-term manifestations of PD treatment, such as the levodopa-induced dyskinesias (LID) are the result of an irreversible brain sensitization to L-DOPA. There is only one marketed antidyskinetic drug, amantadine, a glutamate antagonist (Goetz, *Movement Disorders*, 2002) which has poor and short-lasting clinical efficacy. Current strategies to minimize LID involve reducing the L-DOPA dosage and combining it with a dopaminomimetic (e.g. DA receptor agonists) to maintain symptomatic benefit. Thus, newly developed drugs that can effectively treat PD symptoms, and delay or even prevent the development of dyskinesia, would provide significant relief to PD patients, and fill in a gap of unmet medical need in PD therapy.

Previous studies by Morari and co-workers have demonstrated the involvement of the nociceptin/orphanin FQ peptide receptor (NOP, known previously as ORL-1 (opioid receptor-like-1 receptor)) in Parkinson's disease, particularly in development of motor symptoms. The NOP receptor, a G protein coupled receptor (GPCR) is the fourth member of the opioid receptor family having close homology to the classical opioid receptors (mu, delta, kappa) (Mollereau et al, *FEBS Lett*, 1994). The endogenous ligand for the NOP receptor is the heptadecapeptide nociceptin/orphanin FQ (N/OFQ), which is similar to the endogenous kappa opioid peptide dynorphin, but has no binding affinity to the three classic opioid receptors. NOP activation by endogenous N/OFQ is implicated in many physiological functions and pathologies, including pain and analgesia, anxiety, learning and memory, as well as modulation of tolerance development and reward.(Toll et al. 2018) The NOP receptor is found throughout the brain, specifically, in brain cortical and subcortical areas, particularly in striatum, globus pallidus and substantia nigra (SNr) neurons.(Mollereau and Mouledous, 2000) Morari et. al. have shown that endogenous N/OFQ contributes to the development of PD symptoms based on the following findings: 1) increased levels of N/OFQ found in the SNr following dopamine (DA) cell loss or impairment of DA transmission (Marti et al, *J Neurosci*, 2005) (Marti et al, *Mov Disord*, 2010), 2) NOP receptor antagonists reverse parkinsonian symptoms in neurodegenerative (6-OHDA hemilesioned rat, MPTP-treated mouse and macaque) and functional (reserpinized or haloperidol-treated animals) models of PD (Marti et al, *J Neurosci*, 2005) (Viaro et al, *Neurobiol Dis*, 2008) (Volta et al, *J Neurochem*, 2010), and 3) genetic deletion of the N/OFQ or NOP genes, or pharmacological blockade of the NOP receptor protects mice from the neurotoxic action of MPTP (Marti et al, *J Neurosci*, 2005)(Arcuri et al. *Neurobiol Dis* 2016). Mechanistic studies revealed that the antiparkinsonian action of NOP antagonists is accomplished through

normalization of the imbalance between excitatory (Glu) and inhibitory (GABA) inputs to the nigro-thalamic neurons, generated by striatal DA deafferentation (Marti, *J Neurosci*, 2007) (Marti et al, *J Neurochem*, 2008). This suggests that NOP antagonists may provide a non-dopaminergic approach for the symptomatic and neuroprotective therapy of PD.

Morari and colleagues also showed that NOP receptor agonists attenuate the expression of abnormal involuntary movements (AIMs, a rodent correlate of LID) in dyskinetic rats and nonhuman primates challenged with L-DOPA, by acting in the striatum where, contrary to SNr, the N/OFQ tone is reduced and NOP receptors are up-regulated following DA cell loss. (Marti et al, *J Neurosci*, 2012)(Arcuri, et al. *BJP* 2018) This action can be dissociated from the typical motor inhibiting effects of NOP agonists since antidyskinetic doses were 100-fold lower than doses that caused hypolocomotion. Morari et al showed that N/OFQ and NOP agonists specifically target dyskinesia pathways, preventing D1 receptor-mediated correlates of LID such as pERK phosphorylation and loss of depotentiation of synaptic plasticity in striatal GABA neurons (Marti et al. *J. Neurosci* 2012)(Arcuri 2018).

Based on the above evidence from NOP antagonists targeting PD symptoms by blocking nigral NOP receptors, and NOP agonists attenuating LID symptoms by stimulating up-regulated striatal NOP receptors, we hypothesized that NOP partial agonists could provide a balanced action in both areas affected in PD. A NOP partial agonist would be expected to function as an antagonist under conditions of high extracellular levels of endogenous N/OFQ (as in SNr) and alleviate parkinsonian motor dysfunction. On the other hand, in areas where the endogenous N/OFQ tone is low or absent and NOP receptors are upregulated (i.e. in striatum during LID development), NOP partial agonists would function as agonists. Based on this, we explored the discovery and structure-activity relationships of NOP partial agonists to identify suitable

compounds to validate our hypothesis that NOP partial agonists may be a viable approach for PD and LID treatment, and to gain further insights into the role of the NOP-N/OFQ system in the pathophysiology of PD and dyskinesias.


### **CHEMISTRY (Nur to Delete the AT-numbers at the very very end, not just yet)**

Zaveri et al previously disclosed NOP ligands with an indolinone motif (piperidinyl-indolinones) that provide both full and partial agonist efficacy and modest selectivity over the mu opioid receptor (MOP) (Zaveri, *J. Med. Chem.*, 2004) (Zaveri et al. AAPS Journal 2005) (Journigan, *Bioorg. Med. Chem.*, 2014). An SAR exploration was initiated with other similar bicyclic heterocycles such as an indole and indoline ring, in place of the indolinone, which resulted in two lead compounds **1** (AT-001) and **2** (AT-001), found to have partial agonist efficacy at NOP and >30-fold selectivity over the classical opioid receptors (Table 1). SAR studies to improve binding affinity and modulate the intrinsic activity of these lead compounds focused on the C(3) substituent of the indole ring.

The synthesis of lead compounds indoline **1** (AT-001) and indole **2** (AT-004) is shown in Scheme I. Commercially available indoline **I-1** and N-Boc-4-piperidone **I-2** were subjected to standard reductive animation to provide intermediate **I-3** in 96% yield. Subsequent Boc removal, followed by a reductive alkylation of the piperidine nitrogen with 4-*i*Pr-cyclohexanone initially provided indoline **1** (AT-001) as a mixture of ca. 1.5:1 *cis*:*trans* isomers (with respect to the disubstituted cyclohexyl ring). The diastereomers could be separated via column chromatography to provide indoline **1** as a single diastereomer in 28% yield over two steps. Oxidation of **1** (AT-001) with MnO<sub>2</sub> in dichloromethane smoothly provided indole **2** (AT-004) in 79% yield. Treating indole **2** (AT-004) with trimethylsilylisocyanate (TMSNCO) in MeCN provided C(3)-substituted amide **20** (AT-103) in modest yield. To explore the SAR of different

functionalities at the C(3) indole ring, aldehyde **I-4** was synthesized from indole **2** (AT-004) using Vilsmeier-Haack conditions (Scheme I). Conversion of aldehyde **I-4** to oxime **21** (AT-143) using standard conditions occurred readily in 96% yield. Oxime **21** was subsequently reduced using Ra(Ni) hydrogenation to afford amine **3** (AT-035) in 78% yield (Scheme I).

**Scheme I**




**Reagents and Conditions:** (a) AcOH, STAB, DCE, rt, 16 h, 96%; (b) i. TFA, CH<sub>2</sub>Cl<sub>2</sub>, 1h, ii. 4-*i*Pr-cyclohexanone, STAB, AcOH, DCE, rt, 1-2 days, 28% over 2 steps; (c) MnO<sub>2</sub>, CH<sub>2</sub>Cl<sub>2</sub>, rt, 16 h, 79%; (d) POCl<sub>3</sub>, DMF, 0 °C, 1 h, 74%; (e) NH<sub>2</sub>OH·HCl, NaOAc·3H<sub>2</sub>O, EtOH:H<sub>2</sub>O (2:1), 110 °C, 20 min, 96%; (f) H<sub>2</sub>(g) (55 psi), 100 wt% Ra-Ni, MeOH, 16 h, 78%; (g) ClSO<sub>2</sub>NCS, MeCN, 0 °C, 30 min, 37%.



Further functionalization of aldehyde **I-4** lead to various oxygen and nitrogen-containing substituents at the C(3) position of the indole ring (Scheme II). Aldehyde **I-4** was readily converted to alcohol **4** (AT-054) via NaBH<sub>4</sub> reduction, or to cyclic and acyclic amines (**5**, **8-13**) (AT-036, 037, 111, 360, 449, 454, 460)) via a reductive amination with the appropriate amine.

### Scheme II




**Reagents and Conditions:** (a) amine, AcOH, STAB, DCE or MeOH, rt, 1-2 days; (b) NaBH<sub>4</sub>, EtOH, rt, 2 h, 86%; (c) H<sub>2</sub>(g) (55 psi), 100 wt% Ra-Ni, MeOH, 16 h, 10-15%; (d) TFA, CH<sub>2</sub>Cl<sub>2</sub>, 0→rt, 20 min, 78%.

The synthesis of amides **14**, **18**, and **19** (AT-122, 129, and 173), were accomplished in good yield from amine **3** (AT-035) via EDCI coupling with the corresponding N-Cbz-protected amino acid, followed by removal of the Cbz protecting group (Scheme III). The use of the Cbz

protecting group was critical in synthesizing amides **18** and **19** (AT-129 and AT-173), as the Boc-protected substrate degraded upon exposure to acidic conditions during deprotection.


### Scheme III



**Reagents and Conditions:** (a) HOBT, EDCI, TEA, N-Cbz amino acid, CH<sub>2</sub>Cl<sub>2</sub>, 0 °C, 67-88%; (b) H<sub>2</sub> balloon, 10 wt% Pd/C (10%), THF or EtOH, 3 h, 41-88%.

To explore longer alkyl substitution and greater flexibility at the indole C(3) position, the route was amended so that the indole ring was constructed using a Larock heteroannulation reaction between a terminally silylated alkyne and a N-substituted iodoaniline (Scheme IV) (Larock, *J. Am. Chem. Soc.*, 1991). The N-Boc protected amino alkynes **IV-3** and **IV-4** were obtained from the corresponding alkynyl alcohols (**IV-1** and **IV-2**, respectively) in four straightforward steps: 1) the alcohol was converted into a tosylate group, 2) S<sub>N</sub>2 displacement of the tosylate group with phthalimide, 3) removal of the phthalimide group using hydrazine, and 4) Boc-protection of the amine.

### Scheme IV



**Reagents and Conditions:** (a) TsCl, TEA, cat. DMAP,  $CH_2Cl_2$ , rt, 16 h, 93-99%; (b) phthalimide,  $K_2CO_3$ , DMF, 50 °C, 6 h, 67-94%; (c)  $NH_2NH_2$ , MeOH, rt, 16 h, 59-70%; (d)  $(Boc)_2O$ , cat. DMAP, THF, rt, 85-88%; (e) AcOH, STAB, DCE, rt, 16 h; (f) TFA,  $CH_2Cl_2$ , 0 °C, 1 h; (g) 4-*i*Pr-cyclohexanone, AcOH, STAB, DCE, rt, 1-2 days, 39% over 3 steps; (h) alkyne **IV-3** or **IV-4**, cat.  $Pd(OAc)_2$ , LiCl,  $K_2CO_3$ , DMF, 102 °C, 2.5 h, 65-76%; (i) AcCl, MeOH, 0 °C, 1 h, then EtOAc, 78-80%; (j) i. BzNCS,  $CH_2Cl_2$ , 0 °C, ii.  $K_2CO_3$ , MeOH, rt, 41-50%.

The synthesis of the requisite iodoaniline **IV-6** and the eventual formation of the indole ring is shown in Scheme IV. Commercially available 2-iodoaniline **IV-5** was converted to iodoaniline intermediate **IV-6** in a 3-step sequence consisting of 1) reductive amination with *N*-Boc-4-piperidone, 2) removal of the Boc group, and 3) reductive amination of the piperidine nitrogen with 4-isopropylcyclohexanone in 39% yield over the three steps. At this stage, the key Larock heteroannulation reaction between iodoaniline **IV-6**, and either alkyne **IV-3** or **IV-4**, proceeded in good yield to provide indole intermediates **IV-7** and **IV-8**, respectively. When indoles **IV-7** and **IV-8** were separately exposed to AcCl/MeOH conditions, clean removal of

both the silyl group and the Boc moiety occurred to provide amines **6** and **7** (AT-147 and 160) *in situ* as bis HCl salts (Scheme IV), and the salts could be isolated pure by simple filtration. Finally, methyl-, ethyl-, and propyl-linked amines **3** (AT-035), **6** (AT-160), and **7** (AT-147) respectively, reacted with trimethylsilyl thioisocyanate (TMSNCS) to give the corresponding methyl-, ethyl-, and propyl-linked thioureas **15** (AT-111), **16** (AT-163), and **17** (AT-155), in consistent yields (Scheme IV).

### ***In Vitro* Binding and Functional Efficacy**

The compounds were tested *in vitro* for binding affinity at the four opioid receptor subtypes, in radioligand displacement assays in membranes from human NOP, MOP, KOP and DOP receptor-transfected chinese hamster ovary (CHO) cells, using methods we have described in detail previously (Journigan et al. 2014 Bioorg Med Chem., Daga et al. JCIM 2014). Binding affinity was calculated as the binding constant  $K_i$  (nM), as shown in Table 1. The intrinsic activity (functional efficacy) was determined using the [<sup>35</sup>S] GTP $\gamma$ S binding assay in membranes prepared from the opioid receptor-transfected CHO cells, using methods we have previously published (Toll, et al. JPET 2009, Journigan et al. 2014 Bioorg Med Chem., Daga et al. JCIM 2014). The intrinsic activities of the partial agonists in the GTP $\gamma$ S binding assay shown in Table 1 are reported as the % stimulation compared to that of the full agonist N/OFQ taken as 100%. The potencies of the functional efficacy are reported as the  $EC_{50}$  (Table 1). All compounds showed weak binding affinity for the  $\delta$  opioid receptor (DOP) and  $\kappa$  opioid receptor (KOP), and were not further tested for functional efficacy at these receptors.

## **RESULTS and DISCUSSION**

Compounds **3-21** were designed to explore the SAR of the indole moiety on the N-substituted piperidine scaffold, which is a common motif in most NOP ligands.(Zaveri, 2016 JMC, Zaveri and Meyer, 2019) The piperidinyl-indoles have not be widely explored for their SAR for NOP binding affinity, opioid selectivity or agonist efficacy. Further, to the best of our knowledge, there is only one other report claiming a NOP partial agonist, having a 1,3,8-triazaspirodecanone scaffold and NOP binding IC<sub>50</sub> of 34 nM.(Ross et al 2015) However, no functional efficacy data was reported for this compound from this group.(Battista 2009) To explore the SAR of the indolylpiperidines, we focused on the C-3 substituent and maintained the N-isopropylcyclohexylpiperidinyl moiety in all the analogs synthesized, as this piperidinyl substituent has afforded high NOP binding affinity in other scaffolds we have investigated.(Zaveri AAPS 2005, Zaveri, Bifunctional I, Journigan Bifunctional 2)

The C(3)-unsubstituted lead compounds indoline **1** (AT-001) and indole **2** (AT-004) show nanomolar affinity for NOP (10 nM), and weak affinity for the three classical opioid receptors (Table 1, entries 1 and 2), giving at least >40-fold selectivity for the NOP receptor. Both compounds show partial agonist efficacy at NOP in the [<sup>35</sup>S]GTPγS functional assay. Indoline **1** (AT-001) showed a three-fold higher potency (EC<sub>50</sub>) as an agonist at NOP, than indole **2** (AT-004). From the C(3)-substituted indoles bearing a polar substituent on a single methylene linker (compounds **3-5**), primary amine **3** (AT-035) has high binding affinity (NOP Ki = 3 nM), ~20-fold selectivity versus MOP, and modest potency as a NOP partial agonist. Alcohol **4** (AT-054) and dimethylamino (tertiary amine substituent) **5** have 10-fold lower affinity than primary amine **3**, although alcohol **4** (AT-054) has comparable intrinsic activity to amine **3** whereas tertiary dimethylamine **5** showed significantly reduced agonist potency. On the other hand, tertiary amine **8**, a pyrrolidinomethylene at C(3), showed high binding affinity for NOP

( $K_i$  1.47 nM), and >200-fold selectivity versus the other opioid receptors, and also had high intrinsic activity, to a near full agonist (Table 1). We performed a chain-length SAR involving amine **3** (AT-035), and the ethyl-, and propyl-linked amines **6** (AT-160) and **7** (AT-147), respectively. Both methyl- (**3**) (AT-035), and ethyl-linked (**6**) (AT-160) amines are optimal for NOP affinity, and the increase in chain length to the ethyl-linked amine **6** (AT-160) also results in a 10-fold higher NOP selectivity compared to methyleneamine **3** (AT-035). Functionally, the two compounds are both partial agonists at NOP, with similar potency. However, a further extension of the chain length to the propylamine **7** was not optimal for NOP binding.

Reducing the freedom of rotation of the primary amine C(3) substituent by using a 4-amino-substituted piperidine moiety (as in compound **9**) resulted in a significant drop in affinity compared to the open-chain analog **7**, although the 4-hydroxypiperidine analog **10** showed higher binding affinity. Interestingly however, 2-hydroxypyrrolidine **11** (AT-449) showed even higher NOP binding affinity than hydroxypiperidine **10** (AT-454), and an increase in intrinsic activity. It appears that the pyrrolidine **8** (AT-360) and hydroxypyrrolidine **11** (AT-449) show higher agonist efficacy than the open-chain primary amines **3**, **6** and **7**, suggesting that these compounds may be interacting with different groups at the NOP receptor and/or binding in different orientations, affecting the activation of the receptor. Further contraction of the hydroxypyrrolidine ring of **11** to azetidine **12** (AT-460), retains modest binding affinity at NOP ( $K_i = 23$  nM), but shows very low agonist potency in the [ $^{35}$ S]GTP $\gamma$ S functional assay ( $EC_{50} = 1559$  nM). In addition, acyclic alkyl amine **13** (AT-036), an acyclic variant of hydroxypyrrolidine **11** (AT-449) showed weak binding affinity toward all four opioid receptors. The SAR comparison of compounds **3-7** to **8** and **11** illustrates the significance of the pyrrolidine moiety at the C-3 of the piperidinyloles for high NOP affinity and full agonism.

Thiourea analogs **15-17** were synthesized to investigate the importance of a basic amine moiety at the C-3 substituent of compounds **3**, **6** and **7**. The thiourea analogs **15-17** showed only a modest (2-3-fold) drop in NOP binding affinity and agonist potency, suggesting that it is likely hydrogen bonding or polar interactions of the C-3 substituent are important for high affinity rather than a charged interaction. This is further consistent with the reasonable affinity of the ethylaminomethylamide analog **14**.

Amides at the C-3 substituent containing an arginine unit (as in **18**) (AT-129) or a lysine unit (as in **19**) (AT-173), were found to have modest NOP affinities (11–23 nM), suggesting that a charged interaction of the guanidinium group in **18** or the amino group in **19** is likely not an important contributor to the binding affinity. Further, both these compounds had poor agonist potencies and showed low to no intrinsic activity.

Compound **20** (AT-103), where the benzylic carbon bound to C(3) is sp<sup>2</sup> hybridized in the form of an amide, showed weak binding affinity for all the opioid receptors (K<sub>i</sub> > 100 nM). However, the C(3)-oxime **21** (AT-143) is a high-affinity (K<sub>i</sub> = 2 nM), NOP-selective (NOP/MOP = ca. 25 fold) potent full agonist. Further SAR development and understanding of NOP binding is needed to explain the observed NOP full agonism for compounds **8** and **21**. Overall, our SAR showed that the C(3)-substituted piperidinyloles afforded selective NOP partial agonists of varying efficacies, ranging from 20% to nearly full agonist efficacy (>80%).

### **In Vitro ADME**

Selected compounds were evaluated *in vitro* in ADME assays to characterize intestinal permeability, blood brain barrier permeability, and rat liver microsomal stability (Table 2). Indoline **1** (AT-001) and indole **2** (AT-004), show high permeability and low efflux in the Caco-2 intestinal permeability assay as well as in the MDR1-MDCK assay for BBB permeability

(Table 2). Amine **3** (AT-035) showed lower permeability in the Caco-2 assay and high efflux in the MDR1-MDCK BBB permeability assay. However, amine **3** has excellent stability in rat liver microsomes, whereas indole **2** has a short half-life in rat liver microsomes. From the in vitro microsomal stability of the amine **3** (AT-035) compared to that of unsubstituted indole **2** (AT-004), it appears that the unsubstituted C(3)-position in indole **2** (AT-004) may be a site of metabolic liability, and that the high permeability of indole **2** (AT-004), which arises from its high lipophilicity, comes at the expense of solubility and metabolism.

### ***In vivo* efficacy assessment of NOP partial agonists in rat models of Parkinson's disease and levodopa-induced dyskinesia**

NOP receptor partial agonists **1**, **2** and **3** were evaluated in vivo for their efficacy in improving parkinsonian motor disabilities in a rodent neurodegenerative model of PD, i.e. the 6-hydroxydopamine-induced hemilesioned rat model, that we have extensively used to characterize NOP antagonists.(Marti et al, JNS 2005, 2007; Marti et al., JNC 2008) The compounds were administered intraperitoneally (i.p.) in a range of doses.

Figure 1 show that NOP partial agonists **1** (AT-001), **2** (AT-004) and **3** (AT-035) attenuate parkinsonian disabilities in 6-OHDA hemilesioned rats. Parkinsonian motor deficits were assessed using three different measures of motor activity (Marti et al, JNS 2004, 2005, 2007; Marti et al., JNC 2008). Acute administration of **2** (AT-004) (0.1-3 mg/kg, i.p.) improved motor deficits, partially reversing (~25%) the increase of immobility time (a measure of akinesia) at the contralateral (parkinsonian) in the bar test (Figure 1 panel D) and doubling stepping activity at the contralateral paw (a measure of akinesia/bradykinesia) in the drag test (Figure 1 panel E). Moreover, it enhanced by ~20%, the time spent on the rotarod (a measure of overall gait ability) (Figure 1 panel F). This profile is predictive of antiparkinsonian activity



since it is replicated by L-DOPA (Marti, JNS 2005, 2007) and by NOP antagonists (Viaro 2008, Marti, 2007, 2008). Compound **1** (AT-001) and **3** also showed similar patterns, improving parkinsonian motor deficits, although with some differences in efficacy in the three tests, at doses of 0.01-1 mg/kg (Figure 1, panels A-C and G-I), suggesting that the NOP partial agonists may actually be counteracting endogenous N/OFQ action at the NOP receptor, as predicted by our working hypothesis.


However, in the same dose ranges, all three compounds had no effect on the ‘expression’ of LID in fully dyskinetic rats challenged with L-DOPA, using methods previously reported by us (Marti, J. Neurosci 2012; Arcuri BJP, 2018) (data not shown). It is possible that the partial agonist efficacy of about 25-35% in all three compounds is too low to elicit an anti-dyskinetic effect seen with NOP full agonists that we have previously reported (Marti et al, J. Neurosci 2012; Arcuri et al. BJP 2018). SAR studies to obtain higher efficacy NOP partial agonists from the piperidinyl-indole and other novel scaffolds are underway to address this hypothesis and will be reported in due course.

## CONCLUSIONS

In summary, we report the discovery and SAR of a novel series of 4-piperidinyl-1*H*-indoles as NOP receptor partial agonists. The SAR studies reported here represent the first systematic SAR study around NOP partial agonists and a novel NOP ligand scaffold, the piperidinyl-indoles. SAR analysis showed that the nature of the C(3) substituent of the indole moiety influences the binding affinity and intrinsic activity of these ligands. Furthermore, the C(3)-substituted piperidinyl indoles showed significantly lower affinity for the classical opioid receptors, resulting in selective NOP partial agonists. Selected NOP partial agonists **1**, **2** and **3** also reduced parkinsonian motor symptoms in the 6-OHDA-hemilesioned rat model but were


ineffective in reducing LID symptoms at the same dose range. Together with our previous studies, these results provide further evidence that the NOP receptor is a possible non-dopaminergic target for the symptomatic treatment of PD and that NOP receptor partial agonists are promising for development of new symptomatic PD therapies. The novel NOP partial agonists and their SAR reported here can support lead optimization efforts in this regard.

**Table 1. Binding affinities and functional efficacies of C(3)-substituted piperidinyloles at human opioid receptors**




		Binding Affinities, $K_i$ (nM) <sup>a</sup>				[S] <sup>35</sup> GTPγS Functional Assay (nM) <sup>b</sup>			
Compound number		NOP	MOP	DOP	KOP	NOP		MOP	
						EC <sub>50</sub>	% Stim	EC <sub>50</sub>	% Stim
1		10.3 ± 0.03	604 ± 5	3070 ± 747	562 ± 150	54.5 ± 17.6	33.0 ± 2.8	ND	—
	<b>R</b>								
2	H	9.80 ± 0.86	376 ± 37	923 ± 413	1594 ± 57	160 ± 64	29.3 ± 9.5	232 ± 12.6 <sup>c</sup>	16.3 ± 7.6 <sup>c</sup>
3		3.27 ± 0.30	65.3 ± 2.4	>10K	1737 ± 172	121 ± 51.7	35.9 ± 5.7	410 ± 105	11.8 ± 2.7
4		44.7 ± 9.1	716 ± 21	>10K	>10K	117 ± 13.7	18.9 ± 0.70	ND	—
5		43.8 ± 7.4	504 ± 173	>10K	>10K	532 ± 185	28.1 ± 2.9	ND	—
6		2.27 ± 0.11	437 ± 118	>10K	2750 ± 383	139 ± 30.3	27.8 ± 7.2	ND	—
7		20.9 ± 7.6	46.2 ± 6.1	>10K	1320 ± 411	249 ± 9	18.6 ± 2.8	1764 ± 118	15.7 ± 3.2
8		1.47 ± 0.57	453 ± 107	>10K	461 ± 135	97.4 ± 7.4	87.9 ± 11.8	ND	—
9		216 ± 12	1308 ± 18	>10K	3456 ± 300	ND	—	ND	—

**Table 1. Binding affinities and functional efficacies of C(3)-substituted piperidinyloles at human opioid receptors**



		Binding Affinities, $K_i$ (nM) <sup>a</sup>				[S] <sup>35</sup> GTP-γS Functional Assay (nM) <sup>b</sup>			
Compound number		NOP	MOP	DOP	KOP	NOP		MOP	
						EC <sub>50</sub>	% Stim	EC <sub>50</sub>	% Stim
10		64.0 ± 21.3	>10K	6028 ± 652	3840 ± 1887	446 ± 130	10.6 ± 1.2	ND	—
11		9.46 ± 1.01	306 ± 77	>10K	>10K	575 ± 157	69.9 ± 12.1	ND	—
12		23.6 ± 3.1	>10K	>10K	>10K	1559 ± 398	34.3 ± 9.3	ND	—
13		136 ± 7.50	973 ± 211	>10K	1490 ± 52	ND	ND	ND	—
14		15.9 ± 3.7	269 ± 12	>10K	1440 ± 342	277 ± 144	15.9 ± 4.9	ND	—
15		10.5 ± 1.25	71.3 ± 16.2	4067 ± 68	1328 ± 288	343 ± 18	51.2 ± 9.7	>10K	9.8 ± 2.1

**Table 1. Binding affinities and functional efficacies of C(3)-substituted piperidinyloindoles at human opioid receptors**



		Binding Affinities, $K_i$ (nM) <sup>a</sup>				[S] <sup>35</sup> GTP $\square$ S Functional Assay (nM) <sup>b</sup>			
Compound number		NOP	MOP	DOP	KOP	NOP		MOP	
						EC <sub>50</sub>	% Stim	EC <sub>50</sub>	% Stim
16		24.4 ± 0.30	227 ± 50.0	>10K	81.0 ± 2.8 <sup>b</sup>	>10K	5.6 ± 3.0	ND	—
17		31.8 ± 1.7	382 ± 129	>10K	830 ± 181	175 ± 23	34.5 ± 3.6	ND	—
18		11.0 ± 2.7	857 ± 109	>10K	674 ± 85	>10K	11.1 ± 1.5	ND	—
19		23.1 ± 8.5	949 ± 74	>10K	>10K	464 ± 124	37.7 ± 8.9	ND	—
20		129 ± 41.4	453 ± 8	2712 ± 300	446 ± 24	ND	ND	ND	—
21		2.17 ± 0.35	56.4 ± 1.7	1690 ± 870	>10K	67.0 ± 11.0	85.9 ± 9.8	3199 ± 260	37.5 ± 2.9


a) Binding affinities were determined using radioligand displacement assays performed in membranes of CHO cells stably expressing the human NOP, MOP, KOP and DOP receptors, and their respective ra-diolioligands, [<sup>3</sup>H]N/OFQ–NOP, [<sup>3</sup>H]U69,593–KOP, [<sup>3</sup>H]DAMGO–MOP and [<sup>3</sup>H]DPDPE–DOP receptor. Equilibrium dissociation constants ( $K_i$ ) were derived from  $IC_{50}$  values using the Cheng–Prusoff equation. Each  $K_i$  value represents the arithmetic mean  $\pm$  SEM from at least three independent experiments, each performed in triplicate. ND = not determined.

b) Compounds with  $K_i$  values  $>100$ nM were not tested in functional assays (N.D.). At the DOP and KOP receptor, all compounds showed binding affinity  $K_i >100$ nM, hence functional efficacy at KOP and DOP not determined.

**Table 2. *In Vitro* ADME data for indoles 1-3**

Compd	Caco-2 Bidirectional <sup>a</sup>			MDR1-MDCK Permeability <sup>b</sup>			Human Liver Microsomal Stability (t <sub>1/2</sub> min)	Rat Liver Microsomal Stability (t <sub>1/2</sub> min)
	P <sub>app</sub> (x 10 <sup>-6</sup> cm/s)		Efflux Ratio (Perm/Efflux)	P <sub>app</sub> (x 10 <sup>-6</sup> cm/s)		Efflux Ratio (Perm/Efflux)		
	A-B	B-A		A-B	B-A			
<b>1</b>	30	25	0.8 (Hi/No)	26.8	33.8	1.3 (Hi/No)	ND	ND
<b>2</b>	24	19.1	0.8 (Hi/No)	28.8	15.2	0.5 (Hi/No)	5.3	5.8
<b>3</b>	4.64	4.92	1.1 (Hi/No)	0.34	14.4	43 (Low/Hi)	>60	>60

(Need Legend or methods Nur to do)



**Figure 1. Effect of NOP partial agonists 1 (AT-001), 2 (AT-004) and 3 (AT-035) on parkinsonian disabilities in the 6-OHDA hemilesioned rat model of PD. Acute systemic administration of 1-3 in a range of doses (panels A-C, 1, AT-001, 0.001-3 mg/kg, i.p., n=17-23**



mice per group) (panels D-F, 2, AT-004, 0.1-3 mg/kg, i.p., n=6-12 mice per group) or (panels G-I, 3, AT-035, 0.01-3 mg/kg, i.p.; n=6-7 mice per group) attenuated motor deficits in three behavioural measures, (A) the bar test (*left graphs*), (B) the drag test (*middle graphs*) and (C) rotarod test (*right graphs*). Data are means  $\pm$  SEM of 6-7 (AT-001 and AT-035) or 17-23 (AT-004) determinations per group. Essential statistical values: treatment effect panels A-C ( $F_{5,2}=4.60$ ,  $p=0.0007$ ,  $F_{5,2}=4.52$ ,  $p=0.0008$ ,  $F_{5,1}=13.72$ ,  $p<0.0001$ , respectively), panels D-F ( $F_{3,2}=4.39$ ,  $p=0.0051$ ,  $F_{3,2}=8.45$ ,  $p<0.0001$ ,  $F_{3,1}=7.59$ ,  $p<0.0001$ , respectively), panels G-I ( $F_{4,2}=14.31$ ,  $p<0.0001$ ,  $F_{4,2}=3.79$ ,  $p=0.0071$ ,  $F_{4,1}=11.51$ ,  $p<0.0001$ , respectively) \* $P < 0.05$ , \*\* $P < 0.01$  different from vehicle (saline + 5% Tween 80 and 1% DMSO).

## EXPERIMENTAL SECTION

### CHEMISTRY METHODS

Thin layer chromatography was performed on Analtech silica gel GF 250 micron TLC plates. The plates were visualized with a 254 nm UV light and iodine. Flash chromatography was carried out on F60 silica gel, 43-60  $\mu$ m (230-400 mesh), 60  $\text{\AA}$  (Silicycle SiliaFlash). All solvents and chemicals were purchased from commercial suppliers and used without further purification. All reactions were capped from the atmosphere unless otherwise stated. NMR was recorded on a Varian Mercury Plus NMR (300 MHz) using  $\text{CDCl}_3$  (7.27 ppm standard) or  $\text{DMSO-}d_6$  (2.50 ppm standard). Data for  $^1\text{H}$  NMR were recorded as follows:  $\delta$  chemical shift (ppm), multiplicity (s, singlet; d, doublet; t, triplet; dd, doublet of doublets; dt, doublet of triplets; q, quartet; dq, doublet of quartets; m, multiplet; br, broad; etc.), coupling constant (Hz), integration. Mass spectra were obtained on a LCQ Fleet Ion Trap LCMS, a micromass ZMD 1000 or PE Sciex API 150EX LCMS using electrospray ionization (ESI), or APCI mode. Elemental analyses were performed by Atlantic Microlabs, Norcross, GA. HRMS analyses were performed by the Mass Spectrometry Service Laboratory, University of Minnesota Department of Chemistry, Minneapolis, MN on a Bruker BioTOF II HRMS using ESI mode. HPLC analysis was performed on a reverse phase Agilent Zorbax SB-Phenyl column (5  $\mu$ m, 2.1 x150 mm), using a binary gradient of 95:5 $\rightarrow$ 5:95 solvent A (95/5  $\text{H}_2\text{O}/\text{ACN}$  + 0.1% formic acid): solvent B (5/95  $\text{H}_2\text{O}/\text{ACN}$  + 0.1% formic acid) for 12 minutes, at a flow rate of 0.40 mL/min. Eluted peaks were monitored at 254 nm with a Shimadzu SPD-10AVP UV-Vis detector.

### SYNTHESIS

**General Procedure A: Reductive Amination.** The aniline or amine substrate (1.00-2.00 equiv) and carbonyl compound (1.00-1.50 equiv) were charged into a round bottom flask. 1,2-

DCE (0.030-0.25M) was added, and the mixture was stirred until both components dissolved. NOTE: If amine was a salt, *i*Pr<sub>2</sub>NEt (2.30 equiv) was added at this stage to freebase. To this solution was added glacial AcOH (1.00-2.30 equiv) at ambient temperature, and the solution was stirred for 60 minutes. At this stage, sodium triacetoxyborohydride (STAB) (1.50-3.00 equiv) was added. The reaction was allowed to stir at room temperature and monitored by TLC (EtOAc:Hexanes). After 1-2 days, the reaction was  $\geq 90\%$  complete by TLC analysis. The reaction was quenched with saturated NaHCO<sub>3</sub> (aq.) and stirred until the reaction mixture was basic and bubbling had ceased. The biphasic layer was separated, and the organic layer was washed 2x with H<sub>2</sub>O, brine, dried with MgSO<sub>4</sub>, filtered and concentrated in vacuo to provide a crude product that was purified via flash chromatography to provide the desired product.

**General Procedure B: Boc Removal (Step 1) and Reductive Amination with 4-isopropylcyclohexanone (Step 2).** *Step 1.* A solution of N-Boc intermediate (1.00 equiv) in CH<sub>2</sub>Cl<sub>2</sub> (0.25-0.30M) was cooled to 0 °C, and then TFA (6-30 equiv) was added over several minutes. Upon completion of addition, the ice-bath was removed and the reaction was allowed to warm to room temperature and monitored by TLC (EtOAc:Hexanes). After 2 hours, the reaction was complete. The reaction was concentrated in vacuo, followed by the addition of EtOAc, which was consequently removed in vacuo to ensure the majority of excess TFA was removed. The resulting oily residue was dissolved in EtOAc and was stirred as saturated NaHCO<sub>3</sub> (aq.) was added until the aqueous layer remained basic. The layers were separated, and the aqueous layer was extracted with EtOAc until UV activity in the aqueous layer was minimal (3-8x). The EtOAc layers were combined, washed with brine, dried with MgSO<sub>4</sub>, filtered, and concentrated in vacuo to provide the piperidine intermediate.

*Step 2.* The piperidine intermediate from the previous step (1.00 equiv) and 4-*i*Pr-cyclohexanone (1.00-1.50 equiv) were dissolved in 1,2-DCE (0.070M). To the reaction was added glacial AcOH (1.00-2.30 equiv), and the reaction was stirred for 30 minutes. After 30 minutes, STAB (1.50-2.30 equiv) was added. An Ar balloon was fitted on top of the reaction, and the reaction was monitored by TLC (MeOH:CH<sub>2</sub>Cl<sub>2</sub>:NH<sub>4</sub>OH (aq.)). After 2-3 days, the reaction was  $\geq$ 95% complete. Saturated NaHCO<sub>3</sub> (aq.) was added until the aqueous layer remained basic. At this stage, the layers were separated, and the aqueous layer was extracted 2x with CH<sub>2</sub>Cl<sub>2</sub>. The organic layers were combined, and washed 2x with H<sub>2</sub>O, brine, dried with MgSO<sub>4</sub>, filtered, and concentrated in vacuo to provide a crude residue that was purified via flash chromatography using EtOAc:Hexanes:NH<sub>4</sub>OH (aq.).

**General Procedure C: Formation of thiourea from amine.** The amine (1.00 equiv) was dissolved in CH<sub>2</sub>Cl<sub>2</sub> (0.10 M) at room temperature, and then the reaction was placed in an ice-bath and cooled to 0 °C. Benzoyl isothiocyanate (1.05-1.10 equiv) was added, and after several minutes of stirring at 0 °C, the reaction was allowed to warm to room temperature and monitored by TLC (EtOAc:Hexanes). After 1-3 hours, the reaction was complete by TLC. The reaction was concentrated in vacuo, the remaining residue was dissolved in MeOH (0.050 M), and to this was added K<sub>2</sub>CO<sub>3</sub> (7.00 equiv). The reaction was stirred at room temperature and monitored by TLC (EtOAc:Hexanes). Once the reaction was complete, it was filtered over a pad of Celite, rinsed with MeOH, and then concentrated in vacuo to provide a crude material that was purified via flash chromatography using EtOAc:Hexanes:NH<sub>4</sub>OH (aq.) to provide the desired thiourea product.

**tert-butyl 4-(indolin-1-yl)piperidine-1-carboxylate (I-3).** See General Procedure A: Indoline **I-1** (10.0 g, 83.9 mmol, 1.00 equiv), N-Boc piperidone **I-2** (17.6 g, 88.1 mmol, 1.05

equiv), AcOH, 4.80 mL, 83.9 mmol, 1.00 equiv), STAB (26.7 g, 12.6 mmol, 1.50 equiv), DCE (352 mL, 0.25M). The crude oil was purified via flash chromatography using 10:90 EtOAc:Hexanes to provide indoline **I-3** as a white solid (24.3 g, 96% yield). <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 7.06 (t, *J* = 6.0 Hz, 2H), 6.03 (t, *J* = 6.0 Hz, 1H), 6.43 (d, *J* = 6.0 Hz, 1H), 4.25 (m, 2H), 3.52 (m, 1H), 3.35 (t, *J* = 6.3 Hz, 2H), 2.79 (m, 2H), 1.80 (d, *J* = 9.3 Hz, 2H), 1.60 (m, 4H), 1.49 (s, 9H); MS (APCI) *m/z*: 303.06 [M+H]<sup>+</sup>.

**cis-1-(1-(4-isopropylcyclohexyl)piperidin-4-yl)indoline (1, AT-001)**. See General Procedure B: Step 1 (Boc Removal): Indoline **I-3** (24.4 g, 80.5 mmol, 1.00 equiv), TFA (38.0 mL, 496 mmol, 6.20 equiv), CH<sub>2</sub>Cl<sub>2</sub> (300 mL, 0.27M). NOTE: Combined EtOAc layers were dried immediately with MgSO<sub>4</sub>, and were not washed with water or brine, as the product is water-soluble. The product was obtained as a grey solid (13.6 g, 84% yield); Step 2 (Reductive Amination): N-H piperidine from the previous step (13.6 g, 67.2 mmol, 1.00 equiv), *i*Pr-cyclohexanone (9.40 g, 67.2 mmol, 1.00 equiv), AcOH (3.85 mL, 67.2 mmol, 1.00 equiv), STAB (21.3 g, 101 mmol, 1.50 equiv), DCE (960 mL, 0.070M). Purified via flash chromatography using 10:90:1.5 EtOAc:Hexanes:NH<sub>4</sub>OH (aq.) to provide compound **1** (AT-001) as a light-gold oil (33% yield). R<sub>f</sub> = 0.25 (20:80:3 drops EtOAc:Hexanes:NH<sub>4</sub>OH (aq.)), UV, pAA, I<sub>2</sub>; <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 7.05 (t, *J* = 5.7 Hz, 2H), 6.60 (t, *J* = 5.7 Hz, 1H), 6.41 (d, *J* = 5.7 Hz, 1H), 3.37 (m, 3H), 3.10 (d, *J* = 8.7 Hz, 2H), 2.94 (t, *J* = 6.3 Hz, 2H), 2.27 (m, 1H), 2.14 (t, *J* = 8.7 Hz, 2H), 1.82-1.54 (m, 11H), 1.38 (m, 2H), 1.13 (m, 1H), 0.88 (d, *J* = 5.1 Hz, 6H); MS(ESI) *m/z*: 327.4 [M+H]<sup>+</sup>. The HCl salt of amine **1** was formed by dissolving the amine in CH<sub>2</sub>Cl<sub>2</sub> (0.10M) at room temperature, and cooling the reaction in a 0 °C ice-bath. To this solution was added HCl (2.0M in Et<sub>2</sub>O, 1.50 equiv), and the resulting mixture was allowed to warm to room temperature and stir for 10 minutes. At this time, the reaction was concentrated in vacuo, Et<sub>2</sub>O was added,

and the mixture was concentrated in vacuo to provide a brown solid (repeat 3x total). This solid was then triturated w/ ca. 5-10% MeOH:Et<sub>2</sub>O to provide the HCl salt of compound **1** as a cream solid. Anal. Calcd for C<sub>22</sub>H<sub>34</sub>N<sub>2</sub>·1.00 HCl·0.10 H<sub>2</sub>O: C, 72.44; H, 9.73; N, 7.68; found: C, 72.39; H, 9.79; N, 7.34.

**cis-1-(1-(4-isopropylcyclohexyl)piperidin-4-yl)-1H-indole (2, AT-004).** Indoline **1** (AT-001) (4.63 g, 14.2 mmol, 1.00 equiv) was dissolved in 180 mL of CH<sub>2</sub>Cl<sub>2</sub>. To this solution was added 4ÅMS (56.8 g, 4g/mmol of indoline), followed by MnO<sub>2</sub> (12.3 g, 142 mmol, 10.0 equiv) and another 20 mL of CH<sub>2</sub>Cl<sub>2</sub>. A balloon filled with argon was fitted on the reaction, and the thick suspension was stirred and monitored by TLC (20:80:3 drops EtOAc:Hexanes:NH<sub>4</sub>OH (aq.)). After 16 hours, the reaction was complete. The mixture was filtered over a large pad of Celite and the remaining solid was washed 5x with CH<sub>2</sub>Cl<sub>2</sub>. The filtrate was concentrated in vacuo to provide a crude oil. This material was dissolved in EtOAc, and 10% HCl (aq.) was added with vigorous stirring, which resulted in a white precipitate. The white solid was filtered, and washed 3x with EtOAc, and was then air-dried over 1 hour. The white solid was then suspended in EtOAc, 70% NaHCO<sub>3</sub> (aq.) was added, and the mixture stirred until >90% of the solid had dissolved (usually overnight). The EtOAc layer was separated, washed with H<sub>2</sub>O, brine, dried with MgSO<sub>4</sub>, filtered, and concentrated in vacuo to provide a thick oil that was purified via flash chromatography using 10:90:1.5 EtOAc:Hexanes:NH<sub>4</sub>OH (aq.) to provide indole **2** (AT-004) as an off-white solid (3.65 g, 79% yield). R<sub>f</sub> = 0.25 (10:90:3 drops EtOAc:Hexanes:NH<sub>4</sub>OH (aq.)), UV, pAA, I<sub>2</sub>; <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 7.64 (d, *J* = 6.0 Hz, 1H), 7.39 (d, *J* = 6.0 Hz, 1H), 7.26 (m, 1H), 7.20 (t, *J* = 6.0 Hz, 1H), 7.11 (t, *J* = 6.0 Hz, 1H), 6.52 (d, *J* = 2.4 Hz, 1H), 4.23 (m, 1H), 3.20 (d, *J* = 9.0 Hz, 2H), 2.30 (m, 3H), 2.08 (m, 4H), 1.51-1.78 (m, 7H), 1.40 (m, 2H), 1.17 (m, 1H), 0.90 (d, *J* = 4.8 Hz, 6H); MS(ESI) *m/z*: 325.4

[M+H]<sup>+</sup>. Anal. Calcd for C<sub>22</sub>H<sub>32</sub>N<sub>2</sub>: C, 81.43; H, 9.94; N, 8.63; found: C, 81.54; H, 9.87; N, 8.45.

**cis-1-(1-(4-isopropylcyclohexyl)piperidin-4-yl)-1H-indole-3-carbaldehyde (I-4)**. To a stirred solution of 25.0 mL DMF at 0 °C was added POCl<sub>3</sub> (3.66 mL, 40.0 mmol, 4.00 equiv). The solution was stirred at 0 °C for 15 minutes. At this stage, indole **2** (3.10 g, 10.0 mmol, 1.00 equiv) was dissolved in 10 mL of DMF with the assistance of heat. The warm solution of indole **2** (AT-004) was then added to the reaction, and the reaction was rinsed with 5.00 mL of DMF. The addition of indole **2** leads to a red solution, and the reaction was allowed to stir for 15-20 minutes at 0 °C. TLC (50:50:3 drops EtOAc:Hexanes:NH<sub>4</sub>OH (aq.)) showed the reaction was complete. The reaction was poured into a saturated NaHCO<sub>3</sub> (aq.) icebath, followed by the addition of CH<sub>2</sub>Cl<sub>2</sub>. The mixture was stirred vigorously for 30 minutes, upon which the layers were separated, and the aqueous layer was extracted with CH<sub>2</sub>Cl<sub>2</sub> until UV activity was minimal (5-6x). The organic layer was then washed 3x with H<sub>2</sub>O, brine, dried with MgSO<sub>4</sub>, filtered, and concentrated in vacuo to provide a dark red oil, which was purified via flash chromatography using 50:50:1.5 EtOAc:Hexanes:NH<sub>4</sub>OH (aq.) to provide aldehyde **I-4** as a light-yellow solid (2.15 g, 74% yield). R<sub>f</sub> = 0.20 (50:50:3 drops EtOAc:Hexanes:NH<sub>4</sub>OH (aq.)), UV, I<sub>2</sub>; <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 10.0 (s, 1H), 8.33 (m, 1H), 7.89 (s, 1H), 7.43 (m, 1H), 7.33 (m, 2H), 4.29 (m, 1H), 3.28 (d, *J* = 7.8 Hz, 2H), 2.40 (m, 3H), 2.19 (m, 3H), 1.55-1.78 (m, 8H), 1.42 (m, 2H), 1.17 (m, 1H), 0.90 (d, *J* = 5.7 Hz, 6H); MS(ESI) *m/z*: 353.1 [M+H]<sup>+</sup>.

**cis-1-(1-(4-isopropylcyclohexyl)piperidin-4-yl)-1H-indole-3-carbaldehyde oxime (21, AT-143)**. Aldehyde **I-4** (2.15 g, 6.10 mmol, 1.00 equiv), NH<sub>2</sub>OH·HCl (551 mg, 7.93 mmol, 1.30 equiv), and NaOAc·3H<sub>2</sub>O (1.08 g, 7.93 mmol, 1.30 equiv) were charged into a round bottom flask. Absolute EtOH (20.5 mL) and 10 mL of H<sub>2</sub>O were added, and the reaction was fitted with

a condenser and an Ar balloon on top. The suspension was heated to reflux (ca. 110 °C oil bath) and monitored by TLC (40:60:3drops EtOAc:Hexanes:NH<sub>4</sub>OH (aq.)). After 2 hours, the reaction was complete. The reaction was allowed to cool to room temperature upon which a white precipitate formed. The mixture was diluted with EtOAc and saturated NaHCO<sub>3</sub> (aq.), and stirred until the mixture became a biphasic solution. The layers were separated, and the organic layer was washed 2x with H<sub>2</sub>O, brine, dried with MgSO<sub>4</sub>, filtered, and concentrated in vacuo to provide oxime **21** (AT-143) as a white solid (2.13 g, 96% yield). The two isomers of the oxime are in ca. 3:2 ratio, and can be enriched via flash chromatography using 40:60:1□50:50:1 EtOAc:Hexanes:NH<sub>4</sub>OH (aq.). R<sub>f</sub> = 0.40 (top spot), 0.45 (bottom spot) (50:50:3 drops EtOAc:Hexanes:NH<sub>4</sub>OH (aq.)), UV, I<sub>2</sub>); Isomer #1: <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 10.8 (br s, 1H), 8.47 (s, 1H), 7.77 (m, 1H), 7.41 (d, *J* = 6.0 Hz, 1H), 7.28-7.20 (m, 3H), 4.31 (m, 1H), 3.30 (d, *J* = 8.7 Hz, 2H), 2.55 (m, 1H), 2.46 (t, *J* = 8.4 Hz, 2H), 2.22 (m, 3H), 1.86 (m, 2H), 1.78-1.57 (m, 6H), 1.43 (m, 2H), 1.18 (m, 1H), 0.91 (d, *J* = 5.1 Hz, 6H); Isomer #2: <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 8.30 (s, 1H), 8.07 (d, *J* = 6.0 Hz, 1H), 7.48 (s, 1H), 7.40 (d, *J* = 6.0 Hz, 1H), 7.28 (m, 1H), 7.20 (t, *J* = 5.4 Hz, 1H), 4.23 (m, 1H), 3.21 (d, *J* = 8.7 Hz, 2H), 2.35 (m, 3H), 2.13 (m, 4H), 1.80-1.55 (m, 7H), 1.43 (m, 2H), 1.18 (m, 1H), 0.90 (d, *J* = 5.1 Hz, 6H); MS(ESI) *m/z*: 368.5 [M+H]<sup>+</sup>. Anal. Calcd for C<sub>23</sub>H<sub>33</sub>N<sub>3</sub>O: C, 75.16; H, 9.05; N, 11.43; found: C, 75.27; H, 8.88; N, 11.30.

**cis-(1-(1-(4-isopropylcyclohexyl)piperidin-4-yl)-1H-indol-3-yl)methanamine (3, AT-035)**. In a 500 mL parr bottle was added oxime **21** (2.00 g, 5.44 mmol, 1.00 equiv), and it was then suspended in MeOH (46 mL). To this mixture was added Raney Nickel (2.00 g, 100 wt%, prewashed with MeOH) and concentrated NH<sub>4</sub>OH (aq.) (23 mL). The reaction mixture was placed on the parr hydrogenator, and the parr vessel was pressurized with H<sub>2</sub> (g) at ca. 20 psi,



then purged under vacuo. This was repeated a total of 3x. The vessel was the pressurized at 50 psi for 18 hours. The reaction mixture was filtered over Celite, and washed thoroughly with MeOH. The resulting filtrate was concentrated in vacuo, and the crude product was purified by flash chromatography using 8:92 MeOH:CH<sub>2</sub>Cl<sub>2</sub> to provide amine **3** as a white solid (1.52 g, 78% yield).  $R_f = 0.30$  (10:90:3 drops MeOH:CH<sub>2</sub>Cl<sub>2</sub>:NH<sub>4</sub>OH (aq.), UV, I<sub>2</sub>). <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>):  $\delta$  7.58 (d,  $J = 6.0$  Hz, 1H), 7.44 (d,  $J = 6.3$  Hz, 1H), 7.37 (s, 1H), 7.10 (t,  $J = 5.7$  Hz, 1H), 6.98 (t,  $J = 5.7$  Hz, 1H), 4.26 (m, 1H), 3.86 (s, 1H), 3.16 (s, 2H), 3.08 (d,  $J = 8.7$  Hz, 3H), 2.27 (br s, 1H), 2.18 (t,  $J = 8.4$  Hz, 2H), 1.91-1.86 (m, 4H), 1.71 (br s, 2H), 1.55-1.50 (m, 3H), 1.38 (dd,  $J = 18.3, 9.3$  Hz, 4H), 1.10 (s, 1H), 0.86 (d,  $J = 5.1$  Hz, 6H); MS(ESI)  $m/z$ : 354.5 [M+H]<sup>+</sup>. The bis HCl salt of amine **3** was formed by dissolving the amine in CH<sub>2</sub>Cl<sub>2</sub> (0.10M) at room temperature, followed by the addition of HCl (2.0M in Et<sub>2</sub>O, 2.20 equiv). The reaction was allowed to stir for 10 minutes. At this time, the reaction was concentrated in vacuo, Et<sub>2</sub>O was added to the mixture, followed by concentrating the mixture in vacuo (repeat 3x total) to provide the HCl salt of amine **3**. Anal. Calcd for C<sub>23</sub>H<sub>35</sub>N<sub>3</sub>·2.00 HCl·1.50 H<sub>2</sub>O: C, 60.92; H, 8.89; N, 9.27; found: C, 60.94; H, 8.76; N, 9.11.

**cis-1-(1-(4-isopropylcyclohexyl)piperidin-4-yl)-1H-indole-3-carboxamide (20, AT-103)**. Indole **2** (AT-004) (60.0 mg, 0.185 mmol, 1.00 equiv) was dissolved in 1.00 mL of MeCN (0.19M) with the aid of heat. The solution was allowed to cool to room temperature, and then it was placed in an ice bath for several minutes. Chlorosulfonylisocyanate (20.1  $\mu$ L, 0.231 mmol, 1.25 equiv) was then added, turning the reaction light-yellow. The reaction was stirred at 0 °C and monitored by TLC (20:80 EtOAc:Hexanes). After 30 minutes, TLC showed full consumption of indole **2**. Water was added to the reaction, temporarily resulting in a white precipitate. The ice-bath was removed, and the mixture was allowed to warm to room

temperature and stir overnight, resulting in a light-yellow solution. The reaction was then quenched with saturated NaHCO<sub>3</sub> (aq.), and diluted with EtOAc. The layers were separated, and the aqueous layer was extracted 2x with EtOAc. The EtOAc layers were combined, washed with brine, dried with MgSO<sub>4</sub>, filtered, and concentrated in vacuo to provide the crude product. Amide **20** was purified via flash chromatography using 2:96:1 MeOH:EtOAc:NH<sub>4</sub>OH (aq.) to provide amide **20** as a white solid (25 mg, 37% yield). R<sub>f</sub> = 0.30 (5:95:3 drops MeOH:EtOAc:NH<sub>4</sub>OH (aq.), UV, I<sub>2</sub>); <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 7.99 (m, 1H), 7.90 (s, 1H), 7.46-7.43 (m, 1H), 7.30-7.27 (m, 2H), 5.78 (br s, 2H), 4.23 (m, 1H), 3.22 (d, *J* = 8.7 Hz, 2H), 2.37 (m, 2H), 2.28 (t, *J* = 8.7 Hz, 2H), 2.1 (m, 4H), 1.75-1.52 (m, 7H), 1.40 (m, 2H), 1.16 (m, 1H), 0.90 (d, *J* = 5.1 Hz, 6H); MS(ESI) *m/z*: 368.5 [M+H]<sup>+</sup>. Anal. Calcd for C<sub>23</sub>H<sub>33</sub>N<sub>3</sub>O·0.50 MeOH: C, 73.59; H, 9.20; N, 10.96; found: C, 73.64; H, 8.82; N, 10.57.

**cis-1-(1-(1-(4-isopropylcyclohexyl)piperidin-4-yl)-1H-indol-3-yl)-N,N-dimethylmethanamine (5, AT-171).** In the reduction of of oxime **21** (AT-143) to amine **3** (AT-035), a non-polar impurity formed. Purification via flash chromatography using 3:97:1 MeOH:CH<sub>2</sub>Cl<sub>2</sub>:NH<sub>4</sub>OH (aq.) provided the byproduct in low yield (10-15%) as a white solid, and the structure was elucidated as dimethyl amine **5**. R<sub>f</sub> = 0.70 (10:90:3 drops MeOH:CH<sub>2</sub>Cl<sub>2</sub>:NH<sub>4</sub>OH (aq.), UV, I<sub>2</sub>); <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 7.73 (d, *J* = 6.0 Hz, 1H), 7.62 (d, *J* = 6.0 Hz, 1H), 7.37 (d, *J* = 6.0 Hz, 1H), 7.22 (d, *J* = 5.1 Hz, 1H), 7.16 (m, 1H), 5.61-5.43 (m, 6H), 4.87 (s, 1H), 4.65 (d, *J* = 3.9 Hz, 1H), 4.59 (d, *J* = 3.6 Hz, 1H), 4.19 (t, *J* = 5.4 Hz, 1H), 3.21 (d, *J* = 7.5 Hz, 2H), 2.38-2.30 (m, 3H), 2.18-1.98 (m, 6H), 1.71-1.60 (m, 5H), 1.43-1.37 (m, 2H), 1.14 (t, *J* = 3.3 Hz, 1H), 0.89 (d, *J* = 5.1 Hz, 6H); MS(ESI) *m/z*: 382.27 [M+H]<sup>+</sup>. HRMS Calcd for C<sub>25</sub>H<sub>39</sub>N<sub>3</sub> (M+H)<sup>+</sup> 382.3144, found 382.2866.

**cis-1-(1-(4-isopropylcyclohexyl)piperidin-4-yl)-3-(pyrrolidin-1-ylmethyl)-1H-indole** (**8**, AT-360). See General Procedure A. Conditions: aldehyde **I-4** (150 mg, 0.426 mmol, 1.00 equiv.), pyrrolidine (57.0  $\mu$ L, 0.682 mmol, 1.60 equiv), glacial AcOH (56.1  $\mu$ L, 0.980 mmol, 2.30 equiv), STAB (208 mg, 0.980 mmol, 2.30 equiv), DCE (4.30 mL, 0.10M). The crude product was purified via flash chromatography using 0:100:1 $\rightarrow$ 2:98:1 MeOH:EtOAc:NH<sub>4</sub>OH (aq.) to provide indole **8** (AT-360) as an oil (93 mg, 53% yield).  $R_f$  = 0.10 (100:3 drops EtOAc:NH<sub>4</sub>OH (aq.), UV, I<sub>2</sub>). <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  7.69 (d,  $J$  = 7.8 Hz, 1H), 7.36 (d,  $J$  = 8.1 Hz, 1H), 7.24 (s, 1H), 7.20 (t,  $J$  = 7.5 Hz, 1H), 7.11 (t,  $J$  = 7.5 Hz, 1H), 4.19 (m, 1H), 3.84 (s, 2H), 3.18 (d,  $J$  = 11.7 Hz, 2H), 2.60 (m, 4H), 2.35-2.19 (m, 3H), 2.12-1.97 (m, 4H), 1.81-1.48 (m, 11H), 1.40 (m, 2H), 1.16 (m, 1H), 0.91 (d,  $J$  = 6.6 Hz, 6H); MS(ESI)  $m/z$ : 408.6 [M+H]<sup>+</sup>. The HCl salt of amine **8** was formed by dissolving the amine in CH<sub>2</sub>Cl<sub>2</sub> (0.10M) at room temperature, and cooling the reaction in a 0 °C ice-bath. To this solution was added HCl (2.0M in Et<sub>2</sub>O, 3.00 equiv), and the reaction was allowed to warm to room temperature and stir for 30 minutes. At this time, the reaction was concentrated in vacuo, Et<sub>2</sub>O was added, and the mixture was concentrated in vacuo (repeat 3x total) to provide an off-white solid. Anal. Calcd for C<sub>27</sub>H<sub>41</sub>N<sub>3</sub>·2.00 HCl·1.50 H<sub>2</sub>O: C, 63.89; H, 9.14; N, 8.28; found: C, 63.92; H, 8.92; N, 8.26.

**cis-1-((1-(1-(4-isopropylcyclohexyl)piperidin-4-yl)-1H-indol-3-yl)methyl)piperidin-4-amine** (**9**, AT-037). *Step 1*. See General Procedure A. Conditions: aldehyde **I-4** (100 mg, 0.284 mmol, 1.00 equiv.), 4-(N-Boc-amino)piperidine (56.0 mg, 0.284 mmol, 1.00 equiv), Note: glacial AcOH not used, STAB (180 mg, 0.851 mmol, 3.00 equiv), DCE (10.0 mL, 0.03M). The crude Boc-protected amine (**II-1**), (0.153 g, 98% yield) was taken directly into the Boc removal step.

*Step 2.* Boc-protected amine **II-1** (85.0 mg, 0.160 mmol, 1.00 equiv) was dissolved in CH<sub>2</sub>Cl<sub>2</sub> (5.0 mL, 0.030M), followed by the addition of excess TFA (4.00 mL). The reaction mixture was stirred for 10 minutes, and then the solvent was evaporated to provide a yellow oily product. This was triturated with Et<sub>2</sub>O to provide the TFA salt of indole **9** as a solid (97.5 mg, 78% yield). <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 7.70 (dd, *J* = 4.2, 2.4 Hz, 2H), 7.53 (dd, *J* = 4.2, 2.4 Hz, 2H), 7.40 (br s, 1H), 4.22 (m, 3H), 3.48 (dd, *J* = 10.5, 5.1 Hz, 4H), 2.08 (s, 14H), 1.69 (m, 2H), 1.43 (m, 3H), 1.35-1.29 (m, 4H), 1.20 (t, *J* = 5.1 Hz, 3H), 0.94-0.88 (m, 6H); MS(ESI) *m/z*: 437.50 [M+H]<sup>+</sup>. Anal. Calcd for C<sub>28</sub>H<sub>44</sub>N<sub>4</sub>·3.00 TFA·1.00 Et<sub>2</sub>O·1.00 CH<sub>2</sub>Cl<sub>2</sub>: C, 50.25; H, 6.22; N, 5.77; found: C, 50.26; H, 6.01; N, 5.60.

**cis-1-((1-(1-(4-isopropylcyclohexyl)piperidin-4-yl)-1H-indol-3-yl)methyl)piperidin-4-ol (10, AT-454).** See General Procedure A. Conditions: aldehyde **I-4** (205 mg, 0.630 mmol, 1.00 equiv.), piperin-4-ol·HCl (173 mg, 1.26 mmol, 2.00 equiv), *i*Pr<sub>2</sub>NEt (253 μL, 1.45 mmol, 2.30 equiv), glacial AcOH (83.0 μL, 1.45 mmol, 2.30 equiv), STAB (307 mg, 1.45 mmol, 2.30 equiv), DCE (6.30 mL, 0.10M). The crude product was purified via flash chromatography using 3:97:1 MeOH:CH<sub>2</sub>Cl<sub>2</sub>:NH<sub>4</sub>OH (aq.) to afford indole **10** as a light-yellow foam (140 mg, 51% yield). *R<sub>f</sub>* = 0.15 (5:95:1 MeOH:CH<sub>2</sub>Cl<sub>2</sub>:NH<sub>4</sub>OH (aq.)), UV, I<sub>2</sub>). <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 7.71 (d, *J* = 7.8 Hz, 1H), 7.37 (d, *J* = 8.1 Hz, 1H), 7.24-7.17 (m, 2H), 7.11 (t, *J* = 6.9 Hz, 1H), 4.42 (m, 1H), 3.75 (m, 3H), 3.19 (d, *J* = 12.0 Hz, 2H), 2.85 (m, 2H), 2.38-2.20 (m, 4H), 2.18-1.85 (m, 4H), 1.78-1.35 (m, 14H), 1.17 (m, 1H), 0.91 (d, *J* = 6.6 Hz, 6H); MS(ESI) *m/z*: 438.7 [M+H]<sup>+</sup>. The bis tartrate salt of amine **10** was formed by dissolving the amine in MeOH (0.10M) at room temperature. To the reaction was added L-(+)-tartaric acid (2.00 equiv) in a MeOH solution. The reaction was allowed to stir for 30 minutes. At this time, the reaction was concentrated in vacuo, and then Et<sub>2</sub>O was added to the mixture, followed by concentrating the

mixture in vacuo (repeat 3x total) to provide a light-yellow solid. Anal. Calcd for  $C_{28}H_{43}N_3O \cdot 2.00$  tartaric acid  $\cdot 1.50$   $H_2O$ : C, 56.53; H, 7.64; N, 5.49; found: C, 56.91; H, 7.83; N, 5.12.

**cis-(R)-1-((1-(1-(4-isopropylcyclohexyl)piperidin-4-yl)-1H-indol-3-yl)methyl)pyrrolidin-3-ol (11, AT-449).** See General Procedure A. Conditions: aldehyde **I-4** (97.0 mg, 0.275 mmol, 1.00 equiv.), pyrrolidin-3-ol (47.9 mg, 0.550 mmol, 2.00 equiv), glacial AcOH (36.2  $\mu$ L, 0.633 mmol, 2.30 equiv), STAB (134 mg, 0.633 mmol, 2.30 equiv), DCE (2.75 mL, 0.10M). The crude product was purified via flash chromatography using 5:95:1 MeOH:CH<sub>2</sub>Cl<sub>2</sub>:NH<sub>4</sub>OH (aq.) to provide indole **11** as a light-yellow foam (112 mg, 97% yield).  $R_f = 0.20$  (5:95:3 drops MeOH:CH<sub>2</sub>Cl<sub>2</sub>:NH<sub>4</sub>OH (aq.), UV, I<sub>2</sub>). <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  7.69 (d,  $J = 6.0$  Hz, 1H), 7.36 (d,  $J = 6.0$  Hz, 1H), 7.21 (m, 2H), 7.11 (t,  $J = 5.1$  Hz, 1H), 4.32 (m, 1H), 4.18 (m, 1H), 3.86 (s, 2H), 3.18 (d,  $J = 9.0$  Hz, 2H), 2.93 (m, 1H), 2.74 (m, 1H), 2.60 (m, 1H), 2.43-1.98 (m, 10H), 1.78-1.52 (m, 8H), 1.57 (m, 2H), 1.15 (m, 1H), 0.90 (d,  $J = 5.1$  Hz, 6H); MS(ESI)  $m/z$ : 424.6 [M+H]<sup>+</sup>. The bis tartrate salt of amine **11** was formed by dissolving the amine in MeOH (0.10M) at room temperature. To the reaction was added L-(+)-tartaric acid (2.50 equiv) in a MeOH solution. The reaction was allowed to stir for 30 minutes. At this time, the reaction was concentrated in vacuo, and then Et<sub>2</sub>O was added to the mixture, followed by concentrating the mixture in vacuo (repeat 3x total) to provide a tan solid. Anal. Calcd for  $C_{27}H_{41}N_3O \cdot 2.50$  tartaric acid  $\cdot 2.00$   $H_2O$ : C, 53.23; H, 7.24; N, 5.03; found: C, 53.60; H, 6.96; N, 4.95.

**cis-1-((1-(1-(4-isopropylcyclohexyl)piperidin-4-yl)-1H-indol-3-yl)methyl)azetid-3-ol (12, AT-460).** See General Procedure A. Conditions: aldehyde **I-4** (40 mg, 0.113 mmol, 1.00 equiv.), azetid-3-ol  $\cdot$  HCl (24.8 mg, 0.226 mmol, 2.00 equiv), *i*Pr<sub>2</sub>NEt (45.3  $\mu$ L, 0.260 mmol,

2.30 equiv), glacial AcOH (14.9  $\mu$ L, 0.260 mmol, 2.30 equiv), STAB (55.1 mg, 0.260 mmol, 2.30 equiv), DCE (1.10 mL, 0.10M). The crude product was purified via flash chromatography using 5:95:1 MeOH:CH<sub>2</sub>Cl<sub>2</sub>:NH<sub>4</sub>OH (aq.) to provide indole **12** (AT-460) as a light-golden oil (37.4 mg, 81% yield).  $R_f$  = 0.15 (5:95:1 MeOH:CH<sub>2</sub>Cl<sub>2</sub>:NH<sub>4</sub>OH (aq.), UV, I<sub>2</sub>). <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  7.65 (d,  $J$  = 5.7 Hz, 1H), 7.58 (br, 1H), 7.36 (d,  $J$  = 6.3 Hz, 1H), 7.21 (m, 2H), 7.12 (t,  $J$  = 5.4 Hz, 1H), 4.44 (p,  $J$  = 4.2 Hz, 1H), 4.19 (m, 1H), 3.84 (s, 2H), 3.69 (m, 2H), 3.18 (d,  $J$  = 9.0 Hz, 2H), 3.07 (m, 2H), 2.36-2.22 (m, 3H), 2.10-1.88 (m, 4H), 1.78-1.46 (m, 7H), 1.40 (m, 2H), 1.14 (m, 1H), 0.90 (d,  $J$  = 5.1 Hz, 6H); MS(ESI)  $m/z$ : 410.6 [M+H]<sup>+</sup>. The bis tartrate salt of amine **12** was formed by dissolving the amine in MeOH:CH<sub>2</sub>Cl<sub>2</sub> (3:2, 0.10M) at room temperature. To the reaction was added L-(+)-tartaric acid (2.00 equiv) in a MeOH solution. The reaction was allowed to stir for 30 minutes. At this time, the reaction was concentrated in vacuo, and then Et<sub>2</sub>O was added to the mixture, followed by concentrating the mixture in vacuo (repeat 3x total) to provide an off-white solid. Anal. Calcd for C<sub>26</sub>H<sub>39</sub>N<sub>3</sub>O·2.00 tartaric acid·2.00 H<sub>2</sub>O: C, 54.75; H, 7.43; N, 5.63; found: C, 54.92; H, 7.27; N, 5.35.

**cis-2-(((1-(1-(4-isopropylcyclohexyl)piperidin-4-yl)-1H-indol-3-yl)methyl)amino)ethan-1-ol (13, AT-036).** See General Procedure A. Conditions: aldehyde **I-4** (100 mg, 0.284 mmol, 1.00 equiv), ethanolamine (17.0  $\mu$ L, 0.284 mmol, 1.00 equiv), Note: glacial AcOH not used, STAB (180 mg, 0.851 mmol, 3.00 equiv), DCE (6.00 mL, 0.050M). The crude product was purified via flash chromatography using 8:92:1 MeOH:CH<sub>2</sub>Cl<sub>2</sub>:NH<sub>4</sub>OH (aq.) to provide indole **13** as a liquid (25 mg, 22% yield).  $R_f$  = 0.30 (10:90 MeOH:CH<sub>2</sub>Cl<sub>2</sub>, UV, I<sub>2</sub>); <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  7.56 (d,  $J$  = 5.7 Hz, 1H), 7.36 (d,  $J$  = 6.3 Hz, 1H), 7.18 (t,  $J$  = 5.1 Hz, 2H), 7.06 (d,  $J$  = 5.7 Hz, 1H), 4.21 (m, 1H), 3.66 (br s, 1H), 3.22 (s, 1H), 2.79 (s, 1H), 2.32 (br s, 3H), 2.10 (s, 4H), 1.72-1.61 (m, 7H), 1.43-1.32 (m, 2H), 1.26 (s, 2H), 1.15 (s, 1H), 0.89 (d,  $J$  =

4.8 Hz, 6H); MS(ESI)  $m/z$ : 398.24 [M+H]<sup>+</sup>. HRMS Calcd for C<sub>25</sub>H<sub>39</sub>N<sub>3</sub>O (M+H)<sup>+</sup> 398.3093, found 398.3173.

**cis-(1-(1-(4-isopropylcyclohexyl)piperidin-4-yl)-1H-indol-3-yl)methanol (4, AT-054).** Aldehyde **I-4** (1.50 g, 4.26 mmol, 1.00 equiv) was taken up in absolute EtOH (50 mL, 0.085M), and sodium borohydride (806 mg, 21.3 mmol, 5.00 equiv) was then added in several portions at room temperature. After bubbling had subsided, the reaction was stirred for an additional 30 minutes and then concentrated in vacuo. The resulting white residue was suspended in a 5% NaOH (aq.) solution (20 mL) and extracted 3x with EtOAc. The EtOAc layer was washed with water, brine, dried with MgSO<sub>4</sub>, filtered, and concentrated in vacuo to provide a crude product that was purified via a 2 inch long filter column using 8:92 EtOAc:Hexanes to provide alcohol **4** as a pale white solid (1.4 g, 86% yield).  $R_f$  = 0.30 (10:90 EA:Hexanes, UV, I<sub>2</sub>); <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  7.73 (d,  $J$  = 5.7 Hz, 1H), 7.37 (d,  $J$  = 6.0 Hz, 1H), 7.26 (s, 1H), 7.22 (d,  $J$  = 6.0 Hz, 1H), 7.13 (t,  $J$  = 5.4 Hz, 1H), 4.87 (s, 2H), 4.20 (m, 1H), 3.15 (d,  $J$  = 8.7 Hz, 2H), 2.33 (br s, 1H), 2.23 (t,  $J$  = 8.4 Hz, 2H), 1.98 (dd,  $J$  = 18.3, 9.0 Hz, 4H), 1.73-1.69 (m, 4H), 1.61 (t,  $J$  = 4.8 Hz, 2H), 1.57 (m, 1H), 1.41 (m, 2H), 1.26 (br s, 1H), 1.02 (br s, 1H), 0.86 (dd,  $J$  = 4.8, 6.9 Hz, 6H); MS(ESI)  $m/z$ : 355.22 [M+H]<sup>+</sup>. Anal. Calcd for C<sub>23</sub>H<sub>34</sub>N<sub>2</sub>O·0.20 CH<sub>2</sub>Cl<sub>2</sub>: C, 75.00; H, 9.33; N, 7.54; found: C, 74.62; H, 9.62; N, 7.34.

**cis-2-amino-5-guanidino-N-((1-(1-(4-isopropylcyclohexyl)piperidin-4-yl)-1H-indol-3-yl)methyl)pentanamide (18, AT-129).** Z-Arg-OH (0.39 g, 0.68 mmol, 1.20 equiv), HOBT (0.099 g, 0.65 mmol, 1.15 equiv), and triethylamine (0.110 mL, 0.79 mmol, 1.40 equiv), were added to 10 mL of CH<sub>2</sub>Cl<sub>2</sub> (0.057M) at room temperature and stirred until the entire solid dissolved. The reaction was cooled in an icebath at 5 °C and EDCI (0.13 g, 0.68 mmol, 1.20 equiv) was added and stirred at 5 °C for 2 hours. At this stage, a solution of amine **3** (0.20 g, 0.57

mmol, 1.00 equiv) in 5 mL of CH<sub>2</sub>Cl<sub>2</sub> was added to the reaction, and then the reaction was allowed to warm to room temperature and was monitored by TLC (10:90:3 drops MeOH:CH<sub>2</sub>Cl<sub>2</sub>:NH<sub>4</sub>OH (aq.)). After 2 hours, the reaction was complete by TLC. The reaction was quenched with saturated NaHCO<sub>3</sub> (aq.), stirred 5 minutes, and the layers were then separated. The aqueous layer was extracted 2x with CH<sub>2</sub>Cl<sub>2</sub>, the organic layers were combined, washed with brine, dried with MgSO<sub>4</sub>, filtered, and concentrated in vacuo to provide a crude residue that was purified via flash chromatography using 3:96:1 MeOH:CH<sub>2</sub>Cl<sub>2</sub>:NH<sub>4</sub>OH (aq.) to provide Cbz intermediate **III-1** as a white solid (0.35 g, 67% yield). R<sub>f</sub> = 0.50 (10:90:3 drops MeOH:CH<sub>2</sub>Cl<sub>2</sub>:NH<sub>4</sub>OH (aq.)), UV, I<sub>2</sub>).

To a solution of Cbz intermediate **III-1** (150 mg, 0.160 mmol, 1.00 equiv) in THF (1.60 mL, 0.10M), was added 10% Pd/C (15.0 mg, 10 wt%), and then the reaction was fitted with a 3-way adapter containing a H<sub>2</sub> (g) balloon. The reaction atmosphere was purged, and then backfilled with H<sub>2</sub> (g). This was repeated 3x total. The reaction was stirred at room temperature and monitored by TLC. After 3 hours, TLC showed full consumption of the starting material. The reaction was poured over a pad of Celite, filtered, and the Celite was washed thoroughly with THF. The resulting filtrate was concentrated in vacuo, and the residue was purified by trituration with CH<sub>2</sub>Cl<sub>2</sub> and Et<sub>2</sub>O to provide amide **18** as a white solid (73 mg, 87% yield). <sup>1</sup>H NMR (300 MHz, DMSO-d<sub>6</sub>): δ 8.32 (s, 2H), 7.72 (br s, 1H), 7.50 (d, *J* = 6.0 Hz, 1H), 7.44 (d, *J* = 6.3 Hz, 1H), 7.41 (s, 1H), 7.08 (t, *J* = 6.0 Hz, 1H), 6.95 (t, *J* = 5.7 Hz, 1H), 4.39 (dd, *J* = 10.8, 3.6 Hz, 1H), 4.31-4.24 (m, 2H), 3.83 (br s, 1H), 3.07 (br s, 3H), 2.90 (m, 1H), 2.66 (s, 2H), 2.28 (d, *J* = 13.5, 2H), 2.16 (t, *J* = 8.4 Hz, 3H), 1.91-1.85 (m, 4H), 1.70 (br s, 2H), 1.54 (br s, 4H), 1.41-1.35 (m, 4H), 1.12 (m, 1H), 0.85 (d, *J* = 4.8 Hz, 6H); MS(ESI) *m/z*: 510.36 [M+H]<sup>+</sup>. Anal.



Calcd for C<sub>29</sub>H<sub>47</sub>N<sub>7</sub>O·2.00 CH<sub>2</sub>Cl<sub>2</sub>: C, 54.79; H, 7.56; N, 14.43; found: C, 54.89; H, 7.31; N, 14.46.

**cis-6-amino-N-((1-(1-(4-isopropylcyclohexyl)piperidin-4-yl)-1H-indol-3-yl)methyl)hexanamide (19, AT-173).** 6-(((benzyloxy)carbonyl)amino)hexanoic acid (0.18 g, 0.68 mmol, 1.20 equiv), HOBT (0.099 g, 0.65 mmol, 1.15 equiv), and triethylamine (0.110 mL, 0.79 mmol, 1.40 equiv) were added to 10 mL of CH<sub>2</sub>Cl<sub>2</sub> (0.057 M) at room temperature and stirred until the entire solid dissolved. The reaction was then cooled in an icebath at 5 °C and EDCI (0.13 g, 0.68 mmol, 1.20 equiv) was added and stirred at 5 °C for 2 h. At this stage, amine **3** (0.20 g, 0.57 mmol, 1.00 equiv) was dissolved in 5 mL of CH<sub>2</sub>Cl<sub>2</sub>, and added to the reaction at 5 °C. The reaction was stirred at 5 °C for 1 hour, and then it was allowed to warm to room temperature and stir for an additional 12 hours. The reaction was quenched with saturated NaHCO<sub>3</sub> (aq.), stirred 5 minutes, and the layers were then separated. The aqueous layer was extracted 2x with CH<sub>2</sub>Cl<sub>2</sub>, the organic layers were combined, washed with brine, dried with MgSO<sub>4</sub>, filtered, and concentrated in vacuo to provide a crude residue that was purified via flash chromatography using 3:96:1 MeOH:CH<sub>2</sub>Cl<sub>2</sub>:NH<sub>4</sub>OH (aq.) to provide carbamate intermediate **III-2** as an oil (0.30 g, 88 % yield). R<sub>f</sub> = 0.50 (10:90:3 drops MeOH:CH<sub>2</sub>Cl<sub>2</sub>:NH<sub>4</sub>OH (aq.), UV, I<sub>2</sub>).

To a solution of Cbz intermediate **III-2** (140 mg, 0.23 mmol, 1.00 equiv) in 2.30 mL of THF (0.10M), was added 10% Pd/C (14.0 mg, 10 wt%), and then the reaction was fitted with a 3-way adapter containing a H<sub>2</sub> (g) balloon. The reaction atmosphere was purged, and then backfilled with H<sub>2</sub> (g). This was repeated 3x total. The reaction was stirred at room temperature and monitored by TLC. After 3 hours, TLC showed full consumption of the starting material. The reaction was poured over a pad of Celite, filtered, and the Celite was washed thoroughly

with THF. The resulting filtrate was concentrated in vacuo, and the residue was purified by trituration with CH<sub>2</sub>Cl<sub>2</sub> and Et<sub>2</sub>O to provide amide **19** as a white solid (42 mg, 39% yield). <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>): δ 7.61 (d, *J* = 7.5 Hz, 1H), 7.36 (d, *J* = 8.1 Hz, 1H), 7.20 (t, *J* = 5.1 Hz, 2H), 7.10 (t, *J* = 7.5 Hz, 1H), 5.93 (s, 1H), 4.59 (d, *J* = 5.1 Hz, 2H), 4.21 (m, 1H), 3.57 (t, *J* = 5.4 Hz, 4H), 3.22 (d, *J* = 11.4 Hz, 2H), 2.56-2.50 (m, 3H), 2.43-2.32 (m, 2H), 2.17 (t, *J* = 7.5 Hz, 2H), 2.12 (s, 2H), 1.78-1.51 (m, 11H), 1.45-1.31 (m, 4H), 1.29-1.16 (m, 1H), 0.90 (d, *J* = 6.6 Hz, 6H); MS(ESI) *m/z*: 467.30 [M+H]<sup>+</sup>. The bis HCl salt of amine **19** was formed by dissolving the amine in CH<sub>2</sub>Cl<sub>2</sub> (0.10M) at room temperature, followed by the addition of HCl (2.0M in Et<sub>2</sub>O, 2.20 equiv). The reaction was allowed to stir for 10 minutes. At this time, the reaction was concentrated in vacuo, Et<sub>2</sub>O was added to the mixture, followed by concentrating the mixture in vacuo (repeat 3x total) to provide the HCl salt of amine **19**. Anal. Calcd for C<sub>29</sub>H<sub>46</sub>N<sub>4</sub>O·2.00 HCl·2.00 CH<sub>2</sub>Cl<sub>2</sub>·2.00 Et<sub>2</sub>O·1.00 H<sub>2</sub>O: C, 53.49; H, 8.52; N 6.40; found: C, 53.73; H, 8.39; N, 6.20.

**cis-2-(ethylamino)-N-((1-(1-(4-isopropylcyclohexyl)piperidin-4-yl)-1H-indol-3-yl)methyl)acetamide (14, AT-122).** Z-glycine-OH (0.20 g, 0.67 mmol, 1.20 equiv), HOBT (0.099 g, 0.65 mmol, 1.15 equiv), and triethylamine (0.110 mL, 0.79 mmol, 1.40 equiv) were added to 10 mL CH<sub>2</sub>Cl<sub>2</sub> (0.057 M) and stirred until the entire solid dissolved. The reaction was then cooled in an icebath at 5 °C and EDCI (0.13 g, 0.68 mmol, 1.20 equiv) was added and stirred at 5 °C for 2 h. At this stage, amine **3** (0.20 g, 0.57 mmol, 1.00 equiv) was dissolved in 5 mL of CH<sub>2</sub>Cl<sub>2</sub>, and added to the reaction at 5 °C. The reaction was allowed to warm to room temperature and stir for an additional 2 hours. The reaction was quenched with saturated NaHCO<sub>3</sub> (aq.), stirred 5 minutes, and the layers were then separated. The aqueous layer was extracted 2x with CH<sub>2</sub>Cl<sub>2</sub>, the organic layers were combined, washed with brine, dried with

MgSO<sub>4</sub>, filtered, and concentrated in vacuo to provide a crude residue that was purified via flash chromatography using 1:98:1 MeOH:CH<sub>2</sub>Cl<sub>2</sub>:NH<sub>4</sub>OH (aq.) to provide Cbz intermediate **III-3** (0.21 g, 68% yield). <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 7.59 (d, *J* = 5.7 Hz, 1H), 7.36 (d, *J* = 6.3 Hz, 1H), 7.32 (s, 5H), 7.24 (m, 2H), 7.11 (t, *J* = 5.7 Hz, 1H), 6.11 (s, 1H), 5.45 (s, 1H), 5.08 (s, 2H), 4.19 (m, 1H), 3.85 (d, *J* = 4.2 Hz, 2H), 3.22 (d, *J* = 8.7 Hz, 2H), 2.44 (s, 1H), 2.31 (d, *J* = 7.8 Hz, 2H), 2.09 (t, *J* = 7.8 Hz, 4H), 1.73 (br s, 2H), 1.65 (m, 3H), 1.43-1.38 (m, 2H), 1.17 (m, 1H), 0.89 (d, *J* = 4.8 Hz, 6H); MS(ESI) *m/z*: 545.28 [M+H]<sup>+</sup>.

To a solution of Cbz intermediate **III-3** (0.15 g, 0.308 mmol, 1.00 equiv) in 3.1 mL of absolute ethanol (0.10M), was added 10% Pd/C (15.0 mg, 10 wt%), and then the reaction was fitted with a 3-way adapter containing a H<sub>2</sub> (g) balloon. The reaction atmosphere was purged, and then backfilled with H<sub>2</sub> (g). This was repeated 3x total. The reaction was stirred at room temperature and monitored by TLC. After 4 hours, TLC showed full consumption of the starting material. The reaction was poured over a pad of Celite, filtered, and the Celite was washed thoroughly with ethanol. The resulting filtrate was concentrated in vacuo, and the residue was purified by flash chromatography on silica gel, using 5:95:1 MeOH:CH<sub>2</sub>Cl<sub>2</sub>:NH<sub>4</sub>OH (aq.) to provide ethylated amine **14** as an oil (55.2 mg, 41% yield). *R<sub>f</sub>* = 0.50 (10:90 MeOH:CH<sub>2</sub>Cl<sub>2</sub>:NH<sub>4</sub>OH (aq.)), UV, I<sub>2</sub>). <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 7.59 (t, *J* = 7.8 Hz, 1H), 7.35 (d, *J* = 8.1 Hz, 1H), 7.20 (t, *J* = 4.8 Hz, 2H), 7.09 (t, *J* = 7.5 Hz, 1H), 5.31 (s, 1H), 4.60 (d, *J* = 5.4 Hz, 2H), 4.32 (br s, 1H), 3.38 (m, 4H), 2.71-2.44 (m, 9H), 2.11 (d, *J* = 11.7 Hz, 2H), 1.88-1.64 (m, 5H), 1.40 (dd, *J* = 10.2, 13.2 Hz, 2H), 1.28-1.20 (m, 1H), 1.07 (t, *J* = 6.9 Hz, 3H), 0.92 (d, *J* = 6.3 Hz, 6H); MS(ESI) *m/z*: 439.28 [M+H]<sup>+</sup>. Anal. Calcd for C<sub>27</sub>H<sub>42</sub>N<sub>4</sub>O·0.50 H<sub>2</sub>O·0.50 CH<sub>2</sub>Cl<sub>2</sub>: C, 67.39; H, 9.05; N, 11.43; found: C, 67.02; H, 9.01; N, 11.69.

#### Synthesis of alkynes **IV-3** and **IV-4**.

**tert-butyl (5-(trimethylsilyl)pent-4-yn-1-yl)carbamate (IV-4).** To a solution of 5-(trimethylsilyl)pent-4-yn-1-ol (2.55 g, 16.3 mmol, 1.00 equiv) (GFS Chemicals) in 65.2 mL of CH<sub>2</sub>Cl<sub>2</sub> (0.25M) at room temperature, was added triethylamine (2.95 mL, 21.2 mmol, 1.30 equiv) and DMAP (99.6 mg, 0.815 mmol, 0.050 equiv). The solution was stirred for several minutes, and then TsCl (3.36 g, 17.6 mmol, 1.08 equiv) was added at room temperature. The reaction was fitted with an Ar balloon and allowed to stir for 16 hours. At this stage, TLC (10:90 EtOAc:Hexanes) showed the reaction was complete. The reaction was concentrated in vacuo to half of the initial reaction volume, then diluted with EtOAc and water, and the biphasic solution stirred for five minutes. The layers were then separated, and the aqueous layer was extracted 2x with EtOAc. The EtOAc layers were combined, washed with brine, dried with MgSO<sub>4</sub>, filtered, and concentrated in vacuo to provide the tosylate intermediate used directly in the following step (4.70 g, 93% yield).

The tosylate intermediate (4.70 g, 15.1 mmol, 1.00 equiv) and phthalimide (2.78 g, 18.9 mmol, 1.25 equiv) were dissolved in 60.0 mL of DMF (0.25M) at room temperature. K<sub>2</sub>CO<sub>3</sub> (6.27 g, 45.4 mmol, 3.00 equiv) was added, the reaction was fitted with an Ar balloon, and the reaction was heated to 50 °C for 6 hours. At this time, TLC (10:90 EtOAc:Hexanes) shows the reaction is complete. The reaction was allowed to cool to room temperature, and then it was diluted with EtOAc and water. The layers were separated, and the aqueous layer was extracted 2x with EtOAc. The EtOAc layers were combined, washed 3x with water, 1x with brine, dried with MgSO<sub>4</sub>, filtered, and concentrated in vacuo to provide semi-crude material. This material was purified via flash chromatography using 8:92 EtOAc:Hexanes to provide the phthalimide intermediate as a white solid (2.89 g, 67% yield). This material was used directly in the following step.

The phthalimide intermediate (2.20 g, 7.71 mmol, 1.00 equiv) from the previous step was dissolved in 77.0 mL of MeOH (0.10M) at room temperature, and then hydrazine (0.966 mL, 30.8 mmol, 4.00 equiv) was added, and the reaction was allowed to stir at room temperature overnight. At this stage, a large amount of a white precipitate had formed. The white precipitate was filtered, washed thoroughly with MeOH, and the resulting filtrate was concentrated in vacuo to provide a solid-liquid mixture. This mixture was diluted with EtOAc, stirred several minutes, and the resulting solids were filtered, and washed thoroughly with EtOAc. The filtrate (EtOAc), was then washed with water, brine, dried with MgSO<sub>4</sub>, filtered, and concentrated in vacuo to provide the desired amine (863 mg, 72% yield), which is used directly in the following step.

The amine intermediate (863 mg, 5.56 mmol, 1.00 equiv) and DMAP (34.0 mg, 0.278 mmol, 0.050 equiv) were dissolved in 37 mL of THF (0.15M) at room temperature. To this solution was added Boc anhydride (1.33 g, 6.11 mmol, 1.10 equiv), and the reaction was allowed to stir overnight. At this stage, TLC (2:98:3 drops MeOH:CH<sub>2</sub>Cl<sub>2</sub>:NH<sub>4</sub>OH (aq.)) showed the reaction was complete. The reaction was concentrated in vacuo, and the crude mixture was purified via flash chromatography using 5:95 EtOAc:Hexanes to provide alkyne **IV-4** as a clear oil (1.25 g, 88% yield). R<sub>f</sub> = 0.10 (5:95 EtOAc:Hexanes, I<sub>2</sub> and pAA); <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 4.80 (br, 1H), 3.23 (m, 2H), 2.29 (t, *J* = 5.4 Hz, 2H), 1.71 (m, 2H), 1.49 (s, 9H), 0.16 (s, 9H).

**tert-butyl (4-(triethylsilyl)but-3-yn-1-yl)carbamate (IV-3)**. Alkyne **IV-3** was obtained as a light-yellow oil using the same 4-step synthesis that provided alkyne **IV-4**. Step 1: 99% yield. Step 2: 94% yield. Step 3: 70% yield. Step 4: 78% yield. R<sub>f</sub> = 0.10 (5:95 EtOAc:Hexanes, I<sub>2</sub> and pAA); <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ 4.80 (br, 1H), 3.28 (q, *J* = 6.6 Hz, 2H), 2.45 (t, *J* = 6.6 Hz, 2H), 1.45 (s, 9H), 0.99 (t, *J* = 7.8 Hz, 9H), 0.58 (q, *J* = 7.8 Hz, 6H).

**cis-N-(2-iodophenyl)-1-(4-isopropylcyclohexyl)piperidin-4-amine (IV-6).**

*i.* See General Procedure A. 2-iodoaniline (15.0 g, 63.3 mmol, 1.00 equiv), N-Boc-piperidone (18.5 g, 95.0 mmol, 1.50 equiv), glacial AcOH (8.40 mL, 146 mmol, 2.30 equiv), STAB (30.9 g, 146 mmol, 2.30 equiv), DCE (250 mL, 0.25M). Purified via flash chromatography using 5:95 EtOAc:Hexanes to provide the secondary aniline intermediate as a white solid (19.1 g, 75% yield).

*ii.* See General Procedure B. **Step 1:** Aniline intermediate (43.5 g, 0.108 mol, 1.00 equiv), TFA (200 mL, 2.61 mol, 24.0 equiv), CH<sub>2</sub>Cl<sub>2</sub> (300 mL, 0.36M). Obtained the N-H piperidine intermediate as a light-tan solid (42.0 g, 128% yield, due to NaTFA), and was used directly in the next step. **Step 2:** N-H piperidine (assume 0.108 mol, 1.00 equiv), 4-*i*Pr-cyclohexanone (22.7 g, 0.162 mol, 1.50 equiv), glacial AcOH (14.2 mL, 0.248 mol, 2.30 equiv), STAB (52.6 g, 0.248 mol, 2.30 equiv), DCE (1.54 L, 0.070M). Compound **IV-6** was separated from its anti-diastereomer via flash chromatography using 6:94:1.5→9:91:1.5 EtOAc:Hexanes:NH<sub>4</sub>OH (aq.) to provide a golden oil (The syn diastereomer has a higher R<sub>f</sub> value compared to the anti diastereomer). Although the product is pure when viewed under UV light (short wave), the product contains non-UV active impurities (viewable with pAA) related to 4-*i*Pr-cyclohexanone. Thus, to remove these impurities the golden oil was dissolved in EtOAc (ca. 300 mL) and transferred to an erlenmeyer flask. At this stage, 10% HCl (aq.) (ca. 200 mL) was added. Upon addition of the 10% HCl (aq.), a white precipitate formed, and the suspension was stirred for 10 minutes. The white precipitate was then filtered, washed 2x with EtOAc, and then air-dried over 1 hour. The white precipitate was then placed in an erlenmeyer flask, suspended in EtOAc, and 70% NaHCO<sub>3</sub> (aq.) was added until basic. The mixture was then stirred overnight. At this stage, the mixture is now a clear biphasic solution (add water and stir if

significant amount of white precipitate remain). The layers were separated, and the EtOAc layer was washed with water, brine, dried with MgSO<sub>4</sub>, filtered, and concentrated in vacuo to provide iodoaniline **IV-6** as a light-gold oil (24.0 g, 39% yield over 3 steps).  $R_f = 0.30$  (10:90:3 drops EtOAc:Hexanes:NH<sub>4</sub>OH (aq.)), UV, I<sub>2</sub>); <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>)  $\delta$  7.65 (dd,  $J = 5.7, 0.9$ , 1H), 7.18 (t,  $J = 6.0$ , 1H), 6.58 (d,  $J = 6.0$ , 1H), 6.41 (dt,  $J = 5.7, 0.9$ , 1H), 4.12 (d,  $J = 5.7$  Hz, 1H), 3.36 (m, 1H), 2.93 (m, 2H), 2.25 (m, 3H), 2.15 (d,  $J = 8.4$  Hz, 2H), 1.47-1.74 (m, 8H), 1.38 (m, 2H), 1.13 (m, 1H), 0.89 (d,  $J = 4.8$  Hz, 6H); MS(ESI)  $m/z$ : 427 [M+H]<sup>+</sup>.

**cis-tert-butyl (2-(1-(1-(4-isopropylcyclohexyl)piperidin-4-yl)-2-(triethylsilyl)-1H-indol-3-yl)ethyl)carbamate (IV-7)**. Iodoaniline **IV-6** (1.00 g, 2.35 mmol, 1.00 equiv) and alkyne **IV-3** (797 mg, 2.81 mmol, 1.20 equiv) were charged into a round-bottom flask. LiCl (99.4 mg, 2.35 mmol, 1.00 equiv) was added, and the mixture was diluted with 33.5 mL of DMF (0.070M). K<sub>2</sub>CO<sub>3</sub> (973 mg, 7.04 mmol, 3.00 equiv) was added to the reaction, followed by a catalytic amount of Pd(OAc)<sub>2</sub> (26.3 mg, 0.117 mmol, 0.050 equiv). The reaction was fitted with an Ar balloon, purged 3x under vacuum and backfilled with Ar, and heated in a 102 °C oil bath. The reaction was monitored by TLC, and was complete at ca. 2.5 hours. NOTE: Pd<sup>0</sup> (black) was observed ca. 2 hours into reaction. Once the reaction was complete, the mixture was allowed to cool to room temperature and was diluted with EtOAc and water. The mixture was stirred for several minutes, and then filtered. The filtrate layers were separated, and the aqueous layer was extracted 1x with EtOAc. The EtOAc layers were combined and washed 2x with water, 1x with brine, dried with MgSO<sub>4</sub>, filtered, and concentrated in vacuo to provide a crude product that was purified via flash chromatography using 7:93:1→10:90:1 EtOAc:Hexanes:NH<sub>4</sub>OH (aq.) to provide indole **IV-7** as a white foam (1.04 g, 76% yield).  $R_f = 0.25$  (10:90:3 drops EtOAc:Hexanes:NH<sub>4</sub>OH (aq.)), UV, I<sub>2</sub>); <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz)  $\delta$  7.69 (d,  $J = 6.3$  Hz, 1H),

7.62 (d,  $J = 5.7$  Hz, 1H), 7.16 (t,  $J = 5.7$  Hz, 1H), 7.06 (t,  $J = 5.7$  Hz, 1H), 4.56 (br, 1H), 4.25 (m, 1H), 3.40 (m, 2H), 3.21 (d,  $J = 8.4$  Hz, 2H), 3.02 (t,  $J = 5.1$  Hz, 2H), 2.71 (dq,  $J = 9.0, 2.1$  Hz, 2H), 2.35 (m, 1H), 2.17 (t,  $J = 9.0$  Hz, 2H), 1.38–1.75 (m, 20H), 1.16 (m, 1H), 0.90–1.03 (m, 21H); MS(ESI)  $m/z$ : 582.8 [M+H]<sup>+</sup>.

**cis-tert-butyl (3-(1-(1-(4-isopropylcyclohexyl)piperidin-4-yl)-2-(trimethylsilyl)-1H-indol-3-yl)propyl)carbamate (IV-8).** Indole **IV-8** was synthesized in a similar manner as indole **IV-7**. Iodoaniline **IV-6** (720 mg, 1.69 mmol, 1.00 equiv), alkyne **IV-4** (475 mg, 1.86 mmol, 1.10 equiv), LiCl (71.6 mg, 1.69 mmol, 1.00 equiv), K<sub>2</sub>CO<sub>3</sub> (700 mg, 5.07 mmol, 3.00 equiv), Pd(OAc)<sub>2</sub> (38.0 mg, 0.169 mmol, 0.100 equiv), and DMF (24.0 mL, 0.070M). Reaction Time of ca. 6 hours. Purified via flash chromatography using 10:90:1 EtOAc:Hexanes:NH<sub>4</sub>OH (aq.) to provide indole **IV-8** in ca. 90% purity as light-beige solid (88% yield). Trituration using MeOH provided pure indole **IV-8** as a white solid (585 mg, 65% yield).  $R_f = 0.10$  (10:90:3 drops EtOAc:Hexanes:NH<sub>4</sub>OH (aq.), UV, I<sub>2</sub>); MS(ESI)  $m/z$ : 554.8 [M+H]<sup>+</sup>.

**cis-2-(1-(1-(4-isopropylcyclohexyl)piperidin-4-yl)-1H-indol-3-yl)ethan-1-amine (6, AT-160).** AcCl (806  $\mu$ L, 11.3 mmol, 6.00 equiv) was added to 19.0 mL of MeOH (0.10M) at 0 °C, and the mixture was stirred for several minutes at 0 °C. At this stage, indole intermediate **IV-7** (1.10 g, 1.89 mmol, 1.00 equiv) was added, resulting in a slurry. The ice-bath was then removed, and the reaction was allowed to warm to room temperature and stir overnight. Once TLC confirmed the reaction was complete, EtOAc (ca. 5x the amount of MeOH) was added and the reaction was allowed to stir for 60 minutes, during which a white precipitate formed. The white precipitate was then filtered and washed 3x with EtOAc. The white solid was then collected, and dried in vacuo to provide the bis HCl salt of indole **6** as a white solid (665 mg, 80% yield).  $R_f = 0.10$  (10:90:3 drops iPrOH:CH<sub>2</sub>Cl<sub>2</sub>:NH<sub>4</sub>OH (aq.), UV, I<sub>2</sub>); <sup>1</sup>H NMR (CDCl<sub>3</sub>,



300 MHz)  $\delta$  7.61 (d,  $J = 6.0$  Hz, 1H), 7.36 (d,  $J = 6.3$  Hz, 1H), 7.21 (t,  $J = 6.0$  Hz, 1H), 7.11 (m, 2H), 4.18 (m, 1H), 3.19 (d,  $J = 8.7$  Hz, 2H), 3.03 (t,  $J = 4.8$  Hz, 2H), 2.93 (t,  $J = 4.8$  Hz, 2H), 2.35 (m, 1H), 2.26 (dt,  $J = 8.4, 1.8$  Hz, 2H), 2.07 (br, 6H), 1.76-1.53 (m, 7H), 1.42 (m, 2H), 1.14 (m, 1H), 0.90 (d,  $J = 5.1$  Hz, 6H); MS(ESI)  $m/z$ : 368.6  $[M+H]^+$ . Anal. Calcd for  $C_{24}H_{37}N_3 \cdot 2.00$  HCl $\cdot 0.50$  H $_2$ O: C, 64.12; H, 8.97; N, 9.35; found: C, 64.23; H, 8.73; N, 9.32.

**cis-3-(1-(1-(4-isopropylcyclohexyl)piperidin-4-yl)-1H-indol-3-yl)propan-1-amine (7, AT-147).** Compound **7** was synthesized in a similar manner as indole **6** (AT-160). Indole intermediate **IV-8** (582 mg, 1.09 mmol, 1.00 equiv), AcCl (466  $\mu$ L, 6.55 mmol, 6.00 equiv), MeOH (7.3 mL, 0.15M). Obtained the bis HCl salt of indole **7** as a white solid (388 mg, 78% yield).  $R_f = 0.10$  (5:95:3 drops MeOH:CH $_2$ Cl $_2$ :NH $_4$ OH (aq.), UV, I $_2$ );  $^1$ H NMR (CD $_3$ OD, 300 MHz)  $\delta$  7.58 (d,  $J = 6.0$  Hz, 1H), 7.50 (d,  $J = 6.3$  Hz, 1H), 7.25 (s, 1H), 7.19 (t,  $J = 6.0$  Hz, 1H), 7.07 (t,  $J = 6.0$  Hz, 1H), 4.72 (m, 1H), 3.74 (d,  $J = 9.6$  Hz, 2H), 3.40-3.30 (m, 5H), 2.98 (t,  $J = 5.7$  Hz, 2H), 2.88 (t,  $J = 5.7$  Hz, 2H), 2.50 (dq,  $J = 9.6, 2.4$  Hz, 2H), 2.28 (d,  $J = 9.9$  Hz, 2H), 2.10-1.95 (m, 6H), 1.80 (m, 3H), 1.57 (t,  $J = 9.9$  Hz, 2H), 1.28 (m, 1H), 0.97 (d,  $J = 5.1$  Hz, 6H); MS(ESI)  $m/z$ : 382.6  $[M+H]^+$ . Anal. Calcd for  $C_{25}H_{39}N_3 \cdot 2.00$  HCl $\cdot 0.50$  H $_2$ O: C, 64.78; H, 9.13; N, 9.07; found: C, 64.93; H, 9.11; N, 9.04.

**cis-1-((1-(1-(4-isopropylcyclohexyl)piperidin-4-yl)-1H-indol-3-yl)methyl)thiourea (15, AT-111).** See **General Procedure C. Step 1.** Amine (200 mg, 0.565 mmol, 1.00 equiv), Benzoyl isothiocyanate (83.6 mg, 0.622 mmol, 1.10 equiv), CH $_2$ Cl $_2$  (5.60 mL, 0.10 M). **Step 2.** K $_2$ CO $_3$  (547 mg, 3.96 mmol, 7.00 equiv), MeOH (5.80 mL, 0.050 M). Obtained thiourea **15** as a light-yellow glue (70.0 mg, 58% yield).  $R_f = 0.50$  (10:90:3 drops MeOH:CH $_2$ Cl $_2$ :NH $_4$ OH (aq.), UV, I $_2$ );  $^1$ H NMR (300 MHz, CDCl $_3$ ):  $\delta$  7.61 (br s, 1H), 7.37 (d,  $J = 6$ Hz, 1H), 7.24 (s, 2H), 7.11 (t,  $J = 5.1$  Hz, 1H), 6.58 (br s, 1H), 5.82 (s, 2H), 4.88 (br s, 1H), 4.45 (br s, 1H), 4.18 (s, 1H),

3.14 (d, J = 6.9 Hz, 2H), 2.33 (s, 1H), 2.22 (t, J = 9.3 Hz, 2H), 2.02-1.97 (m, 4H), 1.71-1.61 (m, 5H), 1.55 (m, 2H), 1.41 (m, 2H), 1.14 (m, 1H), 0.89 (d, J = 4.8 Hz, 6H); MS(ESI) m/z: 413.11 [M+H]<sup>+</sup>. The bis HCl salt of urea **15** was formed by dissolving the amine in CH<sub>2</sub>Cl<sub>2</sub> (0.10M) at room temperature, followed by the addition of HCl (2.0M in Et<sub>2</sub>O, 2.20 equiv). The reaction was allowed to stir for 10 minutes. At this time, the reaction was concentrated in vacuo, Et<sub>2</sub>O was added to the mixture, followed by concentrating the mixture in vacuo (repeat 3x total) to provide the HCl salt of urea **15**. Anal. Calcd for C<sub>24</sub>H<sub>36</sub>N=S·2.00 HCl·1.00 CH<sub>2</sub>Cl<sub>2</sub>: C, 55.76; H, 7.48; N, 10.40; found: C, 55.66; H, 7.52; N, 10.74.

**cis-1-(2-(1-(1-(4-isopropylcyclohexyl)piperidin-4-yl)-1H-indol-3-yl)ethyl)thiourea**

(**16**, AT-163). See General Procedure C. Step 1. Amine (35.0 mg, 95.2 μmol, 1.00 equiv), Benzoyl isothiocyanate (13.5 μL, 100 μmol, 1.05 equiv), CH<sub>2</sub>Cl<sub>2</sub> (950 μL, 0.10 M). Step 2. K<sub>2</sub>CO<sub>3</sub> (92.0 mg, 666 μmol, 7.00 equiv), MeOH (1.90 mL, 0.050 M). Obtained thiourea **16** as a light-yellow glue (20.2 mg, 50% yield). R<sub>f</sub> = 0.15 (60:40:3 drops EtOAc:Hexanes:NH<sub>4</sub>OH (aq.)), UV, I<sub>2</sub>; <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ 7.57 (br s, 1H), 7.37 (d, J = 6.3 Hz, 1H), 7.23 (t, J = 5.4 Hz, 1H), 7.12 (m, 2H), 6.27 (br s, 1H), 5.84 (br s, 2H), 4.20 (m, 1H), 3.22 (d, J = 8.4 Hz, 2H), 3.06 (t, J = 4.8 Hz, 2H), 2.42-2.07 (m, 7H), 1.78-1.56 (m, 7H), 1.41 (m, 2H), 1.27 (m, 2H), 1.16 (m, 1H), 0.91 (d, J = 5.1 Hz, 6H); MS(ESI) m/z: 427.7 [M+H]<sup>+</sup>. HRMS (ESI) calcd for C<sub>25</sub>H<sub>39</sub>N<sub>4</sub>S (M+H), 427.2895; found, 427.2896.

**cis-1-(3-(1-(1-(4-isopropylcyclohexyl)piperidin-4-yl)-1H-indol-3-yl)propyl)thiourea**

(**17**, AT-155). See General Procedure C. Step 1. Amine (20.0 mg, 52.4 μmol, 1.00 equiv), Benzoyl isothiocyanate (7.40 μL, 55.0 μmol, 1.05 equiv), CH<sub>2</sub>Cl<sub>2</sub> (525 μL, 0.10 M). Step 2. K<sub>2</sub>CO<sub>3</sub> (50.0 mg, 0.367 mmol, 7.00 equiv), MeOH (1.05 mL, 0.050 M). Obtained thiourea **17** as a clear oil (10.8 mg, 47% yield). R<sub>f</sub> = 0.20 (50:50:3 drops EtOAc:Hexanes:NH<sub>4</sub>OH (aq.)), UV,

pAA, I<sub>2</sub>); <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz) δ 7.56 (d, J = 6.0 Hz, 1H), 7.36 (d, J = 6.0 Hz, 1H), 7.21 (t, J = 5.7 Hz, 1H), 7.10 (m, 2H), 6.26 (br s, 1H), 5.66 (br s, 2H), 4.18 (m, 1H), 3.18 (m, 3H), 2.85 (t, J = 5.4 Hz, 2H), 2.40-2.25 (m, 3H), 2.12-1.90 (m, 6H), 1.77-1.54 (m, 6H), 1.42 (m, 2H), 1.26 (m, 2H), 1.15 (m, 1H), 0.90 (d, J = 4.8 Hz, 6H); MS(ESI) m/z: 427.7 [M+H]<sup>+</sup>. Anal. Calcd for C<sub>26</sub>H<sub>40</sub>N<sub>4</sub>S·0.50 H<sub>2</sub>O: C, 69.44; H, 9.19; N, 12.46; found: C, 69.43; H, 9.13; N, 12.07.

### **In vitro pharmacological Characterization**

*Cells.* Human NOP, mu, delta, and kappa opioid receptors were individually expressed in Chinese hamster ovary cells stably transfected with the human receptor cDNA, as we have described previously.(Zaveri , et al. 2001; Toll et al. JPET, 2009) Kappa-CN cells were used for KOP radioligand binding assays, while Kappa-FLG19 cells were used in KOP [<sup>35</sup>S]GTPγS functional assays. The cells were grown in Dulbecco's Modified Eagle Medium (DMEM) with 10% fetal bovine serum, in the presence of 0.4 mg/ml G418 and 0.1% penicillin/streptomycin, in 100-mm plastic culture dishes.

*Membrane preparation.* The cell lines are grown to full confluency, then harvested for membrane preparation. The membranes are prepared in 50 mM Tris buffer (pH 7.4). Cells are scraped and centrifuged at 500 × g for 12 mins. The cell pellet is homogenized in 50 mM Tris with a Fisher Scientific PowerGen 125 rotor-stator type homogenizer, centrifuged at 20,000 × g for 25 mins, washed and recentrifuged once more and aliquoted at a concentration of 3 mg/mL protein per vial and stored in a -80 °C freezer till further use.

*Receptor Binding.* The assay is performed in a 96-well polystyrene plate using triplicates of six concentrations of each test compound and tritiated ligands [<sup>3</sup>H]DAMGO (0.2 nM for MOP), [<sup>3</sup>H]DPDPE (0.2 nM for DOP), [<sup>3</sup>H]U69593 (0.2 nM for KOP), or [<sup>3</sup>H]N/OFQ (0.2 nM for NOP). Nonspecific binding was determined by using 1.0 μM of the unlabeled nociceptin for

NOP, 10  $\mu$ M unlabeled DAMGO for MOP, 10  $\mu$ M unlabeled DPDPE for DOP, and 10  $\mu$ M unlabeled U69,593 for KOP. Assays were initiated by addition of membrane homogenates and samples were incubated for 60 min at 25°C in a total volume of 1.0 mL. In NOP receptor experiments, 1 mg/mL BSA is added to the assay buffer. The amount of protein in the binding assay was 15  $\mu$ g. The incubation was terminated by rapid filtration through 0.5% PEI-soaked glassfiber filter mats (GF/C Filtermat A, Perkin-Elmer) on a Tomtec Mach III cell harvester and washed 5 times with 0.5 mL of ice-cold 50 nM Tris-HCl, pH 7.4 buffer. The filters were dried overnight and soaked with scintillation cocktail before counting on a Wallac Beta plate 1205 liquid scintillation counter. Radioactivity was determined as counts per minutes (cpm). Full characterization of compounds includes analysis of the data for IC50 values and Hill coefficients using GraphPad Prism. (ISI, San Diego, CA). Ki values were determined by the method of Cheng and Prusoff.(Cheng and Prusoff, 1973)

*[<sup>35</sup>S]GTP $\gamma$ S Functional assay.* The efficacy of the compounds were determined by the ability to stimulate [<sup>35</sup>S]GTP $\gamma$ S binding to cell membranes, and compared to standard agonists. The functional assay was conducted in Buffer A, containing 20 mM HEPES (4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid), 10 mM magnesium chloride (MgCl<sub>2</sub>) 100 mM sodium chloride (NaCl) at pH 7.4. Membrane prepared as described above was incubated with [<sup>35</sup>S]GTP $\gamma$ S (150,000 cpm/well), guanosine diphosphate (GDP) (10 mM), and the test compound, in a total volume of 1 ml for 60 minutes at 25°C. Samples were filtered over Filtermat A and counted as described for the binding assays. A dose response curve with a prototypical full agonist at the respective receptor was conducted in each experiment to identify full and partial agonist compounds. Typically, the standard full agonists (N/OFQ for NOP, DAMGO for MOP, U69593 for KOP, and DPDPE DOP) showed at least 2-fold to 5-fold

stimulation over basal. The stimulation by the standard full agonists was taken as 100% when comparing stimulation by the test compound (Table 1).

### **In vivo efficacy in the 6-OH hemilesioned rat model of Parkinson's disease**

#### *Animal Subjects*

Experiments were performed in accordance with the ARRIVE guidelines. Experimenters were blinded to treatments. Male Sprague-Dawley rats (150 g, 6 week old; Envigo, S. Pietro al Natisone, Italy) were housed in a standard facility with free access to food (4RF21 standard diet; Mucedola, Settimo Milanese, Milan, Italy) and water, and kept under regular lighting conditions (12 hr dark/light cycle). Animals were housed in groups of 5 for a 55x33x20 cm polycarbonate cage (Tecniplast, Buguggiate, Varese, Italy) with a Scobis Uno bedding (Mucedola, Settimo Milanese, Milan, Italy) and environmental enrichments. The experimental protocols were approved by the Italian Ministry of Health (license n. 170/2013-B). Adequate measures were taken to minimize animal pain and discomfort. At the end of the experiments, rats were sacrificed with an overdose of isoflurane.

#### *Unilateral 6-OHDA lesion*

The unilaterally 6-OHDA lesioned rat, the most popular and best validated model of PD (Duty et al., 2011; Schwarting et al., 1996), was used to assess the ability of NOP partial agonists to improve parkinsonian-like motor deficits. Unilateral lesion of dopaminergic neurons was induced under isoflurane anaesthesia as previously described (Marti et al., 2005, 2007). Eight micrograms of 6-OHDA freebase (dissolved in 0.9% saline solution containing 0.02% ascorbic acid) were stereotaxically injected in the medial forebrain bundle according to following coordinates from bregma: antero-posterior= -4.4 mm, medio-lateral= -1.2, dorso-ventral=-7.8 mm below dura (Paxinos et al., 1986). Animals were pre-treated with antibiotics (Synulox™, 50

$\mu\text{L}/\text{Kg}$ , i.p.). The wound was sutured and infiltrated with 2% lidocaine solution (Esteve™). In order to select rats that were successfully hemilesioned, two weeks after 6-OHDA injection, rats were injected with a test dose of D-amphetamine (5 mg kg<sup>-1</sup>, i.p., dissolved in saline), and those performing >7 turns per minute in the direction ipsilateral to the lesion were enrolled in the study. In fact, this behaviour has been associated with >95% loss of striatal dopaminergic terminals (Marti et al., 2007) and extracellular dopaminergic levels (Marti et al., 2002).

### *Behavioral Tests*

*Bar Test.* The bar test (Marti, et al., 2005), also known as catalepsy test (Sanberg et al., 1988), measures the ability of the animal to react to an externally imposed position. The right and left forepaws were alternatively placed on three blocks of increasing heights (3, 6, 9 cm). The immobility time (in sec) of each forepaw on the blocks was recorded (the cut-off for each step was set at 20 sec).

*Drag Test.* The drag test (Marti et al., 2005), modification of the “wheelbarrow test” (Schallert et al., 1979), measures the ability of the animal to balance its body posture using the forelimbs, in response to backward dragging. Each rat was gently lifted from the abdomen leaving the forepaws on the table and dragged backwards at a constant speed of 20 cm sec<sup>-1</sup> for a fixed distance of 1 m. Two different observers counted the number of touches made by each forepaw.

*Rotarod Test.* This test measures the ability of the animal to run on a rotating cylinder and provides different information on a variety of motor parameters such as coordination, balance, muscle tone, gait and motivation to run (Rozas et al., 1997). The fixed-speed rotarod test was employed using a previously validated protocol (Marti et al., 2005; Marti et al., 2004

JNS paper). Animals were tested starting from 5 rpm, speed was stepwise increased by 5 rpm every 180 sec, and total time spent on the rod was calculated.

## **Experimental protocols and design**

### **Motor behavior in 6-OHDA hemilesioned, L-DOPA naïve rats**

Hemilesioned rats which met the selection criteria after amphetamine testing after 2 weeks were used in the behavioral tests. Ten days later, these rats were subjected to the bar, drag and rotarod tests repeated in a fixed sequence as training. When motor performance was reproducible (usually after 7-10 days), rats were randomized and treated with the respective doses of the test partial agonist, L-DOPA or vehicle. Rats were tested 3-4 times, and a 3-day washout was allowed between treatments. In the bar and drag test, motor activity was assessed 20 and 90 min after compound administration both at the contralateral and ipsilateral paw, and expressed as absolute values for the bar and drag tests, and as percentage of the control session for the rotarod test. Since no motor changes were observed at the ipsilateral paw after compound administration, data have not been shown.

### **Statistical analysis**

Statistical analysis was performed using PRISM 6.0 (GraphPad Software Inc., San Diego, CA) with two-way ANOVA followed by the Bonferroni test. Statistical significance was set at  $p < 0.05$ .

## **SUPPORTING INFORMATION**

SMILES data (Molecular formula strings) for all final compounds along with their in vitro pharmacological data are provided.

## **AUTHOR CONTRIBUTIONS**

U.G.K. and M.E.M. contributed equally to this work.

## **ACKNOWLEDGEMENTS**

This work was supported by grants R01DA014026 (N.T.Z.), R01DA027811 (N.T.Z.), and R43NS070664 (N.T.Z.) from the National Institutes of Health.

## **ABBREVIATIONS USED**

DOP, delta opioid receptor; GPCR, G-protein coupled receptor; KOP, kappa opioid receptor; MOP, mu opioid receptor; NOP, N/OFQ opioid peptide receptor; N/OFQ, nociceptin/orphaninFQ; L-DOPA, levodopa/l-dihydroxyphenylalanine; LID, levodopa-induced dyskinesia.



## TOC GRAPHIC

