

CLINICAL SCIENCE

Is treat-to-target really working in rheumatoid arthritis? a longitudinal analysis of a cohort of patients treated in daily practice (RA BIODAM)

Sofia Ramiro ,^{1,2} Robert BM Landewé,^{2,3} Désirée van der Heijde ,¹ Alexandre Sepriano ,^{1,4} Oliver FitzGerald,⁵ Mikkel Ostergaard,⁶ Joanne Homik,⁷ Ori Elkayam,⁸ J Carter Thorne,⁹ Margaret Larche,¹⁰ Gianfranco Ferracioli,¹¹ Marina Backhaus,¹² Gilles Boire ,¹³ Bernard Combe,¹⁴ Thierry Schaefferbeke,¹⁵ Alain Sarau ,¹⁶ Maxime Dougados,¹⁷ Maurizio Rossini ,¹⁸ Marcello Govoni,¹⁹ Luigi Sinigaglia,²⁰ Alain G Cantagrel,²¹ Cornelia F Allaart,¹ Cheryl Barnabe,²² Clifton O Bingham ,²³ Paul P Tak ,^{3,24,25} Dirkjan van Schaardenburg,³ Hilde Berner Hammer,²⁶ Rana Dadashova,²⁷ Edna Hutchings,²⁷ Joel Paschke,²⁸ Walter P Maksymowich ²⁸

Handling editor Josef S Smolen

For numbered affiliations see end of article.

Correspondence to

Dr Sofia Ramiro, Rheumatology, Leiden University Medical Center, Leiden 2333 ZA, The Netherlands; sofiaramiro@gmail.com

Received 12 December 2019
Revised 25 January 2020
Accepted 27 January 2020
Published Online First
24 February 2020

ABSTRACT

Objectives To investigate whether following a treat-to-target (T2T)-strategy in daily clinical practice leads to more patients with rheumatoid arthritis (RA) meeting the remission target.

Methods RA patients from 10 countries starting/ changing conventional synthetic or biological disease-modifying anti-rheumatic drugs were assessed for disease activity every 3 months for 2 years (RA BIODAM (BIOmarkers of joint DAMAge) cohort). Per visit was decided whether a patient was treated according to a T2T-strategy with 44-joint disease activity score (DAS44) remission (DAS44 <1.6) as the target. Sustained T2T was defined as T2T followed in ≥ 2 consecutive visits. The main outcome was the achievement of DAS44 remission at the subsequent 3-month visit. Other outcomes were remission according to 28-joint disease activity score-erythrocyte sedimentation rate (DAS28-ESR), Clinical Disease Activity Index (CDAI), Simplified Disease Activity Index (SDAI) and American College of Rheumatology/ European League Against Rheumatism (ACR/EULAR) Boolean definitions. The association between T2T and remission was tested in generalised estimating equations models.

Results In total 4356 visits of 571 patients (mean (SD) age: 56 (13) years, 78% female) were included. Appropriate application of T2T was found in 59% of the visits. T2T (vs no T2T) did not yield a higher likelihood of DAS44 remission 3 months later (OR (95% CI): 1.03 (0.92 to 1.16)), but sustained T2T resulted in an increased likelihood of achieving DAS44 remission (OR: 1.19 (1.03 to 1.39)). Similar results were seen with DAS28-ESR remission. For more stringent definitions (CDAI, SDAI and ACR/EULAR Boolean remission), T2T was consistently positively associated with remission (OR range: 1.16 to 1.29), and sustained T2T had a more pronounced effect on remission (OR range: 1.49 to 1.52).

Conclusion In daily clinical practice, the correct application of a T2T-strategy (especially sustained T2T) in patients with RA leads to higher rates of remission.

Key messages

What is already known about this subject?

- ▶ Randomised controlled trials have demonstrated the efficacy of treat-to-target approaches in rheumatoid arthritis. Real life data from cohorts are still needed to support the widespread implementation of treat-to-target (T2T) in clinical practice.

What does this study add?

- ▶ In daily clinical practice, the correct application of a T2T-strategy in patients with rheumatoid arthritis (RA) leads to higher rates of remission as compared with not following it.
- ▶ Not only in early RA, but also in established RA, following a T2T-strategy leads to higher remission rates.

How might this impact on clinical practice or future developments?

- ▶ Rheumatologists should be encouraged to follow a T2T-strategy to contribute to the achievement of higher rates of remission for their patients.

INTRODUCTION

Early diagnosis, prompt commencement of disease modifying anti-rheumatic drug (DMARD) treatment and applying treat-to-target (T2T) strategies are now engrained in rheumatoid arthritis (RA) treatment paradigms. These approaches have substantially improved the outcomes of patients with RA.¹ Remission has been defined and agreed on as the optimal target when managing a patient with RA.^{2,3} Reaching the state of remission is associated with reduced radiographic progression and improved functional ability.⁴

Thoroughly monitoring disease activity, adjusting treatment according to a fixed protocol and aiming



© Author(s) (or their employer(s)) 2020. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Ramiro S, Landewé RBM, van der Heijde D, *et al.* *Ann Rheum Dis* 2020;**79**:453–459.

at a predefined treatment goal, the so-called T2T-strategy, has advantages over usual care.^{5,6} Several strategy studies provide the basis of this evidence, namely the TICORA (Tight Control of RA study)⁷ and CAMERA (Computer Assisted Management in Early RA)⁸ studies. Subsequently, several strategy studies have incorporated a T2T-strategy in their treatment algorithm in the formal comparison of specific therapies, such as was done in the BeSt (Behandel Strategiën) study.⁹ However, such evidence was gathered in the setting of randomised controlled trials (RCTs), with strict inclusion and exclusion criteria, following strict protocols and all particularities of RCTs. These studies provide the best evidence for the efficacy of T2T as an intervention, but to some extent compromise the generalisability of the findings, when one wants to consider applying them more broadly.

Having formally demonstrated the efficacy of T2T in RCTs, it is important to assess whether this strategy also improves outcomes in unselected patients from daily clinical practice. The first cohort studies focussed on patients with very early disease and confirmed that following a standardised intensive treatment led to improved achievement of remission.¹⁰ Subsequently, some cohort studies have shown that tight-control treatment leads to more rapid remission and higher remission achievement after 1 or 2 years than usual care.^{11,12} Nevertheless, the conclusions from these two studies were based on an indirect comparison between two different cohorts (one with T2T applied and another with usual care), with different patient characteristics, and focussed on the remission achievement at 1 or 2 years in the two cohorts. Such a comparison should ideally be made within the same cohort of patients, wherein some patients receive a T2T-strategy while others receive usual care. Real life data from cohorts without strict protocol specifications regarding choice of treatment are still needed to support the widespread implementation of T2T in clinical practice. Furthermore, previous studies have focussed on the achievement of remission at a given time point, for example, 1 or 2 years, ignoring whether or not the remission outcome was achieved in each of the visits throughout the follow-up (eg, three monthly visits, per T2T recommendations). A true longitudinal analysis taking all observations over time into account, both in terms of following T2T or not, and achieving remission or not, reflecting daily clinical practice, has not yet been conducted. Additionally, T2T has not yet been investigated in patients with established RA.

The aim of the present study was to investigate whether following a T2T-strategy leads to more patients with RA meeting the treatment target (remission) in daily clinical practice.

METHODS

Study population

Patients from RA BIODAM (BIOmarkers of joint DAMage), which has been previously described, were included.¹³ In brief, RA BIODAM is a 2-year multinational prospective observational study, including patients with a clinical diagnosis of RA and also fulfilling the 2010 RA Classification Criteria,¹⁴ recruited in daily practice from 10 countries from October 2011 to April 2015. To be eligible, patients presented with active disease (44-joint disease activity score, DAS44 >2.4)¹⁵ and were to be started on or changing DMARD treatment, including conventional synthetic DMARDs (csDMARDs) and a first tumour necrosis factor inhibitor (TNFi); patients who had prior biological DMARD (bDMARD) experience were excluded. All patients were included in this analysis. The database used for this analysis was locked in April 2017. The study fulfilled Good Clinical Practice Guidelines and all patients provided informed consent.

Remission

Remission was the outcome of interest. According to the study protocol, patients were monitored every 3 months using DAS44 calculated with the erythrocyte sedimentation rate (ESR).¹⁵ DAS44 remission, that is, DAS44 <1.6¹⁶ was therefore chosen as the main outcome for this analysis. Alternative definitions of remission were also used, namely the 28-joint disease activity score¹⁷ (DAS28-ESR) remission (ie, DAS28-ESR <2.6),¹⁸ the Clinical Disease Activity Index (CDAI) remission (ie, CDAI ≤2.8),¹⁹ the Simplified Disease Activity Index (SDAI) remission (SDAI ≤3.3)²⁰ and the American College of Rheumatology/European League Against Rheumatism (ACR/EULAR) Boolean remission (ie, tender joint count (TJC) ≤1, swollen joint count (SJC) ≤1, C-reactive protein (CRP) ≤1 mg/dL and patient global assessment (PGA) (0 to 10) ≤1).² All definitions of remission were binary (yes/no).

Treat-to-target

Participating rheumatologists were required by protocol to follow a T2T-strategy with DAS44 remission (DAS44 <1.6) as benchmark. In order to define whether a T2T-strategy was appropriately followed or not, every visit was checked according to predefined criteria. T2T was considered appropriate: (i) if a patient had already a disease activity score below the target (DAS <1.6) and treatment was not intensified; or (ii) if treatment was intensified on a DAS ≥1.6. Treatment intensification was defined as increasing dosage or adding a drug from the following categories: csDMARDs, bDMARDs or corticosteroids. T2T was considered incorrectly applied if: (i) the target was met and treatment was nevertheless intensified; or (ii) the target was not met and treatment was not intensified.

Additional definitions for T2T were also considered for sensitivity analyses: (i) T2T without corticosteroids, that is, without considering corticosteroids as a treatment intensification; (ii) T2T less strict, that is, considering T2T as adequate as long as the target, DAS44 remission, is met, regardless of whether treatment is nevertheless intensified or not; (iii) T2T-low disease activity (T2T-LDA) using LDA (ie, DAS <2.4)²¹ instead of remission as the benchmark.

Furthermore, 'sustained T2T' strategy was defined as following T2T in at least two consecutive visits.

Other relevant clinical information

Age, gender, disease duration, rheumatoid factor (RF) and anti-citrullinated protein antibodies (ACPA) status (positive/negative) and being DMARD-naïve (yes vs no), all collected at baseline, were considered in this analysis as potential effect modifiers or confounders of the relationship of interest. Country of residence was also considered as a potential confounder.

Statistical analysis

The relationship between following T2T at a given visit and meeting the target of remission at the subsequent visit 3 months later was investigated using time-lagged generalised estimating equations (GEE) models. GEE is a suitable technique to make use of all available observations from each patient while adjusting for inherent within-subject correlations of the repeated measurements. Models were time-lagged to allow investigation of the effect of the main predictor of interest (ie, following T2T) on the outcome (ie, remission) with a lag of 3 months; in other words, with the outcome occurring 3 months later. The same analyses were conducted to investigate the effect of sustained T2T on meeting the target of remission. The 'exchangeable' working correlation structure, demonstrating the best fit to the data, was used.

Table 1 Baseline characteristics

	n=571 mean (SD) or n (%)
Age, years	55.7 (12.9)
Female gender	434 (76.0%)
Disease duration, years	6.5 (8.0)
Education, years	12.6 (3.8)
Number of comorbidities	1.2 (1.3)
Rheumatoid factor positivity	370 (68.0%)
Anti-CCP positivity	388 (69.3%)
RF and/or anti-CCP positivity	431 (77.7%)
DAS44 (0–10)	3.8 (1.0)
DAS28-ESR (0–10)	5.2 (1.2)
CDAI (0–76)	26.9 (11.6)
SDAI (0–86)	28.5 (12.4)
Patient global (0–10)	5.7 (2.3)
HAQ (0–3)	1.1 (0.7)
SJC (0–44)	8.4 (6.1)
TJC (0–53)	13.6 (9.1)
ESR (mm/h)	28.7 (22.2)
CRP (mg/dL)	1.5 (2.3)
Number of prior DMARDs	0.9 (1.1)
DMARD naïve	274 (48.0%)
Smoking status	
Never smoker	282 (49.4%)
Current smoker	161 (28.2%)
Ex-smoker	128 (22.4%)
Treatment csDMARD/TNFi started at baseline	
Both	196 (34.6%)
csDMARD only	334 (58.9%)
TNFi only	36 (6.3%)
None	1 (0.2%)
Treatment with oral corticosteroids started at baseline	255 (45%)

anti-CCP, anti-citrullinated protein; CDAI, Clinical Disease Activity Index; CRP, C-reactive protein; csDMARD, conventional synthetic DMARD; DAS44, 44-joint disease activity score; DAS28-ESR, 28-joint disease activity score (with ESR); DMARD, disease modifying anti-rheumatic drug; ESR, erythrocyte sedimentation rate; HAQ, Health Assessment Questionnaire; RF, rheumatoid factor; SDAI, Simple Disease Activity Index; SJC, swollen joint count; TJC, tender joint count; TNFi, tumour necrosis factor inhibitor.

As treatment intensification has a central role in T2T, we sought to investigate the extent to which the components of the disease activity scores contributed to it. We therefore investigated the effect of TJC >1, SJC >1, PGA >1 and CRP >1 mg/dL on treatment intensification (yes/no). This analysis was also conducted with GEE, including all above-mentioned disease activity components in one multivariable model.

For each model, interactions between the T2T variable and age, gender, disease duration and RF/ACPA positivity were tested, and if significant ($p < 0.15$) and clinically relevant the model was fitted in each subgroup. If these proved to be not relevant, final models were adjusted for potential confounders selected a priori: age, gender, disease duration and country of residence. Stata/SE V.12 was used.

RESULTS

In total, 571 patients were included with a mean age of 56 (SD 13) years, 78% females and a mean disease duration of 6.5 (8.0) years, 37% with a disease duration up to 2 years (table 1). In total, 78% of the patients were RF and/or ACPA positive, and 48% were DMARD-naïve at baseline (mean disease duration of 3.6 (5.6), 50% with ≤ 2 year disease duration). At the end of the baseline visit, almost 60% of the patients were on treatment with csDMARDs only, 35% of the patients on a TNFi with a csDMARD and only 6% on TNFi monotherapy. Almost half of the patients were on corticosteroids after the baseline visit.

T2T was appropriately applied in 59% of 4356 visits. This included 31% of patient visits where DAS44 remission was met and treatment was not intensified, and 28% of visits where treatment was appropriately escalated. In 3% of visits (9% of those with treatment intensification), treatment intensification took place even though DAS44 remission was met (making a total of 31% of the visits with treatment intensification). In the remaining 38% of visits T2T was not being followed as there was no treatment intensification despite active disease (DAS44 ≥ 1.6) (figure 1).

Throughout the 2-year follow-up period an increasing proportion of patients met remission definitions. At 3 months 24% of the patients were in DAS44 and DAS28-ESR remission, and 8% in ACR/EULAR Boolean remission. At 24 months 52% of the patients were in DAS44 remission, also 52% in DAS28-ESR remission and 27% in ACR/EULAR Boolean remission (figure 2).




Figure 1 Proportion of the visits (n=4356) in which treat-to-target strategy (with DAS44 <1.6 as benchmark) is followed versus not and the details regarding the proportion of visits with target achievement and/or treatment intensification. Treatment intensification was defined as start or dosage increase of a conventional synthetic or biological disease modifying anti-rheumatic drug or of a corticosteroid. DAS44: 44-joint disease activity score.




Figure 2 Proportion of achievement of the different remission outcomes throughout the 2-year follow-up. ACR, American College of Rheumatology; CDAI, Clinical Disease Activity Index; DAS28, 28-joint disease activity score; DAS44, 44-joint disease activity score; EULAR, European League Against Rheumatism; SDAI, Simplified Disease Activity Index (SDAI).

T2T on remission outcomes

Following a T2T-strategy, as compared with not following it, was not significantly associated with a DAS44 or DAS28-ESR remission 3 months later (OR (95% CI) 1.03 (0.92 to 1.16) and 1.03 (0.91 to 1.16), respectively), but was significantly associated with ACR/EULAR Boolean remission (OR 1.16 (1.01 to 1.34)) and also with CDAI remission (OR 1.29 (1.12 to 1.49)) and SDAI remission (OR 1.24 (1.08 to 1.41)) (table 2). Results of the sensitivity analyses were similar, except for a slightly stronger association between T2T and remission (REM) outcomes for both ‘T2T without corticosteroids’ and ‘T2T-REM less strict’. With T2T-LDA, with LDA as the benchmark, there was a significant association between T2T and all remission outcomes (OR between 1.3 and 1.4). None of the tested effect modifiers, namely age, gender, disease duration, seropositivity or DMARD naïve (vs not), modified the relationships of interest.

Sustained T2T on remission outcomes

Following a sustained T2T-strategy compared with not following it was associated with remission 3 months later according to all definitions, for example, DAS44 remission OR 1.19 (1.03 to 1.39) or ACR/EULAR Boolean remission (OR 1.49 (1.24 to 1.81)) (table 3).

Relationship between disease activity components and treatment intensification

All disease activity components were significantly associated with treatment intensification, with SJC and TJC showing the strongest associations, also in a multivariable model including all the components: OR ‘SJC >1’ 3.42 (2.89 to 4.05), OR ‘TJC >1’ 3.35 (2.72 to 4.11), OR ‘PGA >1’ 2.14 (1.71 to 2.68) and OR ‘CRP >1’ 2.00 (1.66 to 2.42).

Table 2 Effect of following treat-to-target strategies on remission outcomes 3 months later*

	DAS44 remission (OR (95% CI))	DAS28-ESR remission (OR (95% CI))	ACR/EULAR Boolean remission (OR (95% CI))	CDAI remission (OR (95% CI))	SDAI remission (OR (95% CI))
T2T	1.03 (0.92 to 1.16)	1.03 (0.91 to 1.16)	1.16 (1.01 to 1.34)	1.29 (1.12 to 1.49)	1.24 (1.08 to 1.41)
T2T without corticosteroids	1.07 (0.95 to 1.20)	1.11 (0.98 to 1.26)	1.23 (1.06 to 1.42)	1.37 (1.18 to 1.59)	1.34 (1.17 to 1.53)
T2T-REM less strict	1.06 (0.94 to 1.19)	1.07 (0.95 to 1.21)	1.32 (1.13 to 1.53)	1.43 (1.22 to 1.67)	1.34 (1.17 to 1.54)
T2T-LDA	1.26 (1.10 to 1.44)	1.36 (1.17 to 1.56)	1.27 (1.09 to 1.47)	1.39 (1.18 to 1.64)	1.36 (1.17 to 1.59)

T2T without corticosteroids: without considering corticosteroids in treatment intensification. T2T-REM less strict: considering T2T as adequate as long as the target, DAS44 remission, is met, regardless of whether treatment nevertheless intensified or not.

*All models adjusted for age, gender, disease duration and country. T2T was considered being followed: (i) if a patient had already a disease activity score below the target (DAS <1.6; DAS <2.4 for LDA definition) and treatment was correctly not intensified; or (ii) if treatment was intensified on a DAS ≥1.6 (or DAS ≥2.4 for LDA definition).

ACR, American College of Rheumatology; CDAI, Clinical Disease Activity Index; DAS44, 44-joint disease activity score; DAS28-ESR, 28-joint disease activity score (with ESR); ESR, erythrocyte sedimentation rate; EULAR, European League Against Rheumatism; SDAI, Simple Disease Activity Index; T2T, treat-to-target; T2T-LDA, T2T-low disease activity; T2T-REM, T2T-remission.

Table 3 Effect of following a sustained treat-to-target strategy on remission outcomes 3 months later*

	DAS44 sustained remission (OR (95% CI))	DAS28-ESR sustained remission (OR (95% CI))	ACR/EULAR Boolean sustained remission (OR (95% CI))	CDAI sustained remission (OR (95% CI))	SDAI sustained remission (OR (95% CI))
Sustained T2T	1.19 (1.03 to 1.39)	1.23 (1.06 to 1.44)	1.49 (1.24 to 1.81)	1.45 (1.19 to 1.77)	1.52 (1.27 to 1.82)

*All models adjusted for age, gender, disease duration and country. Sustained treat-to-target was considered followed if T2T was followed in ≥ 2 subsequent visits. T2T was considered being followed: (i) if a patient had already a disease activity score below the target (DAS <1.6; DAS <2.4 for LDA definition) and treatment was correctly not intensified; or (ii) if treatment was intensified on a DAS ≥ 1.6 (or DAS ≥ 2.4 for LDA definition).

ACR, American College of Rheumatology; CDAI, Clinical Disease Activity Index; DAS44, 44-joint disease activity score; DAS28-ESR, 28-joint disease activity score; ESR, erythrocyte sedimentation rate; EULAR, European League Against Rheumatism; LDA, low disease activity; SDAI, Simple Disease Activity Index; T2T, treat-to-target.

DISCUSSION

In the present study we have shown that following a T2T-strategy, and particularly sustained T2T, in daily clinical practice leads to more patients with RA meeting the most stringent remission criteria over time. This is the first comprehensive analysis that considers all available visits of unselected patients who were followed by protocol for a period of 2 years. The results of the analysis provide direct evidence that following T2T, and particularly sustained T2T, immediately results in a higher likelihood of remission at the next visit, 3 months later (the longitudinal interpretation of a T2T-strategy). Moreover, we have for the first time shown that following T2T is also efficacious in patients with established RA, while previous studies focussed on the effect of T2T in patients with early RA.

The strictly temporal relationship between following a T2T-strategy and meeting remission was statistically significant for almost all remission outcomes and for the different T2T definitions used. The exceptions were the DAS44 and DAS28-ESR remission definitions with an interval of 3 months only, while for sustained T2T the relationship with all remission outcomes was statistically significant. The explanation is rather technical: the independent variable (T2T with DAS44 as benchmark) and the outcome (ie, DAS44 remission) include exactly the same disease activity score, which implies that the model becomes inherently auto-regressive. Such a scenario effectively removes the variability in the data necessary to demonstrate efficacy of an intervention. The other definitions of remission are slightly different from the benchmark definition and allow more statistical separation. An alternative explanation is that DAS44 and DAS28-ESR definitions are more lenient in comparison to ACR/EULAR Boolean, CDAI and SDAI remission and are more frequently met even if T2T is not applied.² Nevertheless, the signal that a T2T-strategy, and particularly sustained T2T-strategy, increases the likelihood of stringent remission is clear and consistent. Also, these findings became even more evident throughout the follow-up of this study. The proportion of patients achieving remission, regardless of its definition, increased substantially through follow-up (figure 2). Even after 2 years, a plateau has not yet been reached, reassuring clinicians that if we measure disease activity and treat patients effectively over time, high remission rates can be achieved.

These findings come from a population of patients with an average of 6.5 years of disease duration. One may speculate that the effect of following T2T could be even better in early disease. In this study, we have not found any differences between patients DMARD naïve versus not and also according to disease duration, but a lack of statistical power cannot be excluded. Additionally, even patients who were DMARD naïve had a relatively long disease duration (average of 3.6 years), not being the most representative DMARD naïve patients.

If T2T is so clearly associated with clinical remission, as shown here and in the literature,^{5,6} why, then, is a T2T-strategy not always followed in clinical practice? Even in this study, with a protocol requiring implementation of T2T, this strategy was 'only' followed in less than two-thirds of the visits. Also within the RA BIODAM cohort, we have shown that, among other factors, the absence of objective signs of inflammation (eg, swollen joints) implied a lower likelihood to follow T2T.²² Also, in the 10-year follow-up of the BeSt trial, non-adherence to the protocol has been assigned to disagreement with how DAS reflects disease activity (felt to overestimate the real disease activity) and disagreement with the subsequently required step in the protocol.²³ Many clinicians find regularly measuring disease activity too time consuming endeavour and consider it an additional barrier to implementation of T2T.^{24,25}

In order to launch new strategies or interventions in clinical practice, the formulation of recommendations, like the T2T recommendations,²⁶ does not suffice and implementation should actively be promoted. Studies like ours may further corroborate the message that T2T leads to more stringent remission and may help implementation in clinical practice. Appropriate education may also help. The intervention of the TRACTION trial included one educational face-to-face meeting and monthly webinars on the principles and practical advice on implementation of T2T. A substantial improvement in the adherence to T2T was demonstrated with improvement of 46% in the arm following the training programme compared with 14% in the control arm.²⁷ Still, rheumatologists may report compliance with recommendations but in practice do not always follow them.²⁸

Some limitations of this study need to be considered. First, it is designed as an observational study reflecting daily clinical practice with unselected patients contrasting with the reality of RCTs from which most evidence on T2T originates to date. However, one may question how close to daily clinical practice the RA BIODAM cohort really is, with participation from only a few centres per country, several being tertiary referral centres, and with rheumatologists mandated to follow a strict T2T protocol. As in principle, rheumatologists were required to follow T2T per protocol, we have in this study in essence compared the visits in which the protocol was followed to others in which protocol was violated. One can therefore not exclude a bias intrinsic to this comparison. Additionally, detailed reasons for not following T2T have not been adequately registered precluding additional analysis of adherence to T2T versus taking the physician's reasoning into account. Moreover, only patients with active disease were included, and the average baseline disease activity was high. This may preclude the generalisability of the findings to patients with low disease activity, and not answer the question of whether following a T2T-strategy is beneficial in patients

already in low disease activity, given the risks of overtreatment.^{29,30} Lastly, when investigating the impact of following a T2T-strategy, one is not only analysing the impact of treatment intensification but implicitly one is evaluating visits in which patients are already in remission, which have accentuated the benefit of T2T. However, it was our aim to investigate the impact of following the T2T-strategy in its whole and not parts of it, as well as to take all disease activity measurements into account as the longitudinal technique chosen properly does. As a main strength, this is a multinational observational study, including unselected patients reflecting daily clinical practice, with the first truly longitudinal analysis addressing the impact of following a (sustained) T2T-strategy.

In conclusion, following a T2T-strategy, and especially sustained T2T, works in daily clinical practice and leads to more patients meeting the target, that is, remission. Rheumatologists should be encouraged to follow a T2T-strategy to contribute to the achievement of higher rates of remission for their patients.

Author affiliations

¹Department of Rheumatology, Leiden University Medical Center, Leiden, The Netherlands

²Department of Rheumatology, Zuyderland Medical Center, Heerlen, The Netherlands

³Department of Rheumatology, Amsterdam Rheumatology Center, Amsterdam, The Netherlands

⁴NOVA Medical School, Universidade Nova de Lisboa, Lisbon, Portugal

⁵St Vincent's University Hospital and Conway Institute for Biomolecular Research, University College Dublin, Dublin, Ireland

⁶Copenhagen Center for Arthritis Research, Center for Rheumatology and Spine Diseases, Rigshospitalet, Glostrup and Department of Clinical Medicine, University of Copenhagen, Copenhagen, Denmark

⁷Department of Medicine, University of Alberta, Edmonton, Alberta, Canada

⁸Tel Aviv Sourasky Medical Center and the "Sackler" Faculty of Medicine, Tel Aviv University, Tel Aviv, Israel

⁹Southlake Regional Health Centre, University of Toronto, Toronto, Ontario, Canada

¹⁰Departments of Medicine and Pediatrics, Divisions of Rheumatology, Clinical Immunology and Allergy, McMaster University, Hamilton, Ontario, Canada

¹¹Catholic University of the Sacred Heart, Roma, Italy

¹²Park-Klinik Weissensee, Academic Hospital of the Charité, Berlin, Germany

¹³Department of Medicine/Division of Rheumatology, Centre intégré universitaire de santé et de services sociaux de l'Estrie – Centre hospitalier universitaire de Sherbrooke (CIUSSS de l'Estrie-CHUS), Université de Sherbrooke, Sherbrooke, Quebec, Canada

¹⁴CHU Montpellier and Montpellier University, Montpellier, France

¹⁵Department of Rheumatology, FHU ACRONIM, University Hospital of Bordeaux, University of Bordeaux, Bordeaux, France

¹⁶Rheumatology, CHU Brest, Brest, France

¹⁷Rheumatology Department, Paris Descartes University, Cochin Hospital, Assistance Publique-Hôpitaux de Paris, INSERM (U1153): Clinical Epidemiology and Biostatistics, PRES Sorbonne Paris-Cité, Paris, France

¹⁸Rheumatology Unit, Department of Medicine, University of Verona, Verona, Italy

¹⁹Rheumatology Unit, S. Anna Hospital and University of Ferrara, Ferrara, Italy

²⁰Department of Rheumatology, Gaetano Pini Institute, Milan, Italy

²¹Department of Rheumatology, Paul Sabatier University, Toulouse, France

²²Departments of Medicine and Community Health Sciences, University of Calgary, Calgary, Alberta, Canada

²³Johns Hopkins University, Baltimore, Maryland, USA

²⁴Department of Rheumatology, Ghent University, Ghent, Belgium

²⁵Department of Medicine, Cambridge University, Cambridge, United Kingdom

²⁶Diakonhjemmet Hospital, Oslo, Norway

²⁷CaRE Arthritis LTD, Edmonton, Alberta, Canada

²⁸CaRE Arthritis LTD, University of Alberta, Edmonton, Alberta, Canada

Acknowledgements The authors sincerely thank the staff and all patients of the RA BIODAM study.

Contributors All authors made contributions to conception and/or implementation of the study, were involved in reviewing and revising the manuscript and gave final approval to the version to be published.

Funding BIODAM was financially supported by an unrestricted grant from AbbVie. AS is supported by a doctoral grant from 'Fundação para a Ciência e Tecnologia' (SFRH/BD/108246/2015).

Competing interests SR: Received research grants and/or consultancy fees from AbbVie, Eli Lilly, MSD, Novartis and Sanofi. RL: Received consulting fees from AbbVie, BMS, Celgene, Eli-Lilly, Galapagos, Gilead, Glaxo-Smith-Kline, Janssen, Merck, Novartis, Pfizer, Roche, UCB and is Director of Rheumatology Consultancy BV. DvdH: Received consulting fees from AbbVie, Amgen, Astellas, AstraZeneca, BMS, Boehringer Ingelheim, Celgene, Daiichi, Eli-Lilly, Galapagos, Gilead, Glaxo-Smith-Kline, Janssen, Merck, Novartis, Pfizer, Regeneron, Roche, Sanofi, Takeda, UCB Pharma and is Director of Imaging Rheumatology BV. AS: Received speaking fees from Novartis. OFG: Received research grants and/or consultancy fees from AbbVie, Eli Lilly, Celgene, Novartis, UCB, Pfizer, Amgen and Janssen. MØ: Received research support, consultancy fees and/or speaker fees from Abbvie, BMS, Boehringer-Ingelheim, Celgene, Eli-Lilly, Hospira, Janssen, Merck, Novartis, Novo, Orion, Pfizer, Regeneron, Roche, Sandoz, Sanofi and UCB. OE: Received grants and personal fees from Pfizer, Abbvie, Roche, Janssen, Novartis and Lilly. JCT: Received research grants and/or consulting fees from AbbVie, Amgen, Celgene, Centocor, Lilly, Medexus/Medac, Merck, Novartis, Pfizer, Sandoz and Sanofi. ML: Received advisory board honoraria from AbbVie, Actelion, Boehringer Ingelheim, BMS, Janssen, Novartis, Pfizer, Sanofi, Roche and an unrestricted educational grant from AbbVie. GF: Received research grants and/or consultancy fees from Novartis, Roche, BMS, Pfizer, AbbVie, Lilly, Janssen, MSD, Boehringer-Ingelheim and UCB. MB: Received research grants from AbbVie Co.KG, Roche, Novartis and Pfizer; received speaking fees from Actelion, AbbVie Co.KG, BMS, Celgene, Gilead, Janssen, Lilly, MSD, Novartis, Sanofi Genzyme, Roche, Pfizer and UCB. GB: Received consulting fees from Eli Lilly, Janssen, Novartis and Pfizer; received speaker fees from BMS, Merck and Pfizer. Industry support for investigator-initiated research initiatives from Abbvie, Amgen, BMS, Celgene, Eli Lilly, Janssen, Merck, Pfizer, Roche and UCB. BC: Received consulting fees from AbbVie, BMS, Eli-Lilly, Gilead, Janssen, Merck, Novartis, Pfizer, Roche-Chugai, Sanofi and UCB. TS: Received consulting fees from Lilly, Novartis, Pfizer and Sanofi; received research financial support from Pfizer and Lilly. AS: Received research grants and/or consultancy fees from AbbVie, BMS, Chugai, Eli Lilly, MSD, Nordic pharma, Novartis, Pfizer, Sanofi and UCB. MD: Received research grants and/or honorarium fees for his participation at advisory boards and/or symposium organised by Pfizer, AbbVie, Lilly, Novartis, BMS, Roche, UCB and Merck. MR: Received advisory board honoraria, consultancy fees and/or speaker fees from Abiogen, Amgen, Abbvie, BMS, Celgene, Eli-Lilly, MSD, Novartis, Pfizer, Sanofi, Sandoz and UCB. MG: Received advisory board honoraria, consultancy fees and/or speaker fees from Abbvie, Alfa-Sigma, BMS, Celgene, MSD, Novartis, Pfizer, Roche and Sanofi. LS: Received speaker fees from Amgen, Eli Lilly, Abbvie, Roche and BMS. AGC: Received consultancy fees from BMS, Chugai, Janssen, Lilly, MSD, Novartis, Pfizer, Roche, Sanofi, UCB, AbbVie, Celgene and Nordic Pharma; received research grants from Pfizer, UCB, AbbVie and Celgene. CB: Received advisory board honoraria from Gilead, Pfizer, Novartis, Eli Lilly, Roche and Amgen; speaking fees from UCB, BMS and Abbvie. COB III: Consultant to Abbvie, BMS, Gilead, Lilly, Pfizer, Genentech/Roche and Sanofi/Regeneron. Grant support from BMS and Janssen. PPT: Currently an employee of Kintai Therapeutics, Cambridge MA. Kintai Therapeutics has not been involved in this work. HBH: Received fees for speaking and/or consulting from AbbVie, BMS, Pfizer, UCB, Roche, MSD and Novartis. WPM: Received consulting fees from AbbVie, Boehringer Ingelheim, Celgene, Eli-Lilly, Galapagos, Janssen, Novartis, Pfizer and UCB Pharma and is Chief Medical Officer of CARE Arthritis Limited.

Patient consent for publication Not required.

Ethics approval The study received ethical approval from the local ethics committees.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data may be obtained from a third party and are not publicly available.

ORCID iDs

Sofia Ramiro <http://orcid.org/0000-0002-8899-9087>

Désirée van der Heijde <http://orcid.org/0000-0002-5781-158X>

Alexandre Sepriano <http://orcid.org/0000-0003-1954-0229>

Gilles Boire <http://orcid.org/0000-0003-2481-5821>

Alain Saraux <http://orcid.org/0000-0002-8454-7067>

Maurizio Rossini <http://orcid.org/0000-0001-9692-2293>

Clifton O Bingham <http://orcid.org/0000-0002-4752-5029>

Paul P Tak <http://orcid.org/0000-0002-3532-5409>

Walter P Maksymowych <http://orcid.org/0000-0002-1291-1755>

REFERENCES

- Kievit W, Fransen J, de Waal Malefijt MC, *et al.* Treatment changes and improved outcomes in RA: an overview of a large inception cohort from 1989 to 2009. *Rheumatology* 2013;52:1500–8.
- Felson DT, Smolen JS, Wells G, *et al.* American College of Rheumatology/European League against rheumatism provisional definition of remission in rheumatoid arthritis for clinical trials. *Ann Rheum Dis* 2011;70:404–13.

- 3 Smolen JS, Landewé R, Bijlsma J, *et al.* EULAR recommendations for the management of rheumatoid arthritis with synthetic and biological disease-modifying antirheumatic drugs: 2016 update. *Ann Rheum Dis* 2017;76:960–77.
- 4 van Tuyl LHD, Felson DT, Wells G, *et al.* Evidence for predictive validity of remission on long-term outcome in rheumatoid arthritis: a systematic review. *Arthritis Care Res* 2010;62:108–17.
- 5 Schoels M, Knevel R, Aletaha D, *et al.* Evidence for treating rheumatoid arthritis to target: results of a systematic literature search. *Ann Rheum Dis* 2010;69:638–43.
- 6 Schipper LG, van Hulst LTC, Grol R, *et al.* Meta-Analysis of tight control strategies in rheumatoid arthritis: protocolized treatment has additional value with respect to the clinical outcome. *Rheumatology* 2010;49:2154–64.
- 7 Grigor C, Capell H, Stirling A, *et al.* Effect of a treatment strategy of tight control for rheumatoid arthritis (the TICORA study): a single-blind randomised controlled trial. *The Lancet* 2004;364:263–9.
- 8 Verstappen SMM, Jacobs JWG, van der Veen MJ, *et al.* Intensive treatment with methotrexate in early rheumatoid arthritis: aiming for remission. computer assisted management in early rheumatoid arthritis (camera, an open-label strategy trial). *Ann Rheum Dis* 2007;66:1443–9.
- 9 Goekoop-Ruiterman YPM, De Vries-Bouwstra JK, Allaart CF, *et al.* Clinical and radiographic outcomes of four different treatment strategies in patients with early rheumatoid arthritis (the best study): a randomized, controlled trial. *Arthritis Rheum* 2005;52:3381–90.
- 10 Vermeer M, Kuper HH, Hoekstra M, *et al.* Implementation of a treat-to-target strategy in very early rheumatoid arthritis: results of the Dutch rheumatoid arthritis monitoring remission induction cohort study. *Arthritis Rheum* 2011;63:2865–72.
- 11 Schipper LG, Vermeer M, Kuper HH, *et al.* A tight control treatment strategy aiming for remission in early rheumatoid arthritis is more effective than usual care treatment in daily clinical practice: a study of two cohorts in the Dutch rheumatoid arthritis monitoring registry. *Ann Rheum Dis* 2012;71:845–50.
- 12 Brinkmann GH, Norvang V, Norli ES, *et al.* Treat to target strategy in early rheumatoid arthritis versus routine care - A comparative clinical practice study. *Seminars in arthritis and rheumatism* 2018.
- 13 Maksymowych WP, FitzGerald O, Østergaard M, *et al.* The International RA BIODAM cohort for validation of soluble biomarkers in rheumatoid arthritis: cohort description. *J Rheumatol*. In Press 2019;jrheum.190302.
- 14 Aletaha D, Neogi T, Silman AJ, *et al.* 2010 rheumatoid arthritis classification criteria: an American College of Rheumatology/European League against rheumatism collaborative initiative. *Ann Rheum Dis* 2010;69:1580–8.
- 15 van der Heijde DM, van 't Hof MA, van Riel PL, *et al.* Judging disease activity in clinical practice in rheumatoid arthritis: first step in the development of a disease activity score. *Ann Rheum Dis* 1990;49:916–20.
- 16 Prevoo MLL, van Gestel AM, van T Hof MA, *et al.* Remission in a prospective study of patients with rheumatoid arthritis. American rheumatism association preliminary remission criteria in relation to the disease activity score. *Rheumatology* 1996;35:1101–5.
- 17 Prevoo ML, van 't Hof MA, Kuper HH, *et al.* Modified disease activity scores that include twenty-eight-joint counts. development and validation in a prospective longitudinal study of patients with rheumatoid arthritis. *Arthritis Rheum* 1995;38:44–8.
- 18 Franssen J, Creemers MC, Van Riel PL. Remission in rheumatoid arthritis: agreement of the disease activity score (DAS28) with the Ara preliminary remission criteria. *Rheumatology* 2004;43:1252–5.
- 19 Aletaha D, Smolen J. The simplified disease activity index (SDAI) and the clinical disease activity index (CDAI): a review of their usefulness and validity in rheumatoid arthritis. *Clin Exp Rheumatol* 2005;23:S100–8.
- 20 Aletaha D, Ward MM, Machold KP, *et al.* Remission and active disease in rheumatoid arthritis: defining criteria for disease activity states. *Arthritis Rheum* 2005;52:2625–36.
- 21 van Gestel AM, Prevoo ML, van 't Hof MA, *et al.* Development and validation of the European League against rheumatism response criteria for rheumatoid arthritis. Comparison with the preliminary American College of rheumatology and the world health Organization/International League against rheumatism criteria. *Arthritis Rheum* 1996;39:34–40.
- 22 Sepriano A, Ramiro S, FitzGerald O, *et al.* Adherence to Treat-to-Target management in rheumatoid arthritis and associated factors: data from the International RA BIODAM cohort. *J Rheumatol* 2019;jrheum.190303.
- 23 Markuse IM, Dirven L, Han KH, *et al.* Evaluating adherence to a Treat-to-Target protocol in recent-onset rheumatoid arthritis: reasons for compliance and hesitation. *Arthritis Care Res* 2016;68:446–53.
- 24 Pincus T, Segurado OG. Most visits of most patients with rheumatoid arthritis to most rheumatologists do not include a formal quantitative joint count. *Ann Rheum Dis* 2006;65:820–2.
- 25 Yazici Y, Bergman M, Pincus T. Time to score quantitative rheumatoid arthritis measures: 28-Joint count, disease activity score, health assessment questionnaire (HAQ), multidimensional HAQ (MDHAQ), and routine assessment of patient index data (rapid) scores. *J Rheumatol* 2008;35:603–9.
- 26 Smolen JS, Breedveld FC, Burmester GR, *et al.* Treating rheumatoid arthritis to target: 2014 update of the recommendations of an international Task force. *Ann Rheum Dis* 2016;75:3–15.
- 27 Solomon DH, Losina E, Lu B, *et al.* Implementation of Treat-to-Target in rheumatoid arthritis through a learning collaborative: results of a randomized controlled trial. *Arthritis Rheumatol* 2017;69:1374–80.
- 28 Gvozdenović E, Allaart CF, van der Heijde D, *et al.* When rheumatologists report that they agree with a guideline, does this mean that they practise the Guideline in clinical practice? results of the International recommendation implementation study (iris). *RMD Open* 2016;2:e000221.
- 29 Landewé RBM. Overdiagnosis and overtreatment in rheumatology: a little caution is in order. *Ann Rheum Dis* 2018;77:1394–96.
- 30 Bergstra SA, Olivás O, Akdemir G, *et al.* Further treatment intensification in undifferentiated and rheumatoid arthritis patients already in low disease activity has limited benefit towards physical functioning. *Arthritis Res Ther* 2017;19:220.

T2T in real-life helps people get to remission



Rheumatologists should follow a T2T-strategy in patients with both early and established rheumatoid arthritis

INTRODUCTION

Rheumatoid arthritis is a chronic inflammatory disease that can affect a person's joints, and may cause pain and disability. Rheumatoid arthritis affects people of all ages, and is more common in women than men.

'Treat to target' (often shortened to T2T) has worked well in clinical trials in people with rheumatoid arthritis. When a doctor tries to treat to target, they start by setting a goal. This is often remission – meaning you have no signs or symptoms of active disease. They then closely monitor your disease activity, and adjust treatment according to a set of rules to help you get to the goal. T2T has been tested in clinical trials. Clinical trials are good at finding out specific things, but in real life people are more complicated – and doctors may have less time or resources than in a trial. Clinical trials often exclude people with certain types of disease or treatment history, but in real life it is important to be able to treat everybody.

WHAT DID THE AUTHORS HOPE TO FIND?

The authors looked at T2T in real life to see how well it is being followed. They wanted to see how well it works in real life, and whether people who follow T2T achieve remission more often than people who are not on T2T. The authors also wanted to know whether T2T works in people with established rheumatoid arthritis, not just those who are newly diagnosed.

WHO WAS STUDIED?

The study looked at 571 people with rheumatoid arthritis who had taken part in the RA BIODAM study. People were being treated in clinics across 10 countries. Everyone had active disease, and was about to be started on – or change to – a conventional DMARD medicine or a biologic medicine called a TNF inhibitor. People could not take part if they had tried a biologic medicine before.

HOW WAS THE STUDY CONDUCTED?

This was a longitudinal observational study, which means that the researchers simply observed people in normal clinics and recorded information at several time points. People visited the clinic once every 3 months and had their disease activity measured, and notes made about the treatment they were on. The aim of therapy was remission. The authors looked at people's clinical notes to see whether the doctors had followed the T2T strategy properly. This was measured based on whether people's treatment had been changed at the right time according to their disease activity score. If T2T was properly followed in at least two consecutive visits, this was defined as 'sustained T2T'. The authors then looked to see if there was a relationship between following T2T at a clinic check-up, and meeting the target goal of remission at the next visit 3 months later.

WHAT WERE THE MAIN FINDINGS OF THE STUDY?

The main finding was that a T2T-strategy led to more people achieving the goal of remission. There was evidence that following T2T – and especially sustained T2T – made it more likely that a person would be in remission at the next visit, 3 months later.

ARE THESE FINDINGS NEW?

Yes, this is the first study to look in detail at all available visits of real-life T2T patients over 2 years. It is also the first time that T2T has been shown to work in people with established RA.

WHAT ARE THE LIMITATIONS OF THE STUDY?

There are some limitations. First, the patient information came from RA BIODAM, which was set up in specialist centres, where the doctors were following a set T2T programme. This might not reflect normal everyday

practice for people seen in general rheumatology clinics. Also, where doctors were not following T2T, there were not always notes to explain why. There could have been good medical reasons for choosing a different approach, or it could have been the patient's decision not to change treatment at a particular point. Finally, only people with active disease were included, and the disease activity at the start of the study was high. This means the findings might not apply to people with low disease activity, and so it is not possible to say whether following a T2T-strategy is beneficial for people who are already in low disease activity, or if there is a risk of over-treatment.

WHAT DO THE AUTHORS PLAN ON DOING WITH THIS INFORMATION?

This information will be shared with rheumatologists and patients. Doctors should be encouraged to follow a T2T-strategy to help get people to remission, but it is important to properly assess disease activity and use the information to make treatment decisions.

The authors are planning a follow-up study to see whether following a T2T-strategy helps inhibit joint damage, as seen on an X-ray.

WHAT DOES THIS MEAN FOR ME?

If you have rheumatoid arthritis, using a treat-to-target strategy with your doctor will lead to better outcomes. It is important that your disease activity is measured often, and that you talk to your doctor about the results and any treatment changes you need. Being engaged in the management of your disease, and understanding the benefits of treat-to-target, will help you look after yourself better.

If you have any concerns about your disease or its treatment, you should talk to your doctor. It is important that you do not stop taking any medicine you have been prescribed without getting proper medical advice.

Disclaimer: This is a summary of a scientific article written by a medical professional (“the Original Article”). The Summary is written to assist non medically trained readers to understand general points of the Original Article. It is supplied “as is” without any warranty. You should note that the Original Article (and Summary) may not be fully relevant nor accurate as medical science is constantly changing and errors can occur. It is therefore very important that readers not rely on the content in the Summary and consult their medical professionals for all aspects of their health care and only rely on the Summary if directed to do so by their medical professional. Please view our full Website Terms and Conditions. <http://www.bmj.com/company/legal-information/>

Date prepared: March 2020

Summary based on research article published on: 24 February 2020

From: Ramiro S, *et al.* Is treat-to-target really working in rheumatoid arthritis? A longitudinal analysis of a cohort of patients treated in daily practice (RA BIODAM). *Ann Rheum Dis* 2020;79:453–459. doi:10.1136/annrheumdis-2019-216819

Copyright © 2020 BMJ Publishing Group Ltd & European League Against Rheumatism. Medical professionals may print copies for their and their patients and students non commercial use. Other individuals may print a single copy for their personal, non commercial use. For other uses please contact our [Rights and Licensing Team](#).