

## Issue Highlight

### Issue highlight – July 2018

This July issue of Clinical Cytometry B includes 11 original papers, 6 brief communications, 2 letters to the editor and a review article dealing with cancer-related mRNA expression analysis using a novel flow cytometry-based assay. The flow data from Depreter et al. showed a good correlation with the gold standard, RT-qPCR technique (1). Interestingly, *Wilms' tumor 1* (WT1) gene levels were shown to be significantly higher in AML patient samples with WT1 gene overexpression, previously defined by RT-qPCR. Moreover, *WT1* overexpression was distinguishable between heterogeneous cell populations and was also documented in rare leukemic stem cells (LSC). This study also showed *comparable results using fresh, short-term and long-term cryopreserved samples, thus allowing the analysis of RNA samples on a retrospective basis*. The paper by Depreter et al. provides crucial information concerning the ability to detect single-cell or rare cell population-specific gene expression changes in acute myeloid leukemia (AML) samples. In both adult and pediatric AML, persistence of rare LSCs are thought to be causative for the high relapse rates, therefore, an increasing need for advanced instrumental analytical tools, able to elucidate coding and non-coding gene expressions at single-cell level, has emerged, and this study may help in clarifying this aspect (2–5).

Taking into account that the liquid biopsy is becoming a powerful diagnostic tool in solid tumor diagnosis, it is conceivable that this test may play a role for the detection of cancer cells from a tumor that are circulating in the blood and may help find cancer at an early stage. It may also be used to help plan treatment or to find out how well treatment is working or if cancer has come back (6,7). Furthermore, since accumulating biological evidence support the notion that current chemotherapy drugs will not be effective for leukemic stem cells killing, or at least has the capacity to target both the leukemic and normal stem cell populations, new strategies are required that specifically and preferentially kill the malignant stem cell population, while sparing normal stem cells. The possibility to detect LSC using this novel flow cytometry-based assay may allow the identification of suitable drugs targeting specifically LSC (6–8).

Original articles published in this issue were related to four main fields (1) immunophenotyping of lymphoproliferative disorders or normal cells, (2) the detection of a rare acquired genetic disorder Paroxysmal nocturnal hemoglobinuria (PNH), (3) immunology science with particular emphasis on monocytes, and (4) cell function.

In the paper by Starostka et al., a quantitative assessment of informative immunophenotypic markers in mature CD5-positive B-cell neoplasms was performed,



Francesco Lanza

and data showed that this approach may increase the diagnostic value of immunophenotyping in several chronic lymphoproliferative disorders (9). Based on their study, the most informative markers for the distinction of CLL/SLL, MCL, CD5+ MZL were the MFI values of the following markers: CD79b, CD20, CD23, CD43, CD38, CD11c, FMC7, CD200, kappa light chain, and their combinations. CD23 and CD200 were the most discriminant between CLL/SLL and MCL and CD23 plus CD79b between CLL/SLL and CD5+ MZL. However, this quantitative marker investigation failed to accurately distinguish MCL and CD5+ MZL. I, therefore, believe that this paper highlights the data mining methods for the analysis and selection of the most informative immunophenotypic markers and may help designing a predictive model, possibly minimizing the subjectivity of expert based assessment (10–12).

\*Correspondence to: Francesco Lanza, Hematology Institute, University Hospital, Ravenna, 48121- Italy. E-mail: francesco.lanza@auslromagna.it

Published online in Wiley Online Library (wileyonlinelibrary.com). DOI: 10.1002/cyto.b.21644

As far as Dr. Auat's paper is concerned, a flow cytometry measurement of CD307a antigen expression in B-cell malignancies and during normal B-cell maturation has been investigated with the main aim to explore the diagnostic role played by this molecule during normal and leukemic differentiation (13). The results indicate that the flow cytometry assessment of CD307a expression could be helpful to distinguish CLL from MCL, and the latter from MZL. Although these results are not entirely conclusive, they provide a basis for further studies in a larger cohort of patients. Interestingly, B-lymphoblasts from acute lymphoblastic leukemia patients exhibited minimal expression of CD307a, displaying a similar expression pattern to that of normal B-cell precursors. This study may be considered in the construction and use of flow cytometric assays for minimal residual disease detection (14-16). The identification of novel antigenic markers to target the neoplastic cells through engineered monoclonal antibodies remains one of the main goals of the most recent treatment modalities, and this paper may provide valuable information in this research field (16).

The second theme covered in this issue is represented by PNH. PNH is an acquired clonal stem cell disorder characterized by intravascular hemolysis, deep venous thrombosis, bone marrow failure and increased susceptibility to infections. PNH is a stem cell disorder caused by a PIG-A gene mutation (17-19). As a consequence, blood cells completely or partially lack surface proteins that are tethered to the membrane through the glycosylphosphatidylinositol (GPI) anchor. The resulting absence of CD55 and CD59 molecules on red cells renders them sensitive to complement-mediated intravascular hemolysis and associated hemoglobinuria. Over the last 10 years, PNH testing by flow cytometry has become a common laboratory test in the clinical practice (20-22).

In the paper published by Blaha et al., authors tested the monoclonal anti-CD157 antibody clone SY11B5, for high sensitivity detection of PNH clones on the leukocyte population (23). Results showed that this reagent fails to detect a common polymorphic variant encoded by BST-1. The failure of anti-CD157 antibody clone SY11B5 to detect a common SNP can explain some CD157-negative cytometric data. This provides crucial knowledge for laboratories performing PNH analyses as such results can potentially lead to false-positive PNH interpretation. Due to the rare incidence of PNH, as well as rare genetic variations such as CD157, we suggest that flow cytometric analysis of PNH cells must not be based on lack of expression of just a single GPI-anchored marker. Rather a consistent deficiency of at least two markers on at least two different cell types is required.

In the second manuscript related to PNH, one of the leading scientist in this field of investigation, Rob Sutherland, compared high sensitivity 5-, 6, 7-color flow cytometry analysis using Becton-Dickinson and Beckman Coulter cytometers (24). Assessment of > 40 PNH samples showed that the FLAER-based data derive virtually identical data to the non-FLAER results for neutrophils ( $R^2 = 1$ ) and monocytes ( $R^2 = 0.9999$ ). Furthermore, both Canto and Navios platforms gave rise similar data

with 7-, 6-, and 5-color versions of the assay. Interestingly, analysis of non-PNH samples confirmed extremely low background rate of PNH phenotypes (neutrophils and monocytes) with all three approaches. In conclusion, this paper describes, for the first time a series of high-sensitivity, single tube flow assays for the simultaneous detection of GPI-deficient leukocyte subsets across both 3-laser Navios and Canto technologies. The 7-color variants of these assays confirm the earlier findings of Marinov et al. that FLAER is not an absolute requirement for high-sensitivity flow assays to detect PNH clones in PNH and related diseases (25). These results may be regarded as a practice changing approach (20-22,26).

Due to its high sensitivity, Lindemann et al. focused on the role played by the flow cytometry crossmatch (FCXM) for identifying an optimal living donor in kidney transplantation (27). In particular, the effect of ABO incompatibility on T cell flow cytometry crossmatch has been investigated, allowing authors to conclude that ABO incompatibility was associated with higher T-FCXM responses, especially in recipients with blood group O. This finding has major impact on the interpretation of flow crossmatch results. Based on these data, authors have postulated that current cut-off values need to be reassessed in the ABO incompatible setting (28).

Regarding Cecilia Langenskiöld's investigation, authors established a flow cytometry antibody panel that can be used to determine granulocytes, monocytes and lymphocyte subset concentrations in fresh and frozen whole blood using TruCount technology (29). Using this method whole-blood samples can be frozen using a simple preparation method, and stored long-term before accurate determination of cell concentration. This allows for standardized analysis of the samples at a reference laboratory in multi-center studies. This methodological improvement may be helpful in routine flow cytometry analyses (30).

Wonner et al. investigated the effects of acute exercise on monocyte subpopulations in patients with metabolic syndrome (31). Results seem to demonstrate that strenuous exercise, even if it is not sufficient for weight loss, might exert a positive effect on vascular plaque formation by removing pro-inflammatory monocytes from the endothelium. However, as outlined by authors, this hypothesis needs to be further investigated, since strenuous exercise was able to mobilize the same amount of proinflammatory monocytes in MetS patients as in healthy persons; it is conceivable that the elevated basal level of these cells in MetS patients is likely to be caused by enhanced maturation rather than chronic mobilization. The removal of these monocytes from the endothelium might be part of the beneficial effect of exercise on vascular disease in long term studies (32,33).

Conventional data analysis of flow cytometry-based basophil activation testing requires repetitive, labor-intensive analysis that hampers efforts to standardize testing for clinical applications. Using an open-source platform, Patil et al. developed and implemented a

programmatic approach to the analysis of the basophil activation test (BAT) by flow cytometry. This novel method provided a high throughput objective approach to basophil activation analysis (34).

Del Zotto et al. published an interesting paper dealing with a rare genetic disorder, called Fibrodysplasia Ossificans Progressiva (FOP), caused by sporadic heterozygous mutations in *ACVR*, that is a gene which progressively leads to severe heterotopic ossification (35). The peripheral blood mononuclear cell immunophenotyping by flow cytometry in samples from this genetic disorder provided evidence for monocyte DNAM1 up-regulation, as well as significant differences in the expression profile of CXCR1 (CD181), CD62L, CXCR4 (CD184), and HL-DR molecules. Based on the notion that DNAM1 had been previously shown to play a pivotal role in monocyte migration through the endothelial barrier, the increased expression detected in patients' monocytes might suggest a role of this surface receptor during the early phases of FOP flare-ups in which the activation of the immune response is believed to represent a crucial event.

The original paper by Chutvanichkul dealt with labile iron pool (LIP) as a parameter to monitor iron overload and oxidative stress status in the erythrocyte population from  $\beta$ -thalassemia patients (36). LIP is intracellular non-protein bound iron that can generate oxygen radicals via the Fenton reaction resulting in oxidative cell damage. As a consequence, quantitative assessment of LIP may be helpful for detecting and monitoring the toxic iron status in iron overloaded patients. Based on authors results and speculation, it may be hypothesized that LIP assay may represent an alternative test to monitor the magnitude of iron overload and its consequent oxidative stress in  $\beta$ -thalassemia patients. LIP level may also be used as a marker for a careful evaluation of the therapeutic response to iron chelation treatment.

The investigation from the group of North Texas University had the main scope to study the effects of a single high-fat meal on monocyte adhesion molecule expression, CD36 expression and acLDL endocytosis using image-based flow cytometry. Results showed that consumption of a high-fat meal was associated with an increased adhesion molecules expression in both classical and non-classical monocytes, scavenger receptors, and propensity to form foam cells (37).

The current issue of *Cytometry* also includes six interesting brief communications and two letters to the editor, which all deserve a careful evaluation from the readers.

I do believe that the flow cytometry community is today very active, productive, and vital, and in the next years a lot of developments in this area will enrich our knowledge in many research and clinical disciplines, thus enhancing the use of flow cytometry as an ever-increasing powerful diagnostic tool.

Francesco Lanza,  
Hematology Institute  
Romagna Stem Cell Transplantation Programme,  
Ravenna 48121, Italy

## LITERATURE CITED

1. Depreter B, Philippe J, Meul M, Denys B, Vandepoele K, De Moerloose B, Lammens T. Cancer-related mRNA expression analysis using a novel flow cytometry-based assay. *Cytometry B Clin Cytom* (in press).
2. Ikoma MRV, Sandes AF, Thiago LS, Cavalcanti Junior GB, Lorand-Metze IG, Costa ES, Pimenta G, Santos-Silva MC, Bacal NS, Yamamoto M, et al. First proposed panels on acute leukemia for four-color immunophenotyping by flow cytometry from the Brazilian group of flow Cytometry-GBCFLUX. *Cytometry Part B Clin Cytom* 2015;88B:194-203.
3. Meyerson HJ, Osei E, Schweitzer K, Blidaru G, Edinger A, Schlegelmilch J, Awadallah A, Goyal T. CD1c(1) myeloid dendritic cells in myeloid neoplasia. *Cytometry Part B Clin Cytom* 2016;90B:337-348.
4. Mani R, Goswami S, Gopalakrishnan B, Ramaswamy R, Wasmuth R, Tranh M, Mo X, Gordon A, Bucci D, Lucas DM, et al. The interleukin-3 receptor CD123 targeted SL-401 mediates potent cytotoxic activity against CD34+/CD123+ cells from acute myeloid leukemia/myelodysplastic syndrome patients and healthy donors. *Haematologica* (in press). doi:10.3324/haematol.2018.188193.
5. Kornblau SM, Ruvolo PP, Wang RY, Battula VL, Shpall EJ, Ruvolo VR, McQueen T, Qui Y, Zeng Z, Pierce S, et al. Distinct protein signatures of acute myeloid leukemia bone marrow-derived stromal cells are prognostic for patient survival. *Haematologica* 2018;103:810-821. doi:10.3324/haematol.2017.172429.
6. Keup C, Mach P, Aktas B, Tewes M, Kolberg HC, Hauch S, Sprenger-Haussels M, Kimmig R, Kasimir-Bauer S. RNA profiles of circulating tumor cells and extracellular vesicles for therapy stratification of metastatic breast cancer patients. *Clin Chem* (in press). doi:10.1373/clinchem.2017.283531.
7. Ravelli A, Reuben JM, Lanza F, Anfossi S, Spada D, Pedrazzoli P, Martino M, Bottini A, Generali D. Breast cancer circulating biomarkers: Advantages, drawbacks, and new insights. *Tumor Biol* 2015;36:6653-6665. doi:10.1007/s13277-015-3944-7.
8. Lanza F, Campioni D, Hellmann A, Milone G, Wahlin G, Walewski J, Spedini P, Fiamenghi C, Cuneo A, Knopińska W, et al. Individual quality assessment of autografting by probability estimation for clinical endpoints: A prospective validation study from the European group for blood and marrow transplantation. *Biol Blood Marrow Transplant* 2013;19:1670-1676. doi:10.1016/j.bbmt.2013.08.005
9. Starostka D, Kriegova E, Kudelka M, Mikula P, Zehnalova S, Radvansky M, Papajik T, Kolacek D, Chasakova K, Talianova H. Quantitative assessment of informative immunophenotypic markers increases the diagnostic value of immunophenotyping in mature CD5-positive B-cell neoplasms. *Cytometry B Clin Cytom* (in press).
10. Preijers FWMB, van der Velden VHJ, Preijers T, Brooimans RA, Marijt E, Homburg C, van Montfort K, Gratama JW, Gratama JW. Fifteen years of external quality assessment in leukemia/lymphoma immunophenotyping in The Netherlands and Belgium: A way forward. *Cytometry Part B Clin Cytom* 2016;90B:267-278.
11. Wolniak K, Goolsby C, Choi S, Ali A, Serdy N, Stetler-Stevenson M. Report of the results of the International Clinical Cytometry Society and American Society for Clinical Pathology workload survey of clinical flow cytometry laboratories. *Cytometry Part B Clin Cytom* 2017;92B:525-533.
12. Bagwell CB, Hill BL, Wood BL, Wallace PK, Alrazzak M, Kelliher AS, Preffer F. Human B-cell and progenitor stages as determined by probability state modeling of multidimensional cytometry data. *Cytometry Part B Clin Cytom* 2015;88B:214-226.
13. Auat M, Cardoso CC, Santos-Pirath IM, Rudolf-Oliveira RCM, Mاتيollo C, Lange BG, Pires da Silva J, Dامتetto GC, Pirolli MM, Colombo MDHP, Santos-Silva MC. Evaluation of CD307a expression patterns during normal B-cell maturation and in B-cell malignancies by flow cytometry. *Cytometry B Clin Cytom* (in press).
14. Flores-Montero J, de Tute R, Paiva B, Perez JJ, Bottcher S, Wind H, Sanoja L, Puig N, Lecrevisse Q, Vidriales MB, et al. Immunophenotype of normal vs. myeloma plasma cells: Toward antibody panel specifications for MRD detection in multiple myeloma. *Cytometry Part B Clin Cytom* 2016;90B:61-72.
15. Wood BL. Principles of minimal residual disease detection for hematopoietic neoplasms by flow cytometry. *Cytometry Part B Clin Cytom* 2016;90B:47-53.
16. Tembhare PR, Ghogale S, Ghatwai N, Badrinath Y, Kunder N, Patkar NV, Bibi AR, Chatterjee G, Arora B, Narula G, et al. Evaluation of new markers for minimal residual disease monitoring in b-cell precursor acute lymphoblastic leukemia: CD73 and CD86 are the most

- relevant new markers to increase the efficacy of MRD assay. *Cytometry Part B Clin Cytom* 2018;94B:100-111.
17. Luzzatto L, Paroxysmal nocturnal hemoglobinuria: an acquired X-linked genetic disease with somatic-cell mosaicism. *Curr Opin Genet Dev* 2006;16:317-322.
  18. Mannelli F, Bencini S, Peruzzi B, Cutini I, Sanna A, Benelli M, Magi A, Gianfaldoni G, Rotunno G, Carrai V, et al. A systematic analysis of bone marrow cells by flow cytometry defines a specific phenotypic profile beyond GPI deficiency in paroxysmal nocturnal hemoglobinuria. *Cytometry B Clin Cytom* 2013;84:71-81.
  19. Lanza F, Lazzari MC, Brambilla P, Di Martino G, Spedini G. An unusual association of paroxysmal nocturnal hemoglobinuria, myelodysplastic syndrome, and diffuse large B-cell non-Hodgkin lymphoma in a Caucasian man. *Ann Hematol* 2016;35:1555-1557. doi:10.1007/s00277-016-2728-5
  20. Liew M, Farley M, Andreasen J, Parker CJ, Wittwer CT. Rare event counting of cd592 red cells in human blood: A 47-month experience using PNH consensus guidelines for WBC and RBC testing in a reference lab. *Cytometry Part B Clin Cytom* 2015;88B:261-269.
  21. Shih AR, Murali MR. Laboratory tests for disorders of complement and complement regulatory proteins. *Am J Hematol* 2015;90:1180-1186. doi:10.1002/ajh.24209.
  22. Preis M, Lowrey CH. Laboratory tests for paroxysmal nocturnal hemoglobinuria. *Am J Hematol* 2014;89:339-341. doi:10.1002/ajh.23612.
  23. Blaha J, Schwarz K, Fischer C, Schauwecker P, Höchsmann B, Schrezenmeier H, Anliker M. The monoclonal anti-CD157 antibody clone SY11B5, used for high sensitivity detection of PNH clones on WBCs, fails to detect a common polymorphic variant encoded by BST. *Cytometry B Clin Cytom* (in press).
  24. Sutherland DR, Ortiz F, Quest G, Illingworth A, Benko M, Nayyar R, Marinov I. High sensitivity 5-, 6-, and 7-color PNH WBC assays for both CANTO II and NAVIOS platforms. *Cytometry B Clin Cytom* (in press).
  25. Marinov I, Illingworth AJ, Benko M, Sutherland DR. Performance characteristics of a non-fluorescent aerolysin-based paroxysmal nocturnal hemoglobinuria (PNH) assay for simultaneous evaluation of PNH neutrophils and PNH monocytes by flow cytometry, following published PNH guidelines. *Cytometry B Clin Cytom* 2018; 94:257-263. doi:10.1002/cyto.b.21389.
  26. Illingworth A, Marinov I, Sutherland DR, Wagner-Ballon O, Del Vecchio L. ICCS/ESCCA consensus guidelines to detect GPI-deficient cells in paroxysmal nocturnal hemoglobinuria (PNH) and related disorders part 3 - data analysis, reporting and case studies. *Cytometry B Clin Cytom* 2018;94:49-66. doi:10.1002/cyto.b.21609.
  27. Lindemann M, Lenz V, Nyadu B, Heinemann FM, Heinold A, Guberina H, Eisenberger U, Lachmann N, Schönemann C, Kribben A, et al. Effect of ABO incompatibility on T-cell flow cytometry cross-match results prior to living donor kidney transplantation. *Cytometry B Clin Cytom* (in press). doi:10.1002/cyto.b.21496.
  28. Rajab A, Porwit A. Screening bone marrow samples for abnormal lymphoid populations and myelodysplasia-related features with one 10-color 14-antibody screening tube. *Cytometry Part B* 2015;88B:253-260.
  29. Langenskiöld C, Mellgren K, Abrahamsson J, Bemark M. Determination of blood cell subtype concentrations from frozen whole blood samples using TruCount beads. *Cytometry B Clin Cytom* (in press).
  30. Rasmussen SM, Bilgrau AE, Schmitz A, Falgreen S, Bergkvist KS, Tramm AM, Bæch J, Jacobsen CL, Gaihede M, Kjeldsen MK, et al. Stable phenotype of B-cell subsets following cryopreservation and thawing of normal human lymphocytes stored in a tissue biobank. *Cytometry Part B* 2015;88B:40-49.
  31. Wonner R, Wallner S, Orso E, Schmitz G. Effects of acute exercise on monocyte subpopulations in metabolic syndrome patients. *Cytometry B Clin Cytom* (in press).
  32. Hudig D, Hunter KW, Diamond WJ, Redelman D. Properties of human blood monocytes. II. Monocytes from healthy adults are highly heterogeneous within and among individuals. *Cytometry Part B Clin Cytom* 2014;86B:121-134.
  33. Damuzzo V, Pinton L, Desantis G, Solito S, Marigo I, Bronte V, Mandruzzato S. Complexity and challenges in defining myeloid-derived suppressor cells. *Cytometry Part B Clin Cytom* 2015;88B:77-91.
  34. Patil SA, Schneider M, Steinbrecher J, Smith N, Washburn C, Ma A, Shreffler WG. Data-driven programmatic approach to analysis of basophil activation tests. *Cytometry B Clin Cytom* (in press).
  35. Del Zotto G., Antonini F, Azzari I, Ortolani C, Tripodi Gino; Giacomelli F, Cappato S, Moretta L. Peripheral blood mononuclear cell immunophenotyping in fibrodysplasia ossificans progressive patients: evidence for monocyte DNAM1 up regulation. *Cytometry B Clin Cytom* (in press).
  36. Chutvanichkul B, Vattanaviboon P, Sumana Mas-oodi S, U-pratya Y, Wanachiwanawin W. Labile iron pool as a parameter to monitor iron overload and oxidative stress status in  $\beta$ -thalassemic erythrocytes. *Cytometry B Clin Cytom* (in press).
  37. Henning AL, Venable AS, Vingren JL, Hill DW, and McFarlin BK. Consumption of a high-fat meal was associated with an increase in monocyte adhesion molecules, scavenger receptors, and propensity to form foam cells. *Cytometry B Clin Cytom* (in press).