

Litter decomposition in peatlands is promoted by mixed plants

Fabien Leroy, Sébastien Gogo, Alexandre Buttler, Luca Bragazza, Fatima Laggoun-Défarge

► **To cite this version:**

Fabien Leroy, Sébastien Gogo, Alexandre Buttler, Luca Bragazza, Fatima Laggoun-Défarge. Litter decomposition in peatlands is promoted by mixed plants. *Journal of Soils and Sediments*, Springer Verlag, 2018, 18, pp.739-749. 10.1007/s11368-017-1820-3 . insu-01598365

HAL Id: insu-01598365

<https://hal-insu.archives-ouvertes.fr/insu-01598365>

Submitted on 2 Oct 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



1 SOILS, SEC # • RESEARCH ARTICLE

2

3 **Litter decomposition in peatlands is promoted by mixed plants**

4

5 **Fabien Leroy^{1,2,3} • Sébastien Gogo^{1,2,3} • Alexandre Buttler^{4,5,6} • Luca Bragazza^{4,5,7} • Fatima Laggoun-**
6 **Défarge^{1,2,3}**

7

8 ¹Université d'Orléans, ISTO, UMR 7327, 45071, Orléans, France

9 ²CNRS/INSU, ISTO, UMR 7327, 45071 Orléans, France

10 ³BRGM, ISTO, UMR 7327, BP 36009, 45060 Orléans, France

11 ⁴WSL Swiss Federal Institute for Forest, Snow and Landscape Research, Site Lausanne, Station 2, CH-1015
12 Lausanne, Switzerland

13 ⁵Laboratory of Ecological Systems (ECOS), Ecole Polytechnique Fédérale de Lausanne (EPFL), School of
14 Architecture, Civil and Environmental Engineering (ENAC), Station 2, CH-1015 Lausanne, Switzerland

15 ⁶Laboratoire de Chrono-Environnement, UMR CNRS 6249, UFR des Sciences et Techniques, 16 route de Gray,
16 Université de Franche-Comté, F-25030 Besançon, France

17 ⁷Department of Life Science and Biotechnologies, University of Ferrara, Corso Ercole I d' Este 32, I-44121
18 Ferrara, Italy

19

20

21 ✉ Fabien Leroy

22 tel: + 33 2 38 49 45 02

23 fabien.leroy@univ-orleans.fr

24

25 **Abstract**

26 *Purpose.* The carbon sink function of peatlands is primarily driven by a higher production than decomposition of
27 the litter *Sphagnum* mosses. The observed increase of vascular plants in peatlands could alter the decomposition
28 rate and the carbon (C) cycle through a litter mixing effect, which is still poorly studied. Here, we examine the
29 litter mixing effect of a peat moss (*Sphagnum fallax*) and two vascular plants (*Pinus uncinata* and *Eriophorum*
30 *vaginatum*) in the field and laboratory-based experiment.

31 *Materials and methods.* During the laboratory incubation, mass loss, CO₂ production and dissolved organic carbon
32 concentration were periodically monitored during 51 days. The collected data were then processed in a C dynamics
33 model. The calculated enzymatic activity was correlated to the measured β -glucosidase activity in the litter. In the
34 field experiment, mass loss and CO₂ production from litter bags were annually measured for three years.

35 *Results and discussion.* Both laboratory and field experiments clearly show that the litter mixture, i.e. *Sphagnum-*
36 *Pinus-Eriophorum*, had a synergistic effect on decomposition by enhancing the mass loss. Such enhanced mass
37 loss increased the water extractable C and CO₂ production in the litter mixture during the laboratory experiment.
38 The synergistic effect was mainly controlled by the *Sphagnum-Eriophorum* mixture that significantly enhanced
39 both mass loss and CO₂ production. Although the β -glucosidase activity is often considered as a major driver of
40 decomposition, mixing the litters did not cause any increase of the activity of this exo-enzyme in the laboratory
41 experiment suggesting that other enzymes can play an important role in the observed effect.

42 *Conclusions.* Mixing litters of graminoid and *Sphagnum* species led to a synergistic effect on litter decomposition.
43 In a context of vegetation dynamics in response to environmental change, such a mixing effect could alter the C
44 dynamics at a larger scale. Identifying the key mechanisms responsible for the synergistic effect on litter
45 decomposition, with a specific focus on the enzymatic activities, is crucial to better predict the capacity of
46 peatlands to act as C sinks.

47

48 **Keywords** β -glucosidase • Carbon dynamics models • Catalysis • CO₂ production • Dissolved organic carbon •
49 Litter mixture effect

50

51 **1. Introduction**

52 *Sphagnum*-dominated peatlands accumulate organic matter (OM) as peat at a rate of ca. 20 to 30 g C m⁻² year⁻¹
53 (Francez 2000; Rydin et al. 2013). At global scale, peatlands are estimated to store about 270 to 547 Pg C as peat
54 (1 Pg = 10¹⁵ g), representing ca. 15-30% of the world's soil carbon (C) stock in an area accounting for only 3-5%
55 of the land surface (Turunen et al. 2002; Yu et al. 2010). This high rate of C accumulation is due to peculiar
56 environmental and soil conditions, i.e. waterlogging, anoxia, acidity, low temperature, and specific plant species
57 composition that ultimately hamper the microbial litter decomposition so resulting in a net accumulation of OM
58 as peat (e.g. Gorham, 1991; Holden, 2005). In particular, *Sphagnum* mosses have a key role for peat accumulation
59 as they are able to create unfavorable conditions for decomposer activities by producing a recalcitrant litter and by
60 promoting waterlogged and acidic conditions (Van Breemen 1995). As a result, *Sphagnum* mosses gain in
61 competitive ability against vascular plants, whose litter is much more easily decomposable (Bragazza et al. 2007,
62 2009), thereby acting as effective ecosystem engineers (Van Breemen 1995).

63 Human-induced environmental changes are expected to modify plant species abundance in peatlands, in
64 particular by favoring vascular plants at the expense of *Sphagnum* mosses under a warmer climate (Buttler et al.,
65 2015; Dieleman et al., 2015). Although previous studies have shown that warming can modify the rate of litter
66 decomposition in peatlands (Thormann et al., 2004; Bragazza et al. 2016) and that decomposition rates of
67 *Sphagnum* are lower than those of vascular plants (Hoorens et al., 2010), few studies have addressed the issue of
68 a litter mixture effect in peatlands (Hoorens et al. 2010; Krab et al. 2013; Gogo et al. 2016). Most of the studies
69 on litter decomposition in peatlands have focused on single plant species, although in natural conditions litters
70 mainly consist of a mixture of multiple plant species (Salamanca et al. 1998). Litters in mixture can decompose
71 faster (synergistic effect) or slower (antagonistic effect) than the same litter type alone (non-additive effect
72 (Gartner and Cardon, 2004). Almost 70% of mixed-species litter exhibited non-additive mass loss with a
73 prevalence of synergistic effects (Gartner and Cardon, 2004). Such modifications of the decomposition rate due to
74 litter mixture could affect the imbalance between primary production and decomposition in peatlands in response
75 to a vegetation dynamics and, ultimately, the capacity of peatlands to act as C sinks. Despite the crucial role of
76 litter decomposition in controlling C dynamics and peat accumulation in peatlands, we have still little
77 understanding of how the mixture litter affects decomposition.

78 As change in leaf litter quality can affect enzyme production by soil microbes (Hu et al. 2006), litter mixture
79 could also change enzyme activities, which may result in a non-additive effect. For example, β -glucosidase is
80 commonly described as an important exo-enzyme involved in C-cycling, and therefore primarily in the

81 decomposition of cellulose (Kourtev et al. 2002; Sinsabaugh et al. 2002), so that it could be a key-player in the
82 non-additive effect of litter mixture. Nevertheless, the role of enzymes in relation to litter decomposition still needs
83 to be elucidated.

84 In order to understand the effect of a litter mixture of *Sphagnum* mosses and vascular plants on OM
85 decomposition and C dynamics in peatlands, we performed a decomposition experiment under laboratory and field
86 conditions. Experimental laboratory data were then used to calibrate a C dynamics model proposed by Gogo et al.
87 (2014), using three C compartments: the solid (mass loss), the dissolved (Water Extractable Organic Carbon,
88 WEOC) and the gaseous components (CO₂-C). In this model, the litter is catalyzed by exo-enzymes at a rate “c”
89 to produce soluble organic C, i.e. WEOC, which is used for respiration at a rate “r” by microorganisms. The model
90 gives a catalysis rate for each litter that contributes to overall catalytic activity. The field experiment was performed
91 to assess whether there was agreement between laboratory and field decomposition tests by focusing on mass loss
92 and CO₂ production. Overall, the aims of this work were: (i) to determine the occurrence of a litter mixture effect
93 for three different plant species during decomposition; (ii) to elucidate the role of β-glucosidase activity during
94 litter decomposition; (iii) to relate the catalysis rate from the C dynamics modeling to the activity of hydrolytic
95 enzymes.

96

97 **2. Materials and methods**

98 **2.1 Study site and litter sampling**

99 Plant litter material for laboratory and field experiments was collected at the Forbonnet peatland in Frasné (France,
100 N46°49'35" E 6°10'20", 840m). The site is a *Sphagnum*-dominated peatland with a mean annual temperature of
101 7.5°C and annual rainfall amount of 1400 mm (Laggoun-Défarge et al. 2008; Delarue et al. 2011). The vascular
102 plant cover mainly consists of *Eriophorum vaginatum*, *Scheuchzeria palustris*, *Andromeda polifolia*, *Vaccinium*
103 *oxycoccos* and *Carex limosa* (Buttler et al. 2015) with a recent increased abundance of trees of the genus *Pinus* in
104 response to a decline in peat water content, as observed in many bogs in the study region (Frelechoux et al. 2000).
105 Overall, the moss layer is primarily dominated by *Sphagnum fallax* and *S. magellanicum*. Litter samples of
106 *Eriophorum vaginatum* (E) and *Pinus uncinata* (P) were composed, respectively, of senescent leaves and needles.
107 Litter samples of *Sphagnum fallax* (S) corresponded to the decaying part just below the green photosynthesizing
108 apical part (Bragazza et al. 2007). After having been air-dried, sub-samples of each litter type were oven dried at
109 50°C for 48 hours in order to calculate the dry weight.

110

111 2.2 Laboratory experiment

112 **Sample preparation and incubation** - For the laboratory experiment, samples of one gram for seven litter types
113 were prepared. Litter types were formed by single litter of *Sphagnum fallax* (S), *Eriophorum vaginatum* (E) or
114 *Pinus uncinata* (P), and by the following mixtures: *Sphagnum fallax* + *Eriophorum vaginatum* (SE), *Sphagnum*
115 *fallax* + *Pinus uncinata* (SP), *Pinus uncinata* + *Eriophorum vaginatum* (PE) and *Sphagnum fallax* + *Pinus uncinata*
116 + *Eriophorum vaginatum* (SPE). Litter samples received 20 mL of peatland water overnight in order to inoculate
117 microorganisms (Hoorens et al. 2002). The excess water was removed with a tissue and the litter samples were
118 placed in aluminum cups with small holes at the bottom to allow air circulation. The cups were placed on a 0.4 L
119 pot containing 20 mL of saturated solution of K₂SO₄ to maintain moist conditions. Each pot with its cup was
120 covered with perforated aluminum foil to allow air circulation. All pots were placed in an incubator (Aralab 1200)
121 at 20°C and 95% of relative humidity. A total of 105 litter samples (pots) were prepared in order to have 3 replicates
122 for each of the 7 litter types and 5 periodical harvests. At intervals of 0, 1, 14, 28 and 51 days after incubation, we
123 measured CO₂ production, water extractable organic carbon (WEOC) release, litter mass loss, β-glucosidase
124 activities and C content in 3 replicats of each litter type.

125 **Laboratory measurements** - The CO₂ production was measured with a GMP343 Vaisala probe after placing the
126 litter sample (pots) in a 2.34 L chamber. The measurement took 15 minutes. The slope of CO₂ increased over time
127 within the chamber (in μmol CO₂. mol air⁻¹.sec⁻¹) and was used to calculate a cumulative C release (g C g⁻¹ initial
128 C litter). The WEOC from each sample was obtained after rinsing the incubated litter three times with 50 mL of
129 distilled water and followed the method described in Delarue et al. (2011). The extract was filtered through a 0.45
130 μm membrane filter. Dissolved organic carbon (DOC) in water extract was determined with a Shimadzu TOC
131 5000A (Total Organic Carbon Analyzer). The particles on the 0.45 μm filter and the remaining litter were dried at
132 50°C during two days and then weighed to obtain the remaining mass (mass loss = initial mass – remaining mass)
133 in g OM g⁻¹ initial litter mass. The C in each litter sample was then measured with an elementary analyzer
134 (Thermo-FLASH 2000 CHNS/O Analyzer). This normalized the three C pools in g C g⁻¹ initial C litter, thus
135 making possible to input data in the C dynamics models. The activity of β-glucosidase was measured by adding to
136 0.5 mL of water extracted litter, 3.5 mL of 4-Methylumbelliferyl β-D-glucopyranoside, a substrate for β-
137 glucosidase activities which is transformed into 4-Methylumbelliferone (MUF). After one hour of incubation, the
138 concentration of MUF was determined by fluorescence using an excitation wavelength of 330 nm and an emission
139 wavelength of 450 nm and was compared to standard solution. Measured values obtained in the litter mixtures

140 were compared to calculate additive values (expected) after Hoorens et al. (2010), which are means of the values
141 as measured in the decomposition of the single species litter type.

142 **C dynamic modeling** - The experimental design was similar to that published by Gogo et al. (2014, 2016). These
143 authors conceptualized the model assuming that solid OM is catalyzed by exo-enzymes, leading to soluble OM,
144 which is then absorbed by microorganisms and used as an energy source for different microbial functions (enzyme
145 production, maintenance, growth). Then, the soluble organic C is respired and released into the environment in the
146 form of CO₂. The applied model followed Schimel and Weintraub (2003) but was simplified to make it
147 experimentally testable (Supplementary Fig. S6). It is composed of three compartments: (i) the “L” compartment
148 corresponding to the C fraction contained in the litter (solid fraction), (ii) the “W” compartment corresponding to
149 the C fraction in the WEOC (dissolved fraction) and (iii) the “G” compartment corresponding to the C fraction in
150 the cumulative CO₂ released by microbial respiration (gaseous fraction). C flow from the L to the W compartment
151 at a rate corresponding to the exo-enzyme catalysis rate. C flows from the W to the G compartment at a rate
152 corresponding to the respiration rate. Equations describe the simultaneous change in time of the state variable (L,
153 W, and G) and the reaction rates (Gogo et al. 2014; Supplementary Fig. S6). At any time, the sum of all these three
154 fractions is equal to 1. The three fractions corresponding to C stocks (solid, soluble, gaseous) were experimentally
155 measured. The catalysis rate “c” and the respiration rate “r” were tuned at the same time to fit the stock values of
156 the model to those experimentally assessed. When the reaction rates are allowed to change in the course of the
157 experiment, the goodness of fit is improved. The reaction rates were allowed to follow a negative exponential
158 decrease with time. The following parameters describe the shape of the curve: “a+b” = the initial reaction rate, “a”
159 = the final rate, “m” = decay rate with time of the reaction rate (Rovira and Rovira, 2010). Observed rates obtained
160 in litter mixture were compared to calculate additive rates, which are means of the catalysis and respiration rates
161 measured in decomposition of the single species litter type. The root mean square error (RMSE) was calculated
162 for each litter type in order to represent model performance. The RMSE is calculated by squaring individual errors,
163 summing them, dividing the sum by their total number (N), and then taking the square root of this quantity:

$$RMSE = \sqrt{\sum \frac{(y_{pred} - y_{meas})^2}{N}}$$

164
165 The RMSE was normalized to the mean of observed data and multiplied by 100 to obtain the Normalized Root

166 Mean Square Error (NRMSE) in percentage (%):
$$NRMSE = 100 \times \frac{RMSE}{\bar{y}}$$

167

168 2.3 Field experiment

169 The litterbag experiment was performed at the Forbonnet peatland in order to annually measure mass loss and CO₂
170 production during three years of field incubation. Litterbags (0.5 mm mesh) were prepared with one gram of litter
171 of S, P, E and in a mixture of the three species (SPE). For the SPE litter mix, each species contributed equally to
172 the final weight of the litter bag. Litter bags were placed vertically in the moss carpet and buried at 5 cm depth on
173 November 2009 in lawns, which represent the major feature in this bog, avoiding hummocks where there was tree
174 encroachment. Twelve samples were prepared for each of the four litter treatments that were periodically collected
175 after one, two and three years of incubation. Litter mass loss was calculated as percentage of remaining mass of
176 initial litter mass. The CO₂ flux was measured immediately after collection in a 835 cm³ chamber with a Li-Cor
177 LI-8100 analyser and expressed as fluxes (in µg CO₂ h⁻¹ g⁻¹ dry weight).

178

179 2.4 Statistics

180 The role of litter mixture in affecting the measured variables was compared to the corresponding additive
181 calculated values (expected) from the litter decomposition of each single species. Two-way ANOVA was applied
182 to investigate differences in the measured variables for both the field and laboratory experiments where the selected
183 factors were “litter types” (observed vs additive) and “time (years or days) of measurements. Post-hoc Tukey’s
184 Honestly Significant Differences (HSD) tests were performed to determine significant differences within groups.
185 Differences were considered as a trend for $p < 0.1$ (noted with: -) and significant at $p < 0.05$ and referred to by * =
186 $p < 0.05$, ** = $p < 0.01$, *** = $p < 0.001$. Modeled curves for temporal changes of variables in the laboratory
187 incubation experiment were calculated from the C dynamics model. All statistical analyses were done using R
188 3.1.1 software.

189

190 **3. Results**

191 3.1 Laboratory experiment

192 During the laboratory incubation of the single litter type, *Sphagnum fallax* (S) litter was characterized by a mass
193 loss, DOC release and CO₂ production significantly different compared to *Pinus uncinata* (P) and *Eriophorum*
194 *vaginatum* (E) litter (Fig. 1). The S, E and P litters contained, respectively, 42.4, 50.9 and 47.7 % of C (Table 1).
195 These contents were constant in time and between additive (calculated values) and measured values in mixture
196 litter (Table 1). When compared to the decomposition calculated from the single plant litter (additive mean values),
197 the observed *Sphagnum + Pinus + Eriophorum* (SPE) mixture showed a significantly higher mass loss (i.e. lower

198 remaining C) ($p < 0.001$), a higher WEOC ($p < 0.05$) as well as a higher production of CO_2 ($p: 0.063$) (Fig. 2). The
199 litter mixture of *Sphagnum* + *Eriophorum* (SE) significantly ($p < 0.001$) increased mass loss and CO_2 production
200 by 13% after 51 days of incubation, as compared to the additive effect (Fig. 2, Table 2). The C dynamics in the SP
201 and PE mixtures did not significantly differ from the additive effect of the single species (Fig. 2, Table 2) although
202 the litter mixture of *Pinus* and *Eriophorum* (PE) showed a tendency to decrease litter mass loss and CO_2 production
203 ($p: 0.069$ and 0.097 , respectively).

204

205 3.2 Field experiment

206 After three years of field incubation, the decomposition of single litter of *S. fallax* was slower than that of vascular
207 plants ($p < 0.001$) (Fig. 3, Table 3). *S. fallax* litter showed significant differences for CO_2 production compared to
208 *Pinus uncinata* ($p < 0.001$) and *Eriophorum* litter ($p < 0.05$). There was a significant effect of litter mixture during
209 field decomposition, with a mass loss higher (ca. 19%) than that expected from the single species additive effect
210 ($p < 0.01$) (Fig. 4), while the CO_2 production in the mixture litter did not differ from the additive effect (Fig. 4,
211 Table 3).

212

213 3.3 Catalysis of C dynamics models and β -glucosidase activities

214 The model fitted well the measured values with a lower NRMSE (sum between 2.82 and 9.41 %; Supplementary
215 Table S5) compared to Gogo et al. (2014). The catalysis rate decreased with incubation time in each litter type
216 (Table 4). In *Sphagnum* litter, this rate was initially very fast and decreased rapidly to become slower than vascular
217 plant litters. The catalysis rates measured in *Sphagnum-Eriophorum* (SE) and *Sphagnum-Pinus-Eriophorum* (SPE)
218 mixtures were always greater than expected from the mean of single litter species at each time of measurement
219 (Table 4). This corresponded to higher C mass loss, WEOC and CO_2 production.

220 At the beginning of incubation, β -glucosidase activity was similar in all litter types but afterward it significantly
221 increased in *Eriophorum* and *Pinus* litters, while remaining constant in *Sphagnum* litter (Supplementary Fig. S7).
222 Mixing litters did not affect β -glucosidase activities so that there were no significant differences between observed
223 and calculated additive values (Fig. 5). Overall, β -glucosidase activities increased with incubation time, whereas
224 the catalysis rate decreased (Fig. 5; Table 4) so that the catalysis rate was inversely correlated to the β -glucosidase
225 activity ($r^2: 0.52$; $p < 0.001$; Supplementary Fig. S8).

226

227 4. Discussion

228 4.1 Occurrence of a synergistic effect

229 The laboratory incubation experiment clearly showed that the litter mixture of *Sphagnum-Eriophorum* as well
230 as *Sphagnum-Pinus-Eriophorum* had a synergistic effect on mass loss and CO₂ production compared to the
231 corresponding additive effect. Similar results were obtained with a mixture of *Sphagnum* litter and graminoid
232 species (Hoorens et al. 2002; Gogo et al. 2016). Such synergistic interaction on decomposition has been explained
233 by differences in litter chemistry such as N concentration (Hoorens et al. 2002). However, similar N concentration
234 was found in our litter types. Mixing different litters can produce both chemical diversity and microhabitat
235 complexity so supplying an increased diversity of substrates to the decomposers (Gartner and Cardon, 2004). Also,
236 special attention in future studies should be devoted to the improvement of microclimatic condition such as the
237 water content of individual litter in mixture through water flow from the wettest to the driest litter. The hypothesis
238 is that through such water flow the conditions are improved in the driest litter, without decreasing to large extent
239 the conditions of the wettest litter (Gogo et al., 2017).

240 The field experiment also showed an increase of decomposition by mixing *Sphagnum + Eriophorum + Pinus*
241 litter. *Sphagnum fallax* decomposed at a slower rate than *Pinus uncinata* and *Eriophorum vaginatum* litter as
242 observed elsewhere by Hobbie (1996). Curiously, we expected to have higher CO₂ production in combination with
243 higher litter mass loss, but, instead, the highest CO₂ production was measured in *Sphagnum* litter with the slowest
244 mass loss. The high capacity of *Sphagnum* litter to maintain capillary water could have contributed to retaining
245 water that had previously percolated through the upper photosynthetically active centimeters of the *Sphagnum*
246 carpet. This percolating water may have been enriched in labile C that could stimulate CO₂ production.
247 Furthermore, the capillary structure of *Sphagnum* is known to host rich microbial communities, including
248 microbial predators, i.e. amoebae (Jassey et al., 2013) in the living apical part and we cannot exclude the possibility
249 that this might have affected the decaying part in the long field run. This could explain the decoupling observed
250 between mass loss and respiration rate.

251 Laboratory incubation was a short experiment in which environmental factors and C input and output were
252 well controlled. Conversely, field experiments spanned over three years and thus represent litter mass loss as it
253 occurs in natural field conditions. In both laboratory and field experiments, mixing *Sphagnum* with *Eriophorum*
254 and *Pinus* litter had a synergistic effect on litter decomposition. Such non-additive effects have already been
255 reported for other ecosystems (Wu et al. 2013; Zhang et al. 2014), although the reasons are still unclear. Many
256 studies have tried to explain such an effect by nutrient exchanges between litters (Vos et al. 2013), litter chemical
257 quality (Meier and Bowman 2010) or changes in habitat characteristics (Lecerf et al. 2011). Nevertheless, only

258 very few studies have addressed this effect at microbial scale and established links between microbial communities
259 and litter decomposition (Chapman et al. 2013).

260

261 4.2 Role of β -glucosidase in early C dynamics the early decomposition stages???

262

263 Contrary to our hypothesis, no significant increases were noticed in β -glucosidase activity between the
264 observed and the additive values during the laboratory experiment, suggesting that, at least in the early stages of
265 decomposition (i.e. the first 28^h days), the synergistic mixture effect does not originate from the stimulation of this
266 hydrolytic enzyme.

267 By comparing the measured enzymatic activities to the modeled catalysis rates, the *Sphagnum* litter showed
268 values similar to those obtained by Gogo et al. (2014) and Gogo et al. (2016), with a fast rate in the early stage of
269 decomposition, and slower but constant values thereafter ($< 0.002 \text{ gC.g}^{-1}\text{C.d}^{-1}$). Contrary to our hypothesis, a
270 negative link was established between modeled catalysis rates and β -glucosidase activities. Such a link between
271 litter enzyme activities and decomposition rates was already assessed with hydrolases and the results showed that
272 this relationship was weak (Allison and Vitousek, 2004). These authors suggested that mass loss occurred
273 independently of enzyme activities with compounds that may leached out of the litter. As this last one, the really
274 different litter types could not allowed to connect the enzymes activities with the decomposition rates.
275 Furthermore, fast catalysis rate in the initial decomposition stage was not related to the β -glucosidase activity but
276 maybe more to the phenol oxidase activity particularly in the case of *Sphagnum* litter (Sinsabaugh et al. 2002).
277 Indeed, it has been shown that phenol oxidase enzymes by degrading inhibitory phenolic compounds, allow other
278 enzymes to act on soil organic C and as a consequence regulate the stability of a vast C store in the soil following
279 the hypothesis of 'enzyme latch' mechanism (Freeman et al., 2001).

280 Linking microbial extracellular enzyme production and litter decomposition modeled catalysis rates could
281 provide information on the mechanisms of decomposition and non-additive effect (Sinsabaugh et al. 2002).
282 However, using different litter types (which can change litter quality, nutrient availability and pH) such a
283 correlation could not be demonstrated. Enzymatic degradation is still necessary to degrade complex and insoluble
284 litter compound and relating modeled catalysis rates and enzymatic activities are likely to provide powerful models
285 that remains poorly understood.

286

287 4.3 Sensitivity of the WEOC compartment

288 Mixing *Sphagnum* with *Pinus* litters did not increase OM decomposition but affected the WEOC
289 compartment, i.e. the dissolved C pool in the laboratory-based experiment. This dissolved fraction described as
290 the pool that is mostly sensitive to any modification in litter decomposition (e.g. Gogo et al. 2014) is considered
291 as a transitory compartment between solid and gaseous pools, which contains a low C-content compared to the
292 other pools. Thus, an increase in litter decomposition would have a strong effect on this compartment as observed
293 for the *Sphagnum-Eriophorum-Pinus* litter mixture. The *Sphagnum-Eriophorum* mixture did not increase the
294 WEOC content as this pool is a ‘dynamic’ compartment that is rapidly transformed into a gaseous C pool. The
295 WEOC increase in *Sphagnum-Pinus* litter could indicate a modification in litter decomposition, which was still
296 not perceived in the solid or gaseous compartments however.

297

298 4.4 Implication in a scenario of vegetation dynamics

299 Like Hoorens et al. (2002) and Gogo et al. (2016), our study showed that mixing graminoid and *Sphagnum*
300 litters together induces a synergistic effect on decomposition. This has important implication for estimating litter
301 decomposition rates in peatlands. These species-specific interactions appeared to be modulated by the changing
302 climate through modifications of litter chemistry (Hoorens et al., 2002). In the context of global change and its
303 role in causing a shift in the relative abundance of *Sphagnum* and vascular plants in peatlands (Dieleman et al.,
304 2015), synergistic effects of mixed litters on decomposition is likely to play a critical role for the C dynamics at
305 ecosystem level.

306

307 **5 Conclusions**

308 On the global scale, the C-sink capacity of *Sphagnum* peatlands could be affected by the litter mixture effect.
309 This calls for further studies on non-additive effects of litter mixture, with a focus on elucidating the specific
310 enzymatic mechanisms behind such interactions, with the ultimate goal of incorporating them in global
311 biogeochemical models.

312

313 **Acknowledgement** This work was funded as part of the PEATWARM initiative through an ANR (French National
314 Agency for Research) grant (ANR-07-VUL-010) and was also supported by the Labex VOLTAIRE (ANR-10-
315 LABX-100-01). The authors gratefully acknowledge the financial support provided to the PIVOTS project by the
316 Région Centre – Val de Loire (ARD 2020 program and CPER 2015 -2020). The authors are also indebted to the
317 Regional Scientific Council of Natural Heritage of the Franche-Comte Region for permitting access to Le
318 Forbonnet site. The authors would like to thank E. Rowley-Jolivet for revision of the English version.

319

320 **References**

- 321 Allison S, Vitousek P (2004) Extracellular enzyme activities and carbon chemistry as drivers of tropical plant litter
322 decomposition. *Biotropica* 36(3): 285-296.
- 323 Bragazza L, Siffi C, Iacumin P, Gerdol R (2007) Mass loss and nutrient release during litter decay in peatland: the
324 role of microbial adaptability to litter chemistry. *Soil Biol Biochem* 39(1):257-267
- 325 Bragazza L, Buttler A, Siegenthaler A, Mitchell E (2009) Plant litter decomposition and nutrient release in
326 peatlands. In: “Carbon cycling in Northern peatlands”, edited by: Baird A, Belyea L, Comas X, Reeve R, Slater
327 L. *Geophysical Monograph*. 184:99-110
- 328 Bragazza L, Buttler A, Robroek BJM, Albrecht R, Zaccone C, Jassey VEJ, Signarbieux C (2016) Persistent high
329 temperature and low precipitation reduce peat carbon accumulation. *Glob Ch Biol* 22, 4114–4123
- 330 Buttler A, Robroek B, Laggoun-Défarge F, Jassey V, Pochelon C, Bernard G, Delarue F, Gogo S, Mariotte P,
331 Mitchell E, Bragazza L (2015) Experimental warming interacts with soil moisture to discriminate plant
332 responses in an ombrotrophic peatland. *J Veg Sci* 26(5):964-974
- 333 Chapman S, Newman G, Hart S, Schweitzer J, Koch G (2013) Leaf litter mixtures alter microbial community
334 development: mechanisms for non-additive effects in litter decomposition. *PLoS ONE* 8(4):e62671
- 335 Delarue F, Laggoun-Défarge F, Buttler A, Gogo S, Jassey V, Disnar J-R (2011) Effects of short-term ecosystem
336 experimental warming on water-extractable organic matter in an ombrotrophic *Sphagnum*-peatland. *Org*
337 *Geochem* 42:1016-1024
- 338 Dieleman, C, Branfireun B, McLaughlin J, Lindo Z (2015) Climate change drives a shift in peatland ecosystem
339 plant community: implications for ecosystem function and stability. *Glob Change Biol* 21(1):388-395
- 340 Francez A-J (2000) La dynamique du carbone dans les tourbières à *Sphagnum*, de la sphaigne à l’effet de serre.
341 *Année Biologique* 39:205-270

342 Frelechoux F, Buttler A, Gillet F (2000) Dynamics of bog pine dominated mire in the Jura mountains,
343 Switzerland: a tentative scheme based on synusial phytosociology. *Folia Geobotanica* 35(3):273-288

344 Gartner, T. B., & Cardon, Z. G. (2004). Decomposition dynamics in mixed-species leaf litter. *Oikos*, 104(2), 230-
345 246.

346 Gogo S, Francez A-J, Laggoun-Défarge F, Gouélibo N, Delarue F, Lottier N (2014) Simultaneous estimation of
347 actual litter enzymatic catalysis and respiration rates with a simple model of C dynamics of *Sphagnum*-
348 Dominated peatlands. *Ecosystems* 17(2):302-316

349 Gogo S, Laggoun-Défarge F, Merzouki F, Mounier S, Guirimand-Dufour A, Jozja N, Huguet A, Delarue F,
350 Défarge C (2016) In situ and laboratory non-additive litter mixture effect on C dynamics of *Sphagnum rubellum*
351 and *Molinia caerulea* litters. *J Soils Sediments* 16(1):13-27

352 Gogo S, Leroy F, Zocatelli R, Bernard-Jannin L, Laggoun-Défarge F (2017) Role of litter decomposition
353 sensitivity to water content in non-additive litter mixture effect: theoretical demonstration and validation with
354 a peatland litter experiment. *Geophysical Research Abstracts* 19: EGU2017-12297, EGU general assembly 2017

355 Gorham, E. (1991). Northern peatlands: role in the carbon cycle and probable responses to climatic warming.
356 *Ecological applications*, 1(2), 182-195.

357 Hobbie S (1996) Temperature and plant species control over litter decomposition in Alaska tundra. *Ecol Monogr*
358 66(4):503-522

359 Hoorens B, Stroetenga M, Aerts R (2010) Litter mixture interactions at the levels of plant functional types are
360 additive. *Ecosystems* 13:90-98

361 Hoorens B, Aerts R, Stroetenga M (2002) Litter quality and interactive effects in litter mixtures: more negative
362 interactions under elevated CO₂? *J Ecol* 90:1009-1016

363 Hu Y, Wang S, Zeng D (2006) Effects of single Chinese fir and mixed leaf litters on soil chemical, microbial
364 properties and soil enzyme activities. *Plant Soil* 282(1-2):379-386

365 Jassey V, Chiapusio G, Binet P, Buttler A, Laggoun-Défarge F, Delarue F, Bernard N, Mitchell E, Toussaint M-
366 L, Francez A-J, Gilbert D (2013) Above- and belowground linkages in *Sphagnum*-peatland: climate warming
367 affects plant-microbial interactions. *Glob Change Biol* 19:811-823

368 Kourtev P, Ehrenfeld J, Huang W (2002) Enzyme activities during litter decomposition of two exotic and two
369 native plant species in hardwood forests of New Jersey. *Soil Biol Biochem* 34(9):1207-1218

370 Krab E, Berg M, Aerts R, van Logtestijn R, Cornelissen J (2013) Vascular plant litter input in subarctic peat bogs
371 changes Collembola diets and decomposition patterns. *Soil Biol Biochem* 63:106-115

372 Laggoun-Défarage F, Gilbert D, Buttler A, Epron D, Francez, A-J, Grasset L, Guimbaud C, Mitchell E, Roy J-C
373 (2008) Effects of experimental warming on carbon sink function of a temperate pristine mire: the PEATWARM
374 project. Proceedings of 13th International Peat Society Congress, June 2008, Tullamore, Ireland, pp 599-602
375 Lecerf A, Marie G, Kominoski J, Leroy C, Bernadet C, Swan C (2011) Incubation time, functional litter diversity,
376 and habitat characteristics predict litter-mixing effects on decomposition. Ecology 92:160-169
377 Limpens J, Berendse F, Blodau C, Canadell J, Freeman C, Holden J, Roulet N, Rydin H, Schaepman-Strub G
378 (2008) Peatlands and the carbon cycle: from local processes to global implications - a synthesis. Biogeosciences
379 5:1475-1491
380 Meier C, Bowman W (2010) Chemical composition and diversity influence non-additive effects of litter mixtures
381 on soil carbon and nitrogen cycling: implications for plant species loss. Soil Biol Biochem 42:1447-1454
382 Rovira P, Rovira R (2010) Fitting litter decomposition datasets to mathematical curves: towards a generalised
383 exponential approach. Geoderma 155:529-543
384 Rydin H, Jeglum J, Jeglum, J (2013) The biology of peatlands, 2nd ed. Oxford University Press
385 Salamanca E F, Kaneko N, Katagiri S (1998) Effects of leaf litter mixtures on the decomposition of *Quercus*
386 *serrata* and *Pinus densiflora* using field and laboratory microcosm methods. Ecol Eng 10(1):53-73
387 Schimel J, Weintraub, M (2003) The implications of exoenzyme activity on microbial carbon and nitrogen
388 limitation in soil: a theoretical model. Soil Biol Biochem 35(4):549-563
389 Sinsabaugh R, Carreiro M, Alvarez S (2002) Enzyme and microbial dynamics of litter decomposition. Enzymes
390 in the Environment, Activity, Ecology and Applications. Marcel Dekker, New York, Basel, pp 249-265
391 Strakova P, Niemi R, Freeman C, Peltoniemi K, Toberman H, Heiskanen I, Fritze H, Laiho R (2011) Litter type
392 affects the activity of aerobic decomposers in a boreal peatland more than site nutrient and water table regimes.
393 Biogeosciences 8:2741-2755
394 Thormann M, Bayley S, Currah R (2004) Microcosm tests of the effects of temperature and microbial species
395 number on the decomposition of *Carex aquatilis* and *Sphagnum fuscum* litter from southern boreal peatlands.
396 Canadian journal of microbiology 50(10): 793-802.
397 Turunen J, Tomppo E, Toloken K, Reinikainen A (2002) Estimating carbon accumulation rates of undrained mires
398 in Finland-application to boreal and subarctic regions. The Holocene 12:69-80
399 Van Breemen N (1995) How *Sphagnum* bogs down other plants. Trends Ecol Evol 10:270-275
400 Vos V, Van Ruijven J, Berg M, Peeters E, Berendse F (2013) Leaf litter quality drives litter mixing effects through
401 complementary resource use among detritivores. Oecologia 173:269-280

- 402 Wu D, Li T, Wan S (2013) Time and litter species composition affect litter-mixing effects on decomposition rates.
403 Plant Soil 371(1-2):355-366
- 404 Yu Z, Loisel J, Brosseau D, Beilman D, & Hunt S (2010) Global peatland dynamics since the Last Glacial
405 Maximum. Geophys Res Lett 37(13)
- 406 Zhang L, Wang H, Zou J, Rogers W, Siemann E (2014) Non-native plant litter enhances soil carbon dioxide
407 emissions in an invaded annual grassland. PLoS one 9(3):e92301
- 408

409 **Table captions**

410 TAB. 1: C content (%), N content (%) and C:N ratio for the *Sphagnum* (S), *Pinus* (P), *Eriophorum* (E), *Sphagnum-*
411 *Pinus* (SP), *Sphagnum-Eriophorum* (SE), *Pinus-Eriophorum* (PE) and *Sphagnum-Pinus-Eriophorum* (SPE) litters.

412

413 TAB 2. Levels of significance from the post-hoc tests for comparison of *Sphagnum-Pinus* (SP), *Sphagnum-*
414 *Eriophorum* (SE), *Pinus-Eriophorum* (PE) and *Sphagnum-Pinus-Eriophorum* (SPE) litter mixture effect (additive
415 vs observed) by incubation time on the remaining solid C, water extractable organic carbon (WEOC) and cumulated
416 of CO₂ - C production during the laboratory experiments. Asterisks represent significant differences (NS = not
417 significant, - p < 0.1, *p < 0.05, **p < 0.01, ***p < 0.001).

418

419 TAB 3. Levels of significance from the post-hoc tests for comparison of *Sphagnum-Pinus-Eriophorum* (SPE) litter
420 mixture effect (additive vs observed) by incubation time on the remaining solid C and CO₂ production during the
421 field experiments. Asterisks represent significant differences (NS = not significant, - p < 0.1, *p < 0.05, **p <
422 0.01, ***p < 0.001).

423

424 TAB. 4: Catalysis rate (mg C .g⁻¹C.d⁻¹) obtained from the C-fluxes model over time for *Sphagnum* (S), *Pinus* (P),
425 *Eriophorum* (E), *Sphagnum-Pinus* (SP), *Sphagnum-Eriophorum* (SE), *Pinus-Eriophorum* (PE) and *Sphagnum-*
426 *Pinus -Eriophorum* (SPE). The model was calibrated with data from the laboratory incubation experiment.
427 Calculate additive rates values are means from the catalysis rates measured in the decomposition of the single
428 species litter.

429 **Figure captions**

430 FIG. 1: Mean (±SE, n=3) remaining solid C (a), water extractable organic carbon (WEOC) (b) and CO₂ - C
431 production (c) in single species litter decomposition of *Sphagnum fallax* (◐, S), *Pinus uncinata* (Δ, P) and
432 *Eriophorum vaginatum* (◊, E) during the laboratory incubation. Lines represent the corresponding fitted curves
433 from the model for each type of litter. Asterisks represent significant differences (NS = not significant, - p < 0.1,
434 *p < 0.05, **p < 0.01, ***p < 0.001).

435 FIG. 2: Additive (■) and observed (□) mean (\pm SE, n=3) of remaining solid C (i), WEOC (ii) and cumulative
436 CO₂ - C (iii) of *Sphagnum+Pinus+Eriophorum* (a), *Sphagnum+Eriophorum* (b), *Pinus+Eriophorum* (c) and
437 *Sphagnum+Pinus* (d) litter mixture decomposition during the laboratory incubation. Additive values are the
438 weighed means from the values measured in the decomposition of the single species litter. Lines represent the
439 corresponding fitted curves from the model. Asterisks represent significant differences (NS = not significant, - p<
440 0.1, *p < 0.05, **p < 0.01, ***p < 0.001).

441 FIG. 3: Mean (\pm SE, n=12) remaining mass (a) and CO₂ production (b) of *Sphagnum fallax* (○, S), *Pinus*
442 *uncinata* (Δ, P) and *Eriophorum vaginatum* (◇, E) litter from field litter bags experiment. Asterisks represent
443 significant differences (NS = not significant, - p < 0.1, *p < 0.05, **p < 0.01, ***p < 0.001).

444 FIG. 4: Additive (■) and observed (□) mean (\pm SE, n=12) of remaining litter mass (a) and CO₂ production (b)
445 from litter mixtures of *Sphagnum*, *Pinus* and *Eriophorum* (SPE) during field litter bags experiment. Additive
446 values were calculated as the weighed mean of the values from litter decomposition of single plant species litter.
447 Asterisks represent significant differences (NS = not significant, - p < 0.1, *p < 0.05, **p < 0.01, ***p < 0.001).

448 FIG. 5: Additive (■) and observed (□) β -glucosidase activities of *Sphagnum+Pinus+Eriophorum*,
449 *Sphagnum+Eriophorum*, *Pinus+Eriophorum* and *Sphagnum+Pinus* litter mixtures (\pm SD, n=3). Asterisks represent
450 significant differences (NS = not significant, - p < 0.1, *p < 0.05, **p < 0.01, ***p < 0.001). Only values from 0
451 to 28 days of incubation are given because a contamination occurred for the β -glucosidase activities samples at 51
452 days.

453

454 **Tables**

455 Table 1.

Composition	Litters						
	S	P	E	SP	SE	PE	SPE
C (%)	42.4	50.9	47.7	46.8	45.2	49.7	46.9
N (%)	1.8	1.7	1.9	1.6	1.9	1.9	1.7
C:N	23.7	29.2	25.8	30.1	23.3	26.9	26.8

456

457 Table 2.

		Main effect		Post hoc test				
		Litter type (expected vs observed)	Time (days)	0	1	14	28	51
SP	Solid C		***					
	WEOC	-						
	Cumul CO ₂ -C		***					
SE	Solid C	***	***					***
	WEOC							
	Cumul CO ₂ -C	***	***					***
PE	Solid C	-	***					
	WEOC		***					
	Cumul CO ₂ -C	-	***					
SPE	Solid C	*	***					
	WEOC	*	***					
	Cumul CO ₂ -C	-	***					

458

459 Table 3

		Main effect		Post hoc test		
		Litter type (expected vs observed)	Time (years)	1	2	3
SPE	Solid C	**	***			*
	CO ₂ production		***			

460

461

462 Table 4

Litters	Time (days)
---------	-------------

		0	1	14	28	51
S		7.87	6.61	1.91	1.66	1.65
P		2.57	2.54	2.18	1.84	1.40
E		4.02	3.51	2.22	2.20	2.20
SP	observed	3.44	3.33	2.34	1.84	1.54
	additive	3.05	5.18	1.83	1.62	1.61
SE	observed	3.57	3.51	2.86	2.34	1.77
	additive	3.41	3.34	2.57	2.00	1.44
PE	observed	3.79	3.5	1.87	1.62	1.59
	additive	3.47	3.26	2.08	1.88	1.85
SPE	observed	3.29	3.23	2.63	2.11	1.46
	additive	3.10	3.05	2.45	1.93	1.31

463

464 **Figures**

465

466

467

468

469

470

471


472


473 **Fig. 1**

474

475

476





478


479

480 Fig. 2

481

482

483



484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513

Fig. 3





Fig. 4



514

515

516

517

518

519

520

521

522

523

524

525

526

527

528

529 Fig. 5

530

531 **Supplementary data**

532 **Table**

533


534 Supplementary Table S5: Percentage normalized root mean square error (% NRMSE) for the solid (L), dissolved
 535 (W), gaseous (G) pools and the sum of them for the *Sphagnum* (S), *Pinus* (P), *Eriophorum* (E), *Sphagnum-Pinus*
 536 (SP), *Sphagnum-Eriophorum* (SE), *Pinus-Eriophorum* (PE) and *Sphagnum-Pinus -Eriophorum* (SPE) models of
 537 C dynamics.

Pools	Litters						
	S	P	E	SP	SE	PE	SPE
L	0.28	0.19	0.11	0.11	0.13	0.09	0.1
W	0.28	4.08	1.89	0.19	0.48	0.84	5.54
G	5.75	5.14	2.49	2.52	2.52	2.51	1.98
Sum	6.31	9.41	4.49	2.82	3.13	3.45	7.61

538

539 **Figures**

540




541

542 Supplementary FIG. S6: Model of the C flow in the litter decomposition process. Three compartments
 543 corresponding to the solid (litter), aqueous (WEOC), and gaseous (cumulative C-CO₂ respired) forms of C are
 544 indicated. Solid lines indicate the rates of catalysis and respiration. The L pool flows into the WEOC at the catalysis
 545 rate “c” and the WEOC is respired at the rate “r” (adapted from Gogo et al. 2014).


546

547



548

549 Supplementary FIG. S7: β -glucosidase activities in *Sphagnum* (\circ), *Pinus* (Δ) and *Eriophorum* (\diamond) litter during
 550 the laboratory incubation (\pm SD, n=3). Only values from 0 to 28 days of incubation are given because a
 551 contamination occurred for the β -glucosidase activities samples at 51 days.



552

553 Supplementary FIG. S8: Linear regression between catalysis rates and β -glucosidase activities. Each point
 554 represents the β -glucosidase activities measured associated to the catalysis rate obtained for each litter type at a
 555 time t . The r^2 was calculated with all litters at the exception of *Sphagnum* ones (\bullet).