

PAPER • OPEN ACCESS

First release of *LiteBIRD* simulations from an end-to-end pipeline

To cite this article: M. Bortolami *et al* JCAP11(2025)042

View the [article online](#) for updates and enhancements.

You may also like

- [Search for Lensing Signatures in the Gravitational-Wave Observations from the First Half of LIGO–Virgo’s Third Observing Run](#)
R. Abbott, T. D. Abbott, S. Abraham *et al.*
- [Sensor response and radiation damage effects for 3D pixels in the ATLAS IBL Detector](#)
G. Aad, E. Aakvaag, B. Abbott *et al.*
- [Prospects for a survey of the galactic plane with the Cherenkov Telescope Array](#)
S. Abe, J. Abhir, A. Abhishek *et al.*

First release of *LiteBIRD* simulations from an end-to-end pipeline



The LiteBIRD collaboration

M. Bortolami,^{1,2} N. Raffuzzi,¹ L. Pagano,^{1,2,3} G. Puglisi,^{4,5,6} A. Anand,⁷ A.J. Banday,⁸ P. Campeti,^{2,9,10} G. Galloni,^{1,7} A.I. Lonappan,¹¹ M. Monelli,¹² M. Tomasi,^{13,14} G. Weymann-Despres,^{15,16} D. Adak,¹⁷ E. Allys,¹⁸ J. Aumont,⁸ R. Aurvik,¹⁹ C. Baccigalupi,^{20,21,22} M. Ballardini,^{1,2,23} R.B. Barreiro,²⁴ N. Bartolo,^{25,26,27} S. Basak,²⁸ M. Bersanelli,^{13,14} A. Besnard,³ T. Brinckmann,¹ E. Calabrese,²⁹ E. Carinos,⁸ A. Carones,^{20,21} F.J. Casas,²⁴ K. Cheung,^{30,31,32,33} M. Citran,³⁴ L. Clermont,³⁵ F. Columbro,^{36,37} G. Coppi,³⁸ A. Coppolecchia,^{36,37} F. Cuttaia,²³ P. Dal Bo,³⁹ P. de Bernardis,^{36,37} E. de la Hoz,⁴⁰ M. De Lucia,³⁹ S. Della Torre,⁴¹ P. Diego-Palazuelos,⁹ H.K. Eriksen,¹⁹ T. Essinger-Hileman,⁴² C. Franceschet,^{13,14} U. Fuskeland,¹⁹ M. Galloway,¹⁹ M. Gerbino,² M. Gervasi,^{38,41} T. Ghigna,^{43,12} S. Giardiello,²⁹ C. Gimeno-Amo,²⁴ E. Gjerløw,¹⁹ A. Gruppuso,^{23,44} M. Hazumi,^{45,46,12,47} S. Henrot-Versillé,¹⁶ L.T. Hergt,^{48,16} E. Hivon,⁴⁹ B. Jost,¹² K. Kohri,⁴⁵ L. Lamagna,^{36,37} T. Lari,³⁹ M. Lattanzi,² C. Leloup,¹² F. Levrier,¹⁸ M. López-Caniego,^{50,51} G. Luzzi,⁵² J. Macias-Perez,⁵³ B. Maffei,³ E. Martínez-González,²⁴ S. Masi,^{36,37} S. Matarrese,^{25,26,27,54} T. Matsumura,¹² S. Micheli,³⁶ L. Montier,⁸ G. Morgante,²³ L. Mousset,^{18,8} R. Nagata,⁴⁶ T. Namikawa,¹² A. Novelli,³⁶ F. Noviello,²⁹ I. Obata,⁴⁵ A. Occhiuzzi,³⁶ A. Paiella,^{36,37} D. Paoletti,^{23,44} G. Pascual-Cisneros,^{12,24} F. Piacentini,^{36,37} M. Pinchera,³⁹ G. Polenta,⁵² L. Porcelli,⁵⁵ M. Remazeilles,²⁴ A. Ritacco,⁵³ A. Rizzieri,^{15,34} J.A. Rubiño-Martín,^{17,56} M. Ruiz-Granda,^{24,57} J. Sanghavi,^{58,59} V. Sauvage,³ M. Shiraishi,⁶⁰ G. Signorelli,^{61,39} S.L. Stever,^{3,62,12} R.M. Sullivan,¹⁹ Y. Takase,⁶² K. Tassis,^{63,64} L. Terenzi,²³ M. Tristram,¹⁶ L. Vacher,²⁰ B. van Tent,¹⁶ P. Vielva,²⁴ I.K. Wehus,¹⁹ M. Zannoni,^{38,41} and Y. Zhou⁴³

¹Dipartimento di Fisica e Scienze della Terra, Università di Ferrara,
Via Saragat 1, 44122 Ferrara, Italy

²INFN Sezione di Ferrara, Via Saragat 1, 44122 Ferrara, Italy

³Université Paris-Saclay, CNRS, Institut d'Astrophysique Spatiale, 91405, Orsay, France

- ⁴*Dipartimento di Fisica e Astronomia, Università degli Studi di Catania,
Via S. Sofia,64, 95123, Catania, Italy*
- ⁵*INAF, Osservatorio Astrofisico di Catania, via S.Sofia 78, I-95123 Catania, Italy*
- ⁶*INFN, Sezione di Catania, via S.Sofia 64, I-95123, Catania, Italy*
- ⁷*Dipartimento di Fisica, Università di Roma Tor Vergata,
Via della Ricerca Scientifica, 1, 00133, Roma, Italy*
- ⁸*IRAP, Université de Toulouse, CNRS, CNES, UPS, Toulouse, France*
- ⁹*Max Planck Institute for Astrophysics, Karl-Schwarzschild-Str. 1, D-85748 Garching, Germany*
- ¹⁰*Excellence Cluster ORIGINS, Boltzmannstr. 2, 85748 Garching, Germany*
- ¹¹*University of California,
San Diego, Department of Physics, San Diego, CA 92093-0424, U.S.A.*
- ¹²*Kavli Institute for the Physics and Mathematics of the Universe (Kavli IPMU, WPI), UTIAS,
The University of Tokyo,
Kashiwa, Chiba 277-8583, Japan*
- ¹³*Dipartimento di Fisica, Università degli Studi di Milano,
Via Celoria 16 - 20133, Milano, Italy*
- ¹⁴*INFN Sezione di Milano, Via Celoria 16 - 20133, Milano, Italy*
- ¹⁵*Department of Physics, University of Oxford,
Denys Wilkinson Building, Keble Road, Oxford OX1 3RH, U.K.*
- ¹⁶*Université Paris-Saclay, CNRS/IN2P3, IJCLab, 91405 Orsay, France*
- ¹⁷*Instituto de Astrofísica de Canarias, E-38200 La Laguna, Tenerife, Canary Islands, Spain*
- ¹⁸*Laboratoire de Physique de l'École Normale Supérieure, ENS, Université PSL, CNRS,
Sorbonne Université, Université de Paris,
75005 Paris, France*
- ¹⁹*Institute of Theoretical Astrophysics, University of Oslo,
Blindern, Oslo, Norway*
- ²⁰*International School for Advanced Studies (SISSA), Via Bonomea 265, 34136, Trieste, Italy*
- ²¹*INFN Sezione di Trieste, via Valerio 2, 34127 Trieste, Italy*
- ²²*IFPU, Via Beirut, 2, 34151 Grignano, Trieste, Italy*
- ²³*INAF - OAS Bologna, via Piero Gobetti, 93/3, 40129 Bologna, Italy*
- ²⁴*Instituto de Física de Cantabria (IFCA, CSIC-UC), Avenida los Castros SN, 39005, Santander,
Spain*
- ²⁵*Dipartimento di Fisica e Astronomia "G. Galilei", Università degli Studi di Padova,
via Marzolo 8, I-35131 Padova, Italy*
- ²⁶*INFN Sezione di Padova, via Marzolo 8, I-35131, Padova, Italy*
- ²⁷*INAF, Osservatorio Astronomico di Padova, Vicolo dell'Osservatorio 5, I-35122, Padova, Italy*
- ²⁸*School of Physics, Indian Institute of Science Education and Research Thiruvananthapuram,
Maruthamala PO, Vithura, Thiruvananthapuram 695551, Kerala, India*
- ²⁹*School of Physics and Astronomy, Cardiff University,
Cardiff CF24 3AA, U.K.*
- ³⁰*Jodrell Bank Centre for Astrophysics, Alan Turing Building, Department of Physics and
Astronomy, School of Natural Sciences, The University of Manchester,
Oxford Road, Manchester M13 9PL, U.K.*
- ³¹*Department of Physics, University of California, Berkeley,
Berkeley, CA 94720, U.S.A.*

- ³²*Space Sciences Laboratory, University of California, Berkeley, Berkeley, CA 94720, U.S.A.*
- ³³*Lawrence Berkeley National Laboratory (LBNL), Computational Cosmology Center, Berkeley, CA 94720, U.S.A.*
- ³⁴*Université Paris Cité, CNRS, Astroparticule et Cosmologie, F-75013 Paris, France*
- ³⁵*Centre Spatial de Liège, Université de Liège, Avenue du Pré-Aily, 4031 Angleur, Belgium*
- ³⁶*Dipartimento di Fisica, Università La Sapienza, P. le A. Moro 2, Roma, Italy*
- ³⁷*INFN Sezione di Roma, P.le A. Moro 2, 00185 Roma, Italy*
- ³⁸*Physics Department, University of Milano Bicocca, p.zza della Scienza, 3, 20126 Milan, Italy*
- ³⁹*INFN Sezione di Pisa, Largo Bruno Pontecorvo 3, 56127 Pisa, Italy*
- ⁴⁰*CNRS-UCB International Research Laboratory, Centre Pierre Binétruy, UMI2007, Berkeley, CA 94720, U.S.A.*
- ⁴¹*INFN Sezione Milano Bicocca, Piazza della Scienza, 3, 20126 Milano, Italy*
- ⁴²*NASA Goddard Space Flight Center, Greenbelt, MD 20771, U.S.A.*
- ⁴³*International Center for Quantum-field Measurement Systems for Studies of the Universe and Particles (QUP), High Energy Accelerator Research Organization (KEK), Tsukuba, Ibaraki 305-0801, Japan*
- ⁴⁴*INFN Sezione di Bologna, Viale C. Berti Pichat, 6/2 – 40127 Bologna, Italy*
- ⁴⁵*Institute of Particle and Nuclear Studies (IPNS), High Energy Accelerator Research Organization (KEK), Tsukuba, Ibaraki 305-0801, Japan*
- ⁴⁶*Japan Aerospace Exploration Agency (JAXA), Institute of Space and Astronautical Science (ISAS), Sagamihara, Kanagawa 252-5210, Japan*
- ⁴⁷*The Graduate University for Advanced Studies (SOKENDAI), Miura District, Kanagawa 240-0115, Hayama, Japan*
- ⁴⁸*Department of Physics and Astronomy, University of British Columbia, 6224 Agricultural Road, Vancouver, BC V6T1Z1, Canada*
- ⁴⁹*Institut d’Astrophysique de Paris, CNRS/Sorbonne Université, Paris, France*
- ⁵⁰*Aurora Technology for the European Space Agency, Camino bajo del Castillo, s/n, Urbanización Villafranca del Castillo, Villanueva de la Cañada, Madrid, Spain*
- ⁵¹*Universidad Europea de Madrid, 28670, Madrid, Spain*
- ⁵²*Space Science Data Center, Italian Space Agency, via del Politecnico, 00133, Roma, Italy*
- ⁵³*Université Grenoble Alpes, CNRS, LPSC-IN2P3, 53, avenue des Martyrs, 38000 Grenoble, France*
- ⁵⁴*Gran Sasso Science Institute (GSSI), Viale F. Crispi 7, I-67100, L’Aquila, Italy*
- ⁵⁵*Istituto Nazionale di Fisica Nucleare-Laboratori Nazionali di Frascati (INFN-LNF), Via E. Fermi 40, 00044, Frascati, Italy*
- ⁵⁶*Departamento de Astrofísica, Universidad de La Laguna (ULL), E-38206, La Laguna, Tenerife, Spain*
- ⁵⁷*Dpto. de Física Moderna, Universidad de Cantabria, Avda. los Castros s/n, E-39005 Santander, Spain*
- ⁵⁸*Universitäts-Sternwarte, Fakultät für Physik, Ludwig-Maximilians Universität München, Scheinerstr.1, 81679 München, Germany*
- ⁵⁹*GRAPPA, Institute for Theoretical Physics Amsterdam, University of Amsterdam, Science Park 904, 1098 XH Amsterdam, The Netherlands*

⁶⁰*Suwa University of Science, Chino, Nagano 391-0292, Japan*

⁶¹*Dipartimento di Fisica, Università di Pisa,
Largo B. Pontecorvo 3, 56127 Pisa, Italy*

⁶²*Department of Physics, Okayama University, Okayama 700-8530, Japan*

⁶³*Institute of Astrophysics, Foundation for Research and Technology – Hellas,
Vasilika Vouton, GR-70013 Heraklion, Greece*

⁶⁴*Department of Physics and ITCP, University of Crete, GR-70013, Heraklion, Greece*

E-mail: marco.bortolami@unife.it, nicolelia.raffuzzi@unife.it,
luca.pagano@unife.it, giuseppe.puglisi2@unict.it

ABSTRACT: The *LiteBIRD* satellite mission aims at detecting Cosmic Microwave Background *B* modes with unprecedented precision, targeting a total error on the tensor-to-scalar ratio r of $\delta r \sim 0.001$. Operating from the L2 Lagrangian point of the Sun-Earth system, *LiteBIRD* will survey the full sky across 15 frequency bands (34 to 448 GHz) for 3 years. The current *LiteBIRD* baseline configuration employs 4508 detectors sampling at 19.1 Hz to achieve an effective polarization sensitivity of $2 \mu\text{K arcmin}$ and an angular resolution of 31 arcmin (at 140 GHz).

We describe the first release of the official *LiteBIRD* simulations, realized with a new simulation pipeline developed using the *LiteBIRD* Simulation Framework. This pipeline generates 500 full-sky simulated maps at a HEALPix resolution of $N_{\text{side}}=512$. The simulations include also one year of Time Ordered Data (TOD) for approximately one-third of *LiteBIRD*'s total detectors.

KEYWORDS: gravitational waves and CMBR polarization, CMBR experiments, CMBR polarisation

ARXIV EPRINT: [2507.07122](https://arxiv.org/abs/2507.07122)

Contents

1	Introduction	1
2	Instrument model	2
3	Input sky maps	5
4	Simulation pipeline	6
5	Results and validation	9
5.1	TOD	10
5.2	Covariance matrices	11
5.3	Power spectra	14
5.4	Computational cost	14
6	Conclusions	15
A	Products of the pipeline	17
B	Comparison between input and output maps	18

1 Introduction

The standard cosmological model describing our Universe, referred to as the Λ Cold Dark Matter (Λ CDM) model, with Λ referring to the Dark Energy, is strongly supported by the observations of Cosmic Microwave Background (CMB) temperature and polarization anisotropies. A primordial phase of rapid and exponential expansion, known as inflation [1], is currently associated with the mechanism producing primordial density fluctuations together with perturbations of the fabric of space-time, in the form of primordial gravitational waves. While these density fluctuations are associated with the curl-free component of the CMB polarization, E modes, the gravitational waves left a divergence-less pattern and are referred to as B modes [2].

To date, E modes have been characterized by several ground [3–6] and space-based [7, 8] experiments. However, primordial B modes have not yet been detected; only upper limits have been established on their amplitudes, with the tensor-to-scalar ratio, r (the ratio of power in primordial gravitational waves to primordial density perturbations), constrained to $r < 0.032$ at 95% confidence level by [9]. Instrumental systematics and sensitivity together with polarized microwave emission from our own Galaxy pose a challenge for the detection of cosmological B -modes.

The Lite (Light) satellite for the study of B -mode polarization and Inflation from cosmic background Radiation Detection (*LiteBIRD*) is a satellite mission that aims at detecting the CMB B modes with a total error on the tensor-to-scalar ratio of $\delta r \sim 0.001$ [10]. It

will observe the full sky in 15 frequency bands from 34 to 448 GHz for 3 years from the L2 Lagrangian point of the Sun-Earth system, with effective polarization sensitivity of $2.2 \mu\text{K} - \text{arcmin}$ and angular resolution of 31 arcmin (at 140 GHz), employing 4508 detectors sampling at 19.1 Hz [10].

In order to achieve the B -mode requirements on r , the *LiteBIRD* mission requires unprecedented control of instrumental systematics of a typical CMB satellite. Consequently, generating realistic *LiteBIRD* observation simulations that account for these systematic effects are crucial. These simulations are decisive for establishing stringent requirements on instrument parameters associated with systematic errors.

To validate the design of its instruments, the *LiteBIRD* collaboration has developed a simulation pipeline. This computer program emulates the spacecraft’s instrument operations, generating a synthetic data set that mirrors the anticipated *LiteBIRD* observations. Simulation pipelines are crucial for assessing the feasibility of achieving scientific objectives, optimizing observation strategies, and validating ground segment software and data storage systems. Developing a simulation pipeline involves defining scientific goals and instrument requirements, creating a mathematical model of the instrument, translating the mathematical model into a numerical model, developing and implementing the simulation pipeline software, and rigorously validating the accuracy and reliability of the pipeline.

We thus used the *LiteBIRD* Simulation Framework (LBS,¹ [11]) and developed an end-to-end simulation pipeline, designed to generate 500 full sky simulated maps at HEALPix² [12, 13] resolution $N_{\text{side}} = 512$. Given the rapid evolution of the LBS framework due to active community contributions, we note that these simulations utilized version 0.11.0. While the latest version is 0.13.0 at the time of writing, we have ensured backward compatibility with our end-to-end pipeline.

Additionally, the pipeline produces a set of TODs using the *LiteBIRD* Instrument Model database (IMo) that is made available to *LiteBIRD* collaborators for several applications. In particular, we highlight that simulated TODs have been employed to assess the computational feasibility of an end-to-end Bayesian analysis of the *LiteBIRD* experiment within the Cosmoglobe framework [14]. They will be presented in a companion paper to be soon submitted [15]. This paper is organised as follows. In section 2 we describe the instrument model adopted for this first release of simulations. In section 3 we present the beam convolved CMB and foreground maps used as input for the simulation pipeline. In section 4 we illustrate the simulation pipeline. In section 5 we show the outputs of the pipeline along with their validation. In section 6 we draw conclusions. Appendix A lists all products of the pipeline, appendix B shows a comparison between the input and the scanned output maps.

2 Instrument model

For the simulations presented in this work, we employ the official *LiteBIRD* Instrument Model (IMo) database, which provides a detailed description of the current baseline instrument design (version v1.3). *LiteBIRD* is organized into 3 separate telescopes: the Low Frequency

¹https://github.com/litebird/litebird_sim.

²<http://healpix.sourceforge.net>.




Figure 1. Detectors present in *LiteBIRD*'s LFT (left), MFT (middle) and HFT (right). The colors highlight wafers with the same frequency. The black stars indicate the selected pixels.

Telescope (LFT), Medium Frequency Telescope (MFT), and High Frequency Telescope (HFT). The former employs reflective optics, whereas the latter are refractors.

As these simulations are mainly meant to be a pathfinder, we consider 1/3 of the nominal mission time (1 instead of 3 years) and of the detectors in the focal plane. In particular, we used all of the detectors for those channels with a maximum number of 48, whereas for the rest we selected detectors in a configuration preserving the focal plane symmetry. Figure 1 shows the LFT, MFT and HFT focal planes (represented by colored circles), and we marked the camera pixels considered for the simulations with a black star. We remind the reader that each pixel in the focal plane is associated with 2 bolometer detectors.

We report in the first column of table 1 the names of the 22 *LiteBIRD* channels. Table 1 also summarises nominal specifics of each frequency channel employed for our simulations, i.e. the nominal number of detectors $n_{\text{det}}^{\text{IMo}}$, the number of detectors employed in the end-to-end simulation $n_{\text{det}}^{\text{e2e}}$, the beam Full Width Half Maximum (FWHM) in arcminutes, the detector Noise Equivalent Temperature (NET) values NET^{IMo} in $\mu\text{K}\sqrt{s}$ and the detector NET actually employed in the simulations in $\mu\text{K}\sqrt{s}$. The noise levels we use in the simulation, NET^{e2e} , correspond to the IMo noise levels, NET^{IMo} , rescaled to imitate the full focal plane 3-year mission as

$$\text{noise rescaling} = \sqrt{\frac{1}{3\epsilon} \frac{n_{\text{det}}^{\text{e2e}}}{n_{\text{det}}^{\text{IMo}}}}, \quad (2.1)$$

where the 1/3 takes into account the reduced simulated time, $n_{\text{det}}^{\text{e2e}}/n_{\text{det}}^{\text{IMo}}$ the reduced number of simulated detectors and ϵ is the detector efficiency. Given the goals of this round of simulations, we do not include any optical efficiency and assume an ideal detector yield.

For the scanning strategy parameters in these simulations we consistently employ the ones presented in [10] and summarized in table 2 and figure 2. The telescope boresight is positioned at an angle of $\beta = 50^\circ$ relative to the spin-axis, rotating at a rate of 0.05 rpm (20 min.). The spin-axis undergoes precession around the Sun-Earth direction with an angle of $\alpha = 45^\circ$, completing one full rotation in ~ 3.2 hours. Combining three motions (spin-axis rotation, precession around the Sun-Earth axis and 1-year revolution around the Sun) enables the boresight to cover the entire sky within six months. For further details about the *LiteBIRD* scanning strategy refer to [16].

Channel	$n_{\text{det}}^{\text{IMo}}$	$n_{\text{det}}^{\text{e2e}}$	Beam	NET^{IMo}	NET^{e2e}
			[arcmin]	$[\mu\text{K}\sqrt{\text{s}}]$	$[\mu\text{K}\sqrt{\text{s}}]$
L1-040	48	48	70.5	114.6	66.2
L1-060	48	48	51.1	65.3	37.7
L1-078	48	48	43.8	58.6	33.8
L2-050	24	24	58.5	72.5	41.9
L2-068	24	24	47.1	68.8	39.7
L2-089	24	24	41.5	62.3	36.0
L3-068	144	48	41.6	105.6	35.2
L3-089	144	48	33.0	65.2	21.7
L3-119	144	48	26.3	40.8	13.6
L4-078	144	48	36.9	82.5	27.5
L4-100	144	48	30.2	54.9	18.3
L4-140	144	48	23.7	38.4	12.8
M1-100	366	126	37.8	71.7	24.3
M1-140	366	126	30.8	54.0	18.3
M1-195	366	126	28.0	59.6	20.2
M2-119	488	168	33.6	55.7	18.9
M2-166	488	168	28.9	54.4	18.4
H1-195	254	86	28.6	74.0	24.9
H1-280	254	86	22.5	97.3	32.7
H2-235	254	86	24.7	76.1	25.6
H2-337	254	86	20.9	154.6	52.0
H3-402	338	116	17.9	385.6	130.5

Table 1. Channel characteristics. The nomenclature in the first column expresses the telescope identifier (first letter for Low, Medium or High Frequency Telescope), wafer number (first digit), and central frequency (in GHz) each channel is sensitive to. For each channel, $n_{\text{det}}^{\text{IMo}}$ is the number of detectors reported in the IMo (totaling 4508), $n_{\text{det}}^{\text{e2e}}$ is the number of detectors employed for our simulations (totaling 1678), Beam is the beam FWHM in arcminutes, NET^{IMo} is the noise requirement from the IMo, NET^{e2e} is the noise baseline used for our simulations (in units of $\mu\text{K}\sqrt{\text{s}}$).

α	β	Precession period	Spin rate	Sampling rate	HWP rotation rate [rpm]		
					LFT	MFT	HFT
[deg.]	[deg.]	[min.]	[rpm]	[Hz]			
45	50	192.348	0.05	19.1	46	39	61

Table 2. Parameters of the observation strategy and sampling rate of the *LiteBIRD* satellite.




Figure 2. Cartoon representation of the *LiteBIRD* satellite and its scanning parameters. $\alpha = 45^\circ$ is the angle between the spin axis and the Sun-Earth axis, $\beta = 50^\circ$ is the angle separating the boresight from the spin axis. A summary of the scanning parameters is in table 2.

3 Input sky maps

Our simulation pipeline takes as input a set of (I, Q, U) sky maps at HEALPix [12, 13] resolution $N_{\text{side}} = 512$ for each of the 22 *LiteBIRD* frequency channels. We generate the input maps of CMB, Galactic and extragalactic foregrounds separately. The different tools we use to simulate the different components are presented below.

We employ the Map-Based Simulation (MBS) module within the LBS framework to generate maps of Galactic diffuse foregrounds. MBS serves as a wrapper around the Python Sky Model (PySM)³ package [17–19]. We consider the following models:

- d1** thermal dust modelled as a single-component modified black-body with varying temperature and spectral index across the sky, based on the maps from the Planck 2015 analysis [20];
- s1** synchrotron modelled as a power law model with varying spectral index and no curvature, based on Haslam 408 MHz [21] and WMAP data [22, 23];
- a1** two unpolarised spinning dust populations of Anomalous Microwave Emission (AME), based on the templates obtained from *Planck* with Commander methodology [20];
- f1** unpolarised free-free emission with a constant spectral index of -2.14 , based on the templates obtained from *Planck* with Commander methodology [20];

³<https://github.com/galsci/pysm>.

co1 $J : 1 \rightarrow 0, 2 \rightarrow 1, 3 \rightarrow 2$ rotational lines of Galactic CO emission, whose center frequency is 115.3, 230.5 and 345.8 GHz, respectively [24, 25].

We use the WebSky simulations [26] and PySM to generate the maps of extragalactic emissions across the *LiteBIRD* frequency band. We produce maps of thermal (tSZ) and kinetic Sunyaev-Zeldovich (kSZ) emission, Cosmic Infrared Background (CIB) and lensing convergence by populating Dark Matter haloes from an N-body simulation (for further details please refer to [26]). The tSZ effect arises from the inverse Compton scattering of CMB photons off hot electrons in galaxy clusters, distorting the CMB spectrum [27]. The kSZ effect, on the other hand, is caused by the Doppler shift of CMB photons due to the motion of these same galaxy clusters relative to the CMB rest frame [28]. The CIB is produced by the cumulative emission of dust-covered star-forming galaxies across cosmic time [29]. The lensing convergence map represents the projected mass density along the line of sight [30], decisive for modeling and correcting the distortions in the CMB caused by gravitational lensing, allowing for a more accurate reconstruction of the matter distribution and extraction of cosmological information. We also use recent radio sources catalogs to produce frequency-dependent maps [31–33].

We produce 500 Monte Carlo (MC) realisations of the CMB sky from the *Planck* 2018 cosmological parameters [34], lensed with the same WebSky convergence map via the `lenspyx`⁴ package [35]. By using the beam specifications provided by the IMo and reported in table 1, we convolve the input CMB and foreground maps. Then, by scanning the maps, we produce TOD that already include the response to instrumental beams. The beams are assumed to be circular Gaussian with the same FWHM for all detectors in a given channel. We show in figure 3 the coadded input maps of CMB and foregrounds for three different channels chosen from the three telescopes.

4 Simulation pipeline

The pipeline developed for the production of the simulations heavily relies on the LBS, integrated with the IMo described in section 2. The initial step of the pipeline involves loading the instrument parameters summarized in tables 1, 2, including the quaternions describing the rotation from the Ecliptic reference frame to the reference frame of each detector, the rotation frequency of the HWPs⁵ (46/39/61 rpm for LFT/MFT/HFT) and the detector noise levels.

These quantities are firstly used to calculate the pointing information for each simulated detector and sample, employing the *LiteBIRD* scanning strategy (see [10] for further details) and the Ephemeridis tables to get an accurate position of the spacecraft around the Sun at each instant. In particular, we calculate the colatitude θ and longitude ϕ as well as the polarization angle ψ .

⁴<https://github.com/carronj/lenspyx>.

⁵The polarized incident radiation is modulated at four times the rotation frequency of the HWP, effectively shifting the polarized sky signal to a narrow frequency band, specifically above the $1/f$ noise knee frequency. This modulation significantly improves the measurement of CMB polarization by separating the sky signal from the $1/f$ noise.




Figure 3. Coadded input maps in Galactic coordinates. T (left column), Q (middle column), U (right column) coadded input maps of CMB and foregrounds for the first simulation. First row: L1-040 channel of LFT. Second row: M1-140 channel of MFT. Third row: H3-402 channel of HFT.

Subsequently, the beam convolved input maps presented in section 3 are scanned (see appendix B and figure 11) using the pointing information to generate TOD for CMB (tod_cmb) and foregrounds (tod_fg). Additionally, three noise timelines are simulated using the detector noise characteristics $\text{NET}^{\text{e}2\text{e}}$ reported in table 1: tod_wn , which consists of white noise only; tod_wn_1f_30mHz and tod_wn_1f_100mHz , containing white noise and correlated $1/f$ noise with a knee frequency f_{knee} of 30 mHz and 100 mHz, respectively. The tod_wn_1f_30mHz timeline represents a realistic scenario, with a knee frequency close to what is expected from *LiteBIRD*. The tod_wn_1f_100mHz timeline is instead a pessimistic case for which the $1/f$ noise is worse and it is useful for setting requirements in pessimistic conditions. A summary of all products may be found in the appendix A.

We used FFTs to generate the $1/f$ noise, leveraging a simulation length equal to the length of the TOD, which corresponds to one year of data at a sampling rate of 19.1 Hz. To handle the inherent limitation of FFTs, which restrict the simulation to the length of the TOD, we selected a simulation length equal to the TOD length (one year) and padded it

to the nearest power of 2 to optimize FFT performance. By ensuring that the FFT length matches the data length, we avoid issues with periodicity. The decision to use FFTs for generating $1/f$ noise was driven by its computational efficiency, especially when compared to the alternative approach of Markov random walks, which, although more flexible for very low frequencies, would have been significantly more computationally demanding. In frequency domain the $1/f$ noise is:

$$\tilde{d}_i \rightarrow \tilde{d}_i \times \sigma \sqrt{\frac{f_i^\alpha + f_{\text{knee}}^\alpha}{f_i^\alpha + f_{\text{min}}^\alpha}} \quad \text{for } i > 0, \quad \text{with } \tilde{d}_0 = 0 \quad (4.1)$$

where \tilde{d}_i is the Fourier transform of the TOD at frequency index i , f_i is the frequency, f_{knee} is the knee frequency where the $1/f$ noise power equals the white noise power in the power spectrum density (see figure 7), f_{min} is the low-frequency cutoff, σ is the rescaled white noise level (NET^{e2e} in table 1) and $\alpha = 1$ is the low-frequency spectral tilt.

For the first simulation only, we produce a timestream `tod_dip` with the CMB dipole signal. We simulate the dipole signal using the `TOTAL_FROM_LIN_T` model,⁶ following the LBS nomenclature. This is described by:

$$\Delta T = \frac{T_0}{f(x)} \left(\frac{B(\nu\gamma(1 - \vec{\beta} \cdot \hat{n}), T_0)}{(\gamma(1 - \vec{\beta} \cdot \hat{n}))^3 B(\nu, T_0)} \right), \quad (4.2)$$

where ΔT is the dipole signal, T_0 is the CMB temperature [36], $\vec{\beta} = \vec{v}/c$ is the velocity vector (relative to the speed of light c) accounting for Earth's motion in the CMB rest frame, $\gamma = (1 - \vec{\beta} \cdot \vec{\beta})^{-1/2}$, \hat{n} is the line-of-sight direction, ν is the CMB frequency, $x = h\nu/k_B T$ so that:

$$f(x) = \frac{x e^x}{e^x - 1} \quad (4.3)$$

where h is the Planck constant and k_B is the Boltzmann constant, and the black-body spectrum B is given by:

$$B(\nu, T) = \frac{2h\nu^3}{c^2} \frac{1}{e^{h\nu/k_B T} - 1} = \frac{2h\nu^3}{c^2} \frac{1}{e^x - 1}. \quad (4.4)$$

A biner map-making algorithm⁷ is applied to the TOD to produce maps containing CMB, foregrounds, white noise, and $1/f$ noise at a `HEALPix` [12, 13] resolution of $N_{\text{side}} = 512$. Although the biner map-maker inherently assumes uncorrelated noise in the timelines, the presence of the HWP in the polarization case is expected to effectively suppress the $1/f$ noise component. For temperature, the $1/f$ noise becomes negligible due to the extremely favorable signal-to-noise ratio (SNR), as illustrated in figure 4. The top left plot compares the CMB temperature with the noise power spectra. The bottom left panel depicts the SNRs

⁶<https://litebird-sim.readthedocs.io/en/master/dipole.html>.

⁷A biner (or naive) map-maker constructs sky maps from TODs by averaging observed data points into pixels on a sky map. This approach does not incorporate specific noise modeling (e.g., assuming only white noise) or deconvolve instrument effects. While computationally efficient, it is less accurate compared to advanced techniques such as maximum-likelihood or optimal map-makers.

of both levels of $1/f$ noise with respect to the white noise baseline. The SNR is computed as:

$$\text{SNR} = \frac{C_\ell}{\sigma} \quad (4.5)$$

$$\sigma^2 = \frac{2}{2\ell + 1} (C_\ell + N_\ell)^2.$$

The noise power spectra in both panels for the $1/f$ noise levels are obtained by inversely coadding the spectra of the binned output and input map differences across all channels, and averaged over 50 simulations:

$$C_\ell = \left[\sum_{\text{chann}} \frac{1}{C_{\ell, \text{chann}}} \right]^{-1}. \quad (4.6)$$

As a test case, we consider a scenario analogous to the temperature case but without the presence of a HWP. The polarization noise is derived by doubling the power of the temperature noise. This approach is consistent with the observed increase in noise levels for polarization, as demonstrated in figure 8 and by the green curves in figure 9.

We acknowledge that, at the time of submission, a bug was identified in the noise time-stream generation code. This resulted in the unintentional simulation of a higher level of $1/f$ noise, mainly affecting the temperature simulations. However, as evidenced by the unprecedented *LiteBIRD* signal-to-noise ratios presented in figure 4, the impact of this error is considered minor for both polarization and temperature data.

The binner is utilized for all the simulations, in particular producing 500 maps for each of the two cases, with $f_{\text{knee}} = 30$ mHz and with $f_{\text{knee}} = 100$ mHz. Furthermore, only for the first simulation, we stored the TODs, pointing information and noise covariance matrices. In detail, we generated a noise covariance matrix (in pixel space) for each channel, which is crucial for realistic map-based simulations that incorporate pixel-to-pixel correlations in the I , Q , and U maps. Map-based simulations are indispensable for testing and validating the observational system's performance, assessing systematic effects, and ensuring the robustness of data analysis pipelines.

Each result presented in this work is fully reproducible, provided that the noise characteristics outlined in table 1 are adhered to. The repository employing the simulation script and the configuration file has been made publicly available.⁸ The public version⁹ of the IMO is implemented in the LBS, allowing for easy replication of the noise properties. However, while the noise features are reproducible, variations in the focal plane geometry are not fully captured in the IMO version of the code.

5 Results and validation

We present here the results obtained with the simulation pipeline along with their validation. In section 5.1 we show results related to TODs, e.g. their plot for different detectors or components. In appendix B, we present the output maps and their difference with the input ones. In section 5.2, we show the noise covariance matrices. In section 5.3, we discuss the power spectra calculated from the maps. Finally, in section 5.4 we report the computational cost.

⁸<https://github.com/litebird/e2e-simulation>.

⁹<https://github.com/litebird/instrumentdb>.




Figure 4. Top panels: the CMB power spectra for temperature and polarization are compared to the noise spectra in the absence of a half-wave plate (HWP). The fiducial signal is derived from the best-fit cosmological parameters of the *Planck* 2018 data [34], obtained from the TT, TE, EE + lowE + lensing analysis. The noise power spectra are computed as the spectrum of the difference between the binned output and input maps, inversely coadded across all channels and averaged over the first 50 simulations. The various curves represent different levels of $1/f$ noise knee frequencies, along with the white noise-only baseline. For polarization (without HWP), the noise is estimated by scaling the temperature noise by a factor of 2. This scaling is consistent with the elevated noise levels observed in polarization, illustrated in figure 8 and by the green curves in figure 9.

Bottom panels: *LiteBIRD* high signal-to-noise ratio is evident across all $1/f$ noise levels. Cosmic variance dominates the very low- ℓ regime, and the signal remains robust up to multipoles of approximately $\ell \simeq 1100$ (TT), $\ell \simeq 900$ (EE), and $\ell \simeq 500$ (BB).

5.1 TOD

We show in figure 5 the sum of the TODs of CMB, foregrounds, dipole, white noise and $1/f$ noise with $f_{\text{knee}} = 30$ mHz for three different detectors of LFT and MFT and for the first simulation, i.e. the only one for which we saved TODs to disk. We remind the reader that these TODs were also employed for assessing feasibility of Cosmoglobe framework when applied onto *LiteBIRD* data and results will be presented in [15]. The numbers in the legend are related to the selected detectors in figure 1: the counting starts from 1 for the top-left detector marked with a black star and increases going to the right and going to the following row. The letters T or B denote the bolometer pair, identifying each detector as either *top* or *bottom*. For Q-type antennas, the top/bottom designation corresponds to orientations




Figure 5. Sum of TODs of CMB, foregrounds, dipole, white noise and $1/f$ noise with $f_{\text{knee}} = 30$ mHz for three different detectors and for the first simulation. The numbers in the legend refer to the detectors number as explained in the text, while the T (B) letter refer to the detector being top (bottom). Left panel: L1-040 channel of LFT. Right panel: M1-140 channel of MFT.

of $0^\circ/90^\circ$, while for U-type antennas, it indicates orientations of $45^\circ/135^\circ$. Detectors in the same pair (top and bottom) observe the same signal since they are aligned in the same direction, differing only in their noise component, even though it is difficult to distinguish visually. The detectors shown in figure 5 are selected to illustrate this behavior. When examining a single detector’s time stream, various structures become evident. The long time scale modulation observed in figure 5 corresponds to the dipole signal, while the spikes are related to the samples obtained while scanning across the Galactic midplane (e.g., notice the L1-040 channel is strongly dominated by the Synchrotron signal, as the spikes are more prominent than the rest), small scale fluctuations are instead related to CMB and noise. Furthermore, the difference between the LFT and MFT panels, where the signals appear out of phase, is due to the instruments being oriented in opposite directions (recall figure 2). We show in figure 6 the TOD of all the components separately, as described in section 4 and for a single detector of LFT and MFT.

In figure 7 we show the Power Spectral Density (PSD) of the TOD noise shown in figure 6. As expected, the white noise curve remains nearly constant across the entire frequency range. In contrast, the presence of the $1/f$ noise component causes an increase in noise power at lower frequencies, with a more pronounced effect for $f_{\text{knee}} = 100$ mHz compared to $f_{\text{knee}} = 30$ mHz. The plotted PSD reflects the expected behavior, as shown by the vertical and tilted lines corresponding to the two different knee frequencies of the $1/f$ noise.

5.2 Covariance matrices

We present the inverse of the noise covariance matrix, \mathcal{N}_{pp}^{-1} , in figure 8, specifically for the MFT M1-140 channel. This matrix is constructed from simulated pixel correlations, where each pixel’s covariance is a 3×3 matrix. This is readily observable, beginning with mapmaking equations:

$$\hat{\mathbf{m}} = (\mathbf{A}^T \mathbf{N}^{-1} \mathbf{A})^{-1} \mathbf{A}^T \mathbf{N}^{-1} \mathbf{d}, \quad (5.1)$$




Figure 6. TOD of CMB, foregrounds, dipole, white noise, white noise + $1/f$ noise with $f_{\text{knee}} = 30$ mHz and white noise + $1/f$ noise with $f_{\text{knee}} = 100$ mHz for one different detector and for the first simulation. Left panel: L1-040 channel of LFT. Right panel: M1-140 channel of MFT.




Figure 7. Power Spectral Density of the noise TOD shown in figure 6. The solid lines have been obtained from data, while the dashed-dotted lines show the analytical model (eq. (4.1)). The vertical dashed lines show the two knee frequencies used for the two $1/f$ noise timelines. Left panel: L1-040 channel of LFT. Right panel: M1-140 channel of MFT.

where $\hat{\mathbf{m}}$ is the estimated map, \mathbf{A} is the pointing matrix and contains information about how each pixel in the sky contributes to the time-ordered signal \mathbf{d} collected by one detector, $\mathbf{N} = \mathbf{N}_{tt} = \langle \mathbf{n} \mathbf{n}^T \rangle$ is the noise covariance matrix in time domain and $\mathcal{N}_{pp}^{-1} = (\mathbf{A}^T \mathbf{N}_{tt}^{-1} \mathbf{A})^{-1}$ noise covariance matrix in pixel space. This block encompasses the auto-correlations (TT , QQ , UU) and cross-correlations (TQ , QU , TU , QT , UQ , UT). As the *LiteBIRD* scanning strategy is optimized to have redundancy in the pixel cross-linking, each 3×3 block of the \mathcal{N}_{pp}^{-1} matrix is expected to be invertible and almost diagonal. Any departures from symmetry are mainly attributed to numerical round-off errors. The elements of each block are displayed as six separate maps in figure 8.

In all the matrices, the ecliptic poles and equator are discernible due to lower correlations. This effect is particularly noticeable in the auto-correlation fields, but it can also be




Figure 8. Noise covariance matrices in pixel-space in Galactic coordinates for the channel M1-140 of MFT. Note that the value ranges differ between the maps.

distinguished in the cross-correlation fields. This pattern arises from the scanning strategy collecting more samples at the poles and along the equator. Additionally, the QQ and UU matrices exhibit twice the variance of the TT matrix, and the QU covariances are higher than those in the TQ and TU cases. Specifically, the TQ and TU correlations are expected to be exactly zero by construction since detector pairs are always orthogonal, causing terms like $\text{Cos}(2\alpha) + \text{Cos}(2\alpha + \pi)$ and $\text{Sin}(2\alpha) + \text{Sin}(2\alpha + \pi)$ to cancel. The small observed values ($\sim 10^{-8}$) stem from numerical noise. In contrast, QU does not vanish at the measurement level, as terms like $\text{Sin}(2\alpha) \text{Cos}(2\alpha) + \text{Sin}(2\alpha + \pi) \text{Cos}(2\alpha + \pi)$ are not vanishing by construction. While their mean goes to zero with sufficient observations, finite sampling over one year duration and at a certain resolution, leaves a small residual.

We present a validation test for the covariance matrix, focusing on the impact of the mapmaker on temperature and polarization, demonstrating the HWP's effectiveness in reducing the $1/f$ noise in polarization maps. We compare the binned output maps with the input coadded maps (which include CMB, foregrounds, and $1/f$ and white noise) by taking their difference, as shown in the bottom rows of figure 11. This subtraction isolates the noise

contribution. Next, we compute the full sky power spectrum of the map differences using `healpy`'s `anafast` algorithm. In parallel, we generate 50 temperature and polarization maps as Gaussian realizations with a null mean and variances corresponding to the TT , QQ , and UU diagonal terms of the covariance matrices displayed in figure 8. Then we calculate the spectrum for each realization and compute their average. Figure 9 compares the spectra derived from the map differences with the average spectra from the noise realizations based on the covariance matrix. Specifically, it shows the results for the M1-140 channel of MFT, comparing one simulation for both $f_{\text{knee}} = 30$ mHz and $f_{\text{knee}} = 100$ mHz to the average of 50 noise realizations.

As expected, the noise power spectra obtained from the covariance matrix-based maps (solid green line) are flat across all multipoles, representing uncorrelated white noise. In the temperature noise power spectrum, the difference between binned and input maps (solid blue/orange lines) reveals an increase in power towards lower multipoles, driven by the presence of unmitigated $1/f$ noise. However, despite this enhancement at large scales, the *LiteBIRD* data remain signal-dominated (see figure 4) at low multipoles, making this effect negligible in terms of the overall signal-to-noise ratio. Conversely, for polarization, the difference between output and input (solid blue/orange lines) is consistent with a white noise power spectrum, confirming the effective mitigation of the $1/f$ noise due to the use of a HWP.

5.3 Power spectra

In this section we present a comparison among different power spectra. To calculate the spectra, we utilized `healpy`'s `anafast` algorithm, considering the full sky. Figure 10 displays the power spectra derived from the differences between the output maps and the input coadded maps. The results refer to the average of 500 simulations of the M1-140 channel of MFT. As anticipated, the temperature noise spectrum with $f_{\text{knee}} = 100$ mHz is larger than with $f_{\text{knee}} = 30$ mHz. This outcome aligns with expectations, as higher f_{knee} values correspond to increased noise levels in the simulations. Regarding the polarization case, the HWP demonstrates its effectiveness in reducing $1/f$ noise for both knee frequencies as the noise curves virtually overlap.

5.4 Computational cost

The simulations have been produced at the computing facilities at the MARCONI-CINECA in Italy. The production involved 370 computing nodes, accounting for 48 Intel Xeon 8160 (SkyLake) cores per node. The total computational cost for the simulation production and validation is estimated to be 600 thousand CPU-h. The outputs, described in appendix A, encode 500 realizations of maps with one single realization of TODs occupying a total of about 35 TB of disk space. TODs have been currently stored in two different facilities to enable data retrieval and reproducibility for future studies.

We predict the production of full focal plane simulations within the next few years, with an estimated computational cost of 1 million CPU-h. This estimate is derived by scaling the cost of current simulations, assuming a consistent number of Monte Carlo realizations.




Figure 9. Noise-only power spectra from the difference between binned output and input maps (solid blue/orange) for a single simulation, alongside the average power spectra of white noise from 50 realizations (solid green) based on the noise covariance matrices in figure 8. Top panel: TT . Central panel: EE . Bottom panel: BB . The shown quantities refer to the M1-140 channel of MFT and to both the $1/f$ noise levels ($f_{\text{knee}} = 30$ mHz, $f_{\text{knee}} = 100$ mHz).

6 Conclusions

This paper presents the first release of the official *LiteBIRD* simulations. Through the scanning of input convolved CMB and foreground maps, along with the incorporation of white noise, two types of $1/f$ noise (with $f_{\text{knee}} = 30$ and 100 mHz), and the dipole signal, we simulated one year of Time Ordered Data (TOD) for approximately one-third of *LiteBIRD*'s total detectors. To ensure fidelity, the white noise level was carefully rescaled to match the baseline three-year mission and the entire focal plane. The output comprises 500 binned




Figure 10. Full sky spectra obtained from the difference maps between the binned output maps and the input ones for the channel M1-140 of MFT. The plot shows the average of 500 realizations for both $1/f$ noise levels, with the shaded regions representing the corresponding standard deviations.

maps and a noise covariance matrix for each of *LiteBIRD*'s 22 channels. Additionally, we saved TOD and pointing information for the first simulation, for each of the 22 channels and for each component separately. Moreover, the TODs presented in this work have been used within the Cosmoglobe framework and results will be presented in [15].

Notably, our findings demonstrate that the (ideal) HWP effectively mitigates $1/f$ noise (see figures 9, 10), as evidenced by the virtually flat spectra for polarization.

For the next rounds of simulations, we plan to increase the number of detectors considered, thus producing full focal plane simulations and extend the simulation time to the baseline three-year mission. Moreover, our focus will shift towards incorporating several systematic effects, beyond the existing $1/f$ noise currently employed in this simulation effort. These additional systematics encompass a wide range of considerations, such as gain drifts, downtime occurrences, impacts of cosmic rays, as well as beam and HWP systematics. We aim at describing in a publication in the coming year, the simulations encoding these systematic effects that are being produced at the time of writing this work.

Acknowledgments

LiteBIRD (phase A) activities are supported by the following funding sources: ISAS/JAXA, MEXT, JSPS, KEK (Japan); CSA (Canada); CNES, CNRS, CEA (France); DFG (Germany); ASI, INFN, INAF (Italy); RCN (Norway); MCIN/AEI, CDTI (Spain); SNSA, SRC (Sweden); UKSA (U.K.); and NASA, DOE (U.S.A.).

We acknowledge financial support under the National Recovery and Resilience Plan (NRRP), Mission 4, Component 2, Investment 1.1, Call for tender No. 104 published on 2.2.2022 by the Italian Ministry of University and Research (MUR), funded by the European Union — NextGenerationEU — Project Title “SHIFT” — CUP 55723062008 — Grant Assignment Decree No. 962 adopted on June 30th 2023 by the Italian Ministry of Ministry of University and Research (MUR).

We acknowledge the use of `numpy` [37] and `matplotlib` [38] software packages, and the use of computing facilities at CINECA. Some of the results in this paper have been derived using the `healpy` [13] and `HEALPix` [12] packages.

A Products of the pipeline

We summarize all the pipeline products in table 3. As already stated in section 5.4, for the first realization of simulations, we generated six separate TODs for each channel: `tod_cmb`, `tod_fg`, `tod_wn`, `tod_wn_1f_100mHz`, `tod_wn_1f_30mHz`, and `tod_dip`. Each TOD contains only the specific component indicated by its name. Additionally, a noise covariance matrix in pixel space was produced for each channel, computed only once, as it is independent of the specifics of each simulation (see eq. (5.1)). Finally, we generated a total of 1000 binned maps per channel: 500 coadded maps at the `nside=512` pixel resolution, with $f_{\text{knee}} = 30$ mHz and 500 coadded maps with $f_{\text{knee}} = 100$ mHz. Each coadded map includes contributions from the CMB, foregrounds, $1/f$ noise, and white noise.

Products per channel
tod_cmb
tod_fg
tod_wn
tod_wn_1f_100mHz
tod_wn_1f_30mHz
tod_dip
Covariance
500 coadded maps ($f_{\text{knee}} = 30$ mHz)
500 coadded maps ($f_{\text{knee}} = 100$ mHz)

Table 3. For the 500 simulations, we generated 500 coadded maps, including the CMB, foregrounds (dust, synchrotron, free-free and Galactic CO emission), $1/f$ noise and white noise, by binning the observations for both $1/f$ noise knee frequencies. For the first simulation only, we also saved the TOD components and the noise covariances in pixel space to disk.

B Comparison between input and output maps

Here, we present a comparison between the output and input maps. Figure 11 shows the maps produced by the simulation pipeline alongside the input maps, for the first simulation and the M1-140 channel of MFT. Additionally, the fourth and fifth rows display the differences between the output and input maps. The output maps, when compared to their input counterparts (figure 11) and their differences, do not reveal any notable structures, confirming an accurate scanning of the input maps. The anisotropies observed in the two bottom rows primarily stem from the presence of noise.




Figure 11. Comparison between input and output maps for the M1-140 channel of MFT. Left column: T . Central column: Q . Right column: U . From top to bottom rows: input coadded (CMB + foregrounds); binned output coadded + white noise + $1/f$ noise with $f_{\text{knee}} = 100$ mHz; binned output coadded + white noise + $1/f$ noise with $f_{\text{knee}} = 30$ mHz; binned output (CMB + fg + white noise + $1/f$ noise with $f_{\text{knee}} = 30$ mHz) minus coadded input; binned output (CMB + fg + white noise + $1/f$ noise with $f_{\text{knee}} = 100$ mHz) minus coadded input.

References

- [1] A.H. Guth, *The Inflationary Universe: A Possible Solution to the Horizon and Flatness Problems*, *Phys. Rev. D* **23** (1981) 347 [INSPIRE].
- [2] M. Zaldarriaga and U. Seljak, *An all sky analysis of polarization in the microwave background*, *Phys. Rev. D* **55** (1997) 1830 [astro-ph/9609170] [INSPIRE].
- [3] ACTPOL collaboration, *The Atacama Cosmology Telescope: Two-Season ACTPol Spectra and Parameters*, *JCAP* **06** (2017) 031 [arXiv:1610.02360] [INSPIRE].
- [4] POLARBEAR collaboration, *A Measurement of the Cosmic Microwave Background B-Mode Polarization Power Spectrum at Sub-Degree Scales from 2 years of POLARBEAR Data*, *Astrophys. J.* **848** (2017) 121 [arXiv:1705.02907] [INSPIRE].
- [5] SPT collaboration, *Measurements of B-mode Polarization of the Cosmic Microwave Background from 500 Square Degrees of SPTpol Data*, *Phys. Rev. D* **101** (2020) 122003 [arXiv:1910.05748] [INSPIRE].
- [6] BICEP and KECK collaborations, *Improved Constraints on Primordial Gravitational Waves using Planck, WMAP, and BICEP/Keck Observations through the 2018 Observing Season*, *Phys. Rev. Lett.* **127** (2021) 151301 [arXiv:2110.00483] [INSPIRE].
- [7] WMAP collaboration, *Nine-Year Wilkinson Microwave Anisotropy Probe (WMAP) Observations: Final Maps and Results*, *Astrophys. J. Suppl.* **208** (2013) 20 [arXiv:1212.5225] [INSPIRE].
- [8] PLANCK collaboration, *Planck intermediate results. XXXVIII. E- and B-modes of dust polarization from the magnetized filamentary structure of the interstellar medium*, *Astron. Astrophys.* **586** (2016) A141 [arXiv:1505.02779] [INSPIRE].
- [9] M. Tristram et al., *Improved limits on the tensor-to-scalar ratio using BICEP and Planck data*, *Phys. Rev. D* **105** (2022) 083524 [arXiv:2112.07961] [INSPIRE].
- [10] LITEBIRD collaboration, *Probing Cosmic Inflation with the LiteBIRD Cosmic Microwave Background Polarization Survey*, *PTEP* **2023** (2023) 042F01 [arXiv:2202.02773] [INSPIRE].
- [11] LITEBIRD collaboration, *A Simulation Framework for the LiteBIRD Instruments*, [arXiv:2507.04918] [INSPIRE].
- [12] K.M. Górski et al., *HEALPix — A Framework for high resolution discretization, and fast analysis of data distributed on the sphere*, *Astrophys. J.* **622** (2005) 759 [astro-ph/0409513] [INSPIRE].
- [13] A. Zonca et al., *healpy: equal area pixelization and spherical harmonics transforms for data on the sphere in Python*, *J. Open Source Softw.* **4** (2019) 1298 [INSPIRE].
- [14] COSMOGLOBE collaboration, *COSMOGLOBE: Towards end-to-end CMB cosmological parameter estimation without likelihood approximations*, *Astron. Astrophys.* **678** (2023) A169 [arXiv:2306.15511] [INSPIRE].
- [15] LITEBIRD collaboration, *On the computational feasibility of Bayesian end-to-end analysis of LiteBIRD simulations within Cosmoglobe*, [arXiv:2507.05324] [INSPIRE].
- [16] LITEBIRD collaboration, *Multi-dimensional optimisation of the scanning strategy for the LiteBIRD space mission*, *JCAP* **12** (2024) 036 [arXiv:2408.03040] [INSPIRE].
- [17] PAN-EXPERIMENT GALACTIC SCIENCE GROUP collaboration, *Full-sky Models of Galactic Microwave Emission and Polarization at Sub-arcminute Scales for the Python Sky Model*, [arXiv:2502.20452] [INSPIRE].
- [18] A. Zonca, B. Thorne, N. Krachmalnicoff and J. Borrill, *The Python Sky Model 3 software*, *J. Open Source Softw.* **6** (2021) 3783 [arXiv:2108.01444] [INSPIRE].

- [19] B. Thorne, J. Dunkley, D. Alonso and S. Naess, *The Python Sky Model: software for simulating the Galactic microwave sky*, *Mon. Not. Roy. Astron. Soc.* **469** (2017) 2821 [[arXiv:1608.02841](#)] [[INSPIRE](#)].
- [20] PLANCK collaboration, *Planck 2015 results. X. Diffuse component separation: Foreground maps*, *Astron. Astrophys.* **594** (2016) A10 [[arXiv:1502.01588](#)] [[INSPIRE](#)].
- [21] M. Remazeilles et al., *An improved source-subtracted and destriped 408 MHz all-sky map*, *Mon. Not. Roy. Astron. Soc.* **451** (2015) 4311 [[arXiv:1411.3628](#)] [[INSPIRE](#)].
- [22] J. Delabrouille et al., *The pre-launch Planck Sky Model: a model of sky emission at submillimetre to centimetre wavelengths*, *Astron. Astrophys.* **553** (2013) A96 [[arXiv:1207.3675](#)] [[INSPIRE](#)].
- [23] M.-A. Miville-Deschenes et al., *Separation of anomalous and synchrotron emissions using WMAP polarization data*, *Astron. Astrophys.* **490** (2008) 1093 [[arXiv:0802.3345](#)] [[INSPIRE](#)].
- [24] G. Puglisi, G. Fabbian and C. Baccigalupi, *A 3D model for carbon monoxide molecular line emission as a potential cosmic microwave background polarization contaminant*, *Mon. Not. Roy. Astron. Soc.* **469** (2017) 2982 [[arXiv:1701.07856](#)] [[INSPIRE](#)].
- [25] PLANCK collaboration, *Planck 2013 results. XIII. Galactic CO emission*, *Astron. Astrophys.* **571** (2014) A13 [[arXiv:1303.5073](#)] [[INSPIRE](#)].
- [26] G. Stein et al., *The Websky Extragalactic CMB Simulations*, *JCAP* **10** (2020) 012 [[arXiv:2001.08787](#)] [[INSPIRE](#)].
- [27] M. Birkinshaw, *The Sunyaev-Zel'dovich effect*, *Phys. Rept.* **310** (1999) 97 [[astro-ph/9808050](#)] [[INSPIRE](#)].
- [28] R.A. Sunyaev and Y.B. Zeldovich, *Microwave background radiation as a probe of the contemporary structure and history of the universe*, *Ann. Rev. Astron. Astrophys.* **18** (1980) 537 [[INSPIRE](#)].
- [29] G. Lagache, J.-L. Puget and H. Dole, *Dusty infrared galaxies: Sources of the cosmic infrared background*, *Ann. Rev. Astron. Astrophys.* **43** (2005) 727 [[astro-ph/0507298](#)] [[INSPIRE](#)].
- [30] A. Lewis and A. Challinor, *Weak gravitational lensing of the CMB*, *Phys. Rept.* **429** (2006) 1 [[astro-ph/0601594](#)] [[INSPIRE](#)].
- [31] Z. Li, G. Puglisi, M.S. Madhavacheril and M.A. Alvarez, *Simulated catalogs and maps of radio galaxies at millimeter wavelengths in Websky*, *JCAP* **08** (2022) 029 [[arXiv:2110.15357](#)] [[INSPIRE](#)].
- [32] G. Puglisi et al., *Forecasting the Contribution of Polarized Extragalactic Radio Sources in CMB Observations*, *Astrophys. J.* **858** (2018) 85 [[arXiv:1712.09639](#)] [[INSPIRE](#)].
- [33] G. Lagache et al., *Impact of polarised extragalactic sources on the measurement of CMB B-mode anisotropies*, *Astron. Astrophys.* **642** (2020) A232 [[arXiv:1911.09466](#)] [[INSPIRE](#)].
- [34] PLANCK collaboration, *Planck 2018 results. VI. Cosmological parameters*, *Astron. Astrophys.* **641** (2020) A6 [*Erratum ibid.* **652** (2021) C4] [[arXiv:1807.06209](#)] [[INSPIRE](#)].
- [35] M. Reinecke, S. Belkner and J. Carron, *Improved cosmic microwave background (de-)lensing using general spherical harmonic transforms*, *Astron. Astrophys.* **678** (2023) A165 [[arXiv:2304.10431](#)] [[INSPIRE](#)].
- [36] D.J. Fixsen, *The Temperature of the Cosmic Microwave Background*, *Astrophys. J.* **707** (2009) 916 [[arXiv:0911.1955](#)] [[INSPIRE](#)].
- [37] C.R. Harris et al., *Array programming with NumPy*, *Nature* **585** (2020) 357 [[arXiv:2006.10256](#)] [[INSPIRE](#)].
- [38] J.D. Hunter, *Matplotlib: A 2D Graphics Environment*, *Comput. Sci. Eng.* **9** (2007) 90 [[INSPIRE](#)].